### HUNT FOR COLD H<sub>2</sub> MOLECULES

# Combes, F.<sup>1</sup>

Abstract. The bulk of the molecular component in galaxies is made of cold H<sub>2</sub>, which is not observed directly, but which abundance is derived from indirect tracers such as CO emission. The CO to H<sub>2</sub> conversion ratio remains uncertain, and may vary by large factors in special environments with dierent excitation or metallicity. Recent cold gas discoveries (through -rays or cold dust emission) are reviewed and the most promising tracers in the future are discussed, such as the primordial molecules HD and LiH, or the pure rotational lines of excited H<sub>2</sub>.

#### 1 Introduction

The H<sub>2</sub> m olecule has no electric dipole because of its sym metry, and therefore cannot em it line radiation in the radio dom ain, the only ones that could be excited at the low temperature of the interstellar medium (10-15K). The rst rotational line comes from the J = 2 level, through quadrupole radiation, and is 512K above the ground state, S(0) at 28 m in wavelength. The second rotational line of the para H<sub>2</sub> is S(2) at 12 m, and the two rst lines of the ortho H<sub>2</sub> are S(1) and S(3) at 17 m and 9 m respectively. ISO observations of PDR have revealed a large number of H<sub>2</sub> pure rotational lines (T immerm ann et al 1996), and higher resolution observations have con rm ed an unexpectedly large amounts of warm H<sub>2</sub> gas in PDR (Allers et al 2005).

c Paris Observatory 2024

<sup>&</sup>lt;sup>1</sup> Observatoire de Paris, LERMA, 61 Av. de l'Observatoire, F-75014, Paris

### Hunt for M olecules

The only other way to detect cold  $H_2$  directly is through the UV lines in absorption, unfortunately this concerns only the untypical very di use regions, since higher  $H_2$  column densities are associated to high extinction, that prevent to detect the background sources. This was con med by the FUSE survey towards galactic sources (Shull et al 2004). The average m olecular fraction of the di use ISM is of the order of 10%, and the tem perature is warm (100K). Even in "translucent line of sights", there is only evidence of di use clouds (R achford et al 2002).

We rst review our current know ledge of the CO to H $_2$  conversion ratio, which is still the main method to trace cold H $_2$  molecules in galaxies. A fter presenting possible new tracers, we then describe recent discoveries of cold gas, locally in our Galaxy, and in external galaxies or clusters. We conclude with future experiments, which could bring new insights.

### 2 CO and other tracers

#### 2.1 Uncertainties in the CO to H<sub>2</sub> conversion ratio

A lthough the determ ination of this factor has been the goal of multiple studies, it remains largely uncertain; it can vary by a factor 10 or more with metallicity, and also with excitation and local density. When studied for individual clouds, from virial mass arguments, the conversion factor varies by a factor 10 across the mass spectrum of observed clouds. For extrem e starbursts, it is usual to adopt a conversion ratio 5 times lower than the average, and the value to adopt for high-z or primordial galaxies is unknown.

Since the factor could be large in the outer parts of galaxies, at low m etallicity, it is tempting to wonder how much the dark  $H_2$  gas could account for the galaxy dark matter, in particular the dark baryons (i.e. 90% of them). It is indeed known from microlensing experiments that most of them cannot be in compact objects, and must be in gas, either hot or cold.

The rotation curves here serve as an upper lim it on the conversion ratio. In spiral and dwarf galaxies, rotation curves are remarkably t through H I-scaling by a factor 7-10 (H oekstra et al 2001). So inside the H I disk, i.e. inside the last point of directly m easured rotation curve, it is possible to have only 10-20% of the dark baryons in galaxies. The remaining part m ust reside in cosm ic laments.

24

### 2.2 Other possible tracers

To reveal cold  $H_2$  m olecules, we could think of their hyper ne structure (or ultra ne), coming from the interaction between nuclear spin, and the magnetic eld generated by rotation. The ground state has 3 levels (F = I+ J= 0, 1 and 2) and the corresponding lines are in the kilom etric range: F = 1-0 at 546.4 kHz ( = 0.5km) and F = 2-1 at 54.8 kHz ( = 5.5km). The line intensities are extrem ely weak (cfC om bes & P fenniger 1997), the radiation is stopped by the atm osphere, and their detection would require to cover by a loose grid several kilom eters on the M oon! This also assumes that there is a signi cant am ount ofm olecules in the ortho state, even at low tem per-atures, i.e. the m olecules have no time to reach complete equilibrium after their form ation.

At 3K, the pressure of H<sub>2</sub> clum ps is 100 tim es the saturated vapor pressure (C om bes & P fenniger 1997). How ever, since the latent heat is 110K /H<sub>2</sub>, there is no tim e to form much snow in a Hubble tim e. But the conditions for dim erisation m ight be reached in som e very dense clum ps, and continuum em ission is expected from dim ers, through the dipole induced by collisions (Schaefer 1994, 1999). It is di cult to estim ate the am ount of H <sub>2</sub> at such high densities, and the am ount of dim ers form ed.

The HD m olecule has a weak dipole m om ent =  $5.8 \ 10^{-4}$  D ebye (T re er & Gush 1968). The rst rotational level is 130 K above ground state, and the rst J = 1 0 line is at = 112 m (W right et al 1999, Poleham pton et al 2002). The emission could come only from excited regions, and is quite weak, given that HD/H<sub>2</sub> is of the order 10<sup>-5</sup>. HD is therefore not a pow erfultracer. The LiH m olecule could be a better tracer, since its dipole m om ent is m uch stronger = 5.9 D ebye (Law rence et al 1963), and the rst transition J = 1 0 is only 21 K above ground state, at =  $0.67 \ m$  (450G H z). This line cannot be detected from the ground, because of H<sub>2</sub>O absorption. A ssum ing an abundance of LiH/H<sub>2</sub>  $10^{-10}$ , the line becomes optically thick 1 for N (LiH) =  $10^{12} \ cm^{-2} \ or N(H_2) = 10^{22} \ cm^{-2}$ .

The ion  $H_2^+$  (searched for by Encrenaz & Falgarone, as soon as 1971!) has an expected abundance of 10<sup>-11</sup>-10<sup>-10</sup>. It has an hyper ne structure, but not in the N = 0 state, only N = 1, I= 1, S= 1/2. The energy of the upper level is 110K, among the 5 lines expected, the strongest is at 1343 M Hz. If  $H_2^+$  forms through cosm ic ray ionization of  $H_2$ , it disappears through reaction with  $H_2$  to form the molecular ion  $H_3^+$ , which has a key role in ionm olecule interstellar chem istry (e.g. Geballe 2000). The  $H_3^+$  ion has now

### Hunt for Molecules

been detected in the interstellarm edium (G eballe & O ka 1996, M cC allet al 1999), through absorption lines in the infra-red, via a vibrationnally excited line (since only the asymmetric vibration  $_2$  can induce a dipole moment, in this ion which has no permanent dipole). The deuterated ions are more favorable tracers, since they have a weak dipole,  $H_2D^+$  (372G Hz em ission, Stark et al 1999), or  $D_2H^+$  (692G Hz em ission, Vastel et al 2004).

G iven the weakness of all these tracers, it m ight be interesting to keep tracking the CO m olecule, since traces of C and O are still expected in nearly prim ordial gas. A residual 10<sup>-3</sup> solar abundance is observed in Ly forest clouds, due to the rst stars. If the H<sub>2</sub> gas remains clum py enough, and far from photo-dissociating radiation, it is possible to nd CO m olecules, assuming that some heating sources exist to keep the brightness temperature above the background (e.g. Braine & Herpin 2004). Alternatively, the molecules should be searched in absorption, although with a low surface lling factor of less than 1%.

#### 3 Hints of dark gas

### 3.1 rays

A nother way to trace cold gas in the interstellarm edium is the -rays resulting from the interaction of cosm ic-rays (CR) with all nucleons (through the creation of <sup>0</sup>). The emission of E > 100 Mev -rays is the product of total nucleon density and CR density. The CR are re-accelerated by supernovae and are the products of star form ation, their density decreases radially in the G alaxy. It is possible to calibrate their density in the solar neighbourhood, through mapping of all ISM components (H I, CO, H). Until recently, due to the low spatial resolution, it was possible to attribute part of the -rays to point sources (pulsars), but converging observations now, including extinction (B-V) and cold dust emission in the millim eter range, reveals that -rays are tracing dark gas in the local interstellarm edium (G renier et al 2005). This interesting discovery shows that the dark gas is distributed like an envelope around the CO-traced m olecular gas, cfF igure 1.

## 3.2 High Velocity Clouds

Some of the HVC falling towards the G alactic plane appear compatible with an intergalactic origin, with low metallicity (W akker et al 1999) and high

26



F ig. 1. M ap of the local dark gas, in G alactic coordinates centered on  $\models$  70, as traced from -ray intensity, reddening E (B-V) and 94 G H z em ission, from G renier et al (2005)

deuterium abundance (Sem bach et al 2004). A lthough  $H_2$  molecules have been detected through UV absorption by FUSE in HVC, up to now no CO emission nor cold dust emission had been detected. For the rst time, Miville-Deschênes et al (2005) report dust emission through the comparison of SST (Spitzer Space Telescope) infrared maps with the 21cm HI line emission from the GBT (G reen Bank Telescope). The dust is cooler (10K) than the galactic dust in average, compatible with a large distance from the Galaxy. The column density derived is 5 times that corresponding to the HI, suggesting the existence of dense gas clumps associated to the apparent di use HI gas. The clumps would contain dense molecular hydrogen, and would constitute most of the gas accreted by the Galaxy.

### 3.3 Cooling ows

C old gas has also recently been discovered in the center of rich galaxy clusters, where it was long searched for, as the end product of cooling ows. It has been realized that the ux of cooling gas is in reality about 10 times

### Hunt for Molecules

lower than expected in the case of spherical in ow without feedback. The cold in ow fuels an AGN in the central cD galaxy, and the radio lobes feedback then m oderates the cooling, creating bubbles and shocks (Figure 2). The cold gas has still the dynamics and velocity of the cluster, and is not yet settled in the central galaxy potential well (see Salom e et al 2005, and Salom e, this P roceedings).



F ig. 2. M ap of the center of Perseus cluster: the false color-scale is the X-ray emission from Chandra (Fabian et al 2003), the white contours are the radio emission (Pedlar et al 1990), showing the radio lobes of the central AGN, and the black contours are the CO (2-1) emission, observed with the IRAM -30m (from Salom e et al 2005).

### 4 Perspectives

The dark H<sub>2</sub> clumps are not compact enough to be gravitational lenses for LMC stars, however they could act as gaseous lenses (D raine 1998) or produce scintillation of optical light, through refraction of the background stellar emission (M oniez 2003). In the strong di ractive regime, it m ight be possible to distinguish high contrast uctuations of short time-scales from foreground e ects of light propagation through the atm osphere. A project is undertaken to m onitor extra-galactic stars every 10 seconds, to discover dark gas of  $10^{19}$  m olecules/cm<sup>2</sup> per 10 000 km transverse distance (O SER : 0 ptical Scintillation by Extraterrestrial R effactors, M oniez 2005).

The pure rotational lines of the H<sub>2</sub> m olecule m ight be the best tracer afterall. ISO observations have shown that they are ubiquitous. In outer parts of galaxies they were observed surprisingly high by Valentijn & Van der W erf (1999). In the edge-on galaxy NGC 891, the m olecular column density was derived to be N (H<sub>2</sub>) =  $10^{23}$  cm<sup>-2</sup>, with an excitation temperature of T = 80-90 K, i.e. 5-15 N (H I), enough to explain the rotation curve.

ISO observations of lines of sights through the G alaxy encountering no star form ation regions, but only cold di use gas, have shown a rich variety of H<sub>2</sub> rotational lines. The excitation of these lines in the ground vibrational states cannot be due principally to the UV excitation, but has to come from non-therm al excitation (Falgarone et al 2005a). This suggests that within the cold neutral medium, there always exists a few percent of warm gas, due to the interm ittent dissipation of M HD turbulence, and this warm gas can be used as a tracer of the cold gas, independent of metallicity. The detection of this warm gas, at large scale in the G alaxy, and in a survey of nearby galaxies, in particular at large distance from the center, is the main goal of the project H<sub>2</sub>Ex, a space explorerm ission with a dedicated 2m-class telescope, and a high resolution spectro-im ager in the mid-IR (Falgarone et al 2005b).

The rotational lines of  $H_2$  are the main coolant of the primordial gas, and its collapse to form the rst generation of stars in the universe could be traced by their emission, however orders of magnitude gains in sensitivity are required for future instruments (M izusawa et al 2005).

### References

Allers, K.N., Ja e, D.T., Lacy et al.: 2005, ApJ 630, 368

Hunt for M olecules

Braine J., Herpin F .: 2004 Nature 432, 369 Combes F., Pfenniger D .: 1997, A & A 327, 453 D raine B.T: 1998 ApJ 509, L41 Encrenaz P.J., Falgarone E.: 1971 ApL 8, 187 Fabian A C., Sanders J.S., Allen S.W. et al: 2003, MNRAS 344, L43 Falgarone, E., Verstraete, L., Pineau Des Forêts, G., Hily-Blant, P.: 2005a, A & A 433,997 Falgarone, E., Boulanger, F., Combes, F. et al.: 2005b, EAS Publications Series, Vol14, 2005, pp.57-66 Geballe T R .: 2000, Phil. Trans. Royal Soc. A, 358, 2503-2514 Geballe T R ., O ka T .: 1996, Nature 384, 334 Grenier, IA., Casandijan, J.-M., Terrier, R.: 2005, Science 307, 1292 Hoekstra, H., van Albada, T.S., Sancisi, R.: 2001, MNRAS 323, 453 Law rence T R , Anderson C H , R am sey N F .: 1963, Phys Rev 130, 1865 M cCallB J., Geballe T R., Hinkle K H., Oka T.: 1999, ApJ 522, 338 M iville-D eschênes, M .- A ., B ou langer, F ., R each, W . T ., N oriega-C respo, A .: 2005, ApJ 631, L57 M izusawa, H., Nishi, R., Omukai, K.: 2005, PASJ 56, 487 Moniez M .: 2003 A & A 412, 105 Moniez M: 2005 in "Mass Proles and Shapes of Cosmological Structures", 21st IAP colloquium, EdP Sciences, in press Pedlar A., Ghataure H.S., Davies R.D. et al. 1990, MNRAS 246, 477 Poleham pton, E.T., Baluteau, J.P., Ceccarelli, C., Swinyard, B.M., Caux, E.: 2002 A&A 388, L44 Rachford, B.L., Snow, T.P., Tum linson, J. et al: 2002, ApJ 577, 221 Salom e P., Combes F., Edge A. et al: 2005, A&A Letter, in prep. Schaefer J.: 1994, A & A 284, 1015 Schaefer J.: 1999, in H<sub>2</sub> in Space, Eds.: F.Combes, G.Pineau des Forêts, Cambridge Univ.Press Sem bach, K.R., Wakker, B.P., Tripp, T.M. et al: 2004, ApJS 150, 387 Shull, J.M., Anderson, K., Tum linson, J. et al.: 2004, BAAS 204, 6118 Stark, R., van der Tak, F.S., van Dishoeck, E.F.: 1999, ApJ 521, L67 Timmermann, R., Bertoldi, F., Wright, C.M. et al: 1996, A&A 315, L281 Tre er M ., Gush H P .: 1968, Phys Rev Let 20, 708 Valentin, E.A., van der W erf, P.P.: 1999, ApJ 522, L29 Vastel, C., Phillips, T.G., Yoshida, H.: 2004, ApJ 606, L127 W akker, B.P., Howk, J.C., Savage, B.D. et al: 1999, Nature 402, 388

W right, C M ., van D ishoeck, E F ., Cox, P. et al.: 1999, ApJ 515, L29

30