

A B I M A A R R A Y S U R V E Y O F M O L E C U L E S I N C O M E T S L I N E A R (C / 2 0 0 2 T 7) A N D N E A T (C / 2 0 0 1 Q 4)

Anthony J. Remijan^{1,2,3}, D. N. Friedel⁴, Imke de Pater⁵, M. R. Hogerheijde⁶, L. E. Snyder⁴,
M. F. A'Hearn⁷, Geoffrey A. Blake⁸, H. R. Dickel^{4,9}, J. R. Forster⁵, C. Kraybill⁵, L. W.
Looney⁴, Patrick Palmer¹⁰, & M. C. H. Wright⁵

A B S T R A C T

We present an interferometric search for large molecules, including methanol (CH_3OH), methyl cyanide (CH_3CN), ethyl cyanide ($\text{CH}_3\text{CH}_2\text{CN}$), ethanol ($\text{CH}_3\text{CH}_2\text{OH}$), and methyl formate (CH_3OCHO), in comets LINEAR (C/2002 T7) and NEAT (C/2001 Q4) with the Berkeley-Illinois-Maryland Association

¹ NASA Goddard Space Flight Center, Computational and Information Sciences and Technologies Office, Code 606, Greenbelt, MD 20771

² National Research Council Resident Research Associate

³ Current address: National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22901
email: aremijan@nrao.edu

⁴ Department of Astronomy, 1002 W. Green St., University of Illinois, Urbana IL 61801
email: friedel@astro.uiuc.edu, snyder@astro.uiuc.edu, lw@astro.uiuc.edu

⁵ Department of Astronomy, University of California, Berkeley, CA 94720
email: imke@oris.berkeley.edu, rforster@astro.berkeley.edu, ckraybill@astro.berkeley.edu, wright@astro.berkeley.edu

⁶ Leiden Observatory, P.O. Box 9513, 2300 RA, Leiden, The Netherlands
email: michiel@strw.leidenuniv.nl

⁷ Department of Astronomy, University of Maryland, College Park MD 20742-2421
email: ma@astro.umd.edu

⁸ Division of Geological and Planetary Sciences; Division of Chemistry and Chemical Engineering California Institute of Technology 150-21, Pasadena, CA 91125
email: gab@gps.caltech.edu

⁹ Current address: Department of Physics & Astronomy, University of New Mexico, 800 Yale Blvd. NE, Albuquerque, NM 87131
email: h-dickel@phys.unm.edu

¹⁰ Department of Astronomy and Astrophysics, University of Chicago, Chicago, IL 60637
email: ppalmer@oskar.uchicago.edu

(BIMA) array. In addition, we also searched for transitions of the simpler molecules CS, SiO, HNC, HN^{13}C and ^{13}CO . We detected transitions of CH_3OH and CS around Comet LINEAR and one transition of CH_3OH around Comet NEAT within a synthesized beam of $20''$. We calculated the total column density and production rate of each molecular species using the variable temperature and outflow velocity (VTOV) model described by Friedel et al. (2005). Considering the molecular production rate ratios with respect to water, Comet T7 LINEAR is more similar to Comet Hale-Bopp while Comet Q4 NEAT is more similar to Comet Hyakutake. It is unclear, however, due to such a small sample size, whether there is a clear distinction between a Hale-Bopp and Hyakutake class of comet or whether comets have a continuous range of molecular production rate ratios.

Subject headings: comets: individual (LINEAR (C/2002 T7), NEAT (C/2001 Q4)) – molecular processes – techniques: interferometric – radio lines: solar system

1. INTRODUCTION

Comets are believed to contain the most pristine material remaining from the presolar nebula and are providing important insights into the formation mechanisms of complex molecular species. Comets are primarily located in two distinct regions of the solar system. The Oort cloud, the source of long period ($P > 200$ years) comets, is a spherically symmetric distribution of comets that encompasses the solar system out to a distance of nearly 100,000 AU. The Kuiper-Edgeworth belt that lies in the ecliptic plane just beyond the orbit of Pluto out to several hundred AU is the source of short period ($20 < P < 200$ years) comets (Jewitt 2004). Recent models have suggested that the Oort cloud comets may have had origins in the entire giant planet region between Jupiter and Neptune. These models suggest that comets were formed in a much wider range of nebular environments and probably experienced thermal and collisional processing before they were ejected into the Oort cloud (Weissman 1999). This processing may have "homogenized" the cometary nuclei of Oort cloud comets. Observations comparing long and short period comets, which are believed to have had different origins, will allow us to address the molecular diversity in comets (A'Hearn et al. 1995, Mumma et al. 2005).

Furthermore, a key goal in astrochemistry is to learn whether the molecular diversity seen in high and low mass hot molecular cores (HMCs) is similar to the chemistry of the primordial solar nebula. The molecular diversity in comets may provide the link between the formation mechanisms of complex molecular species seen in high mass HMCs and low mass

young stellar objects to the formation of young solar systems. Therefore, only through comprehensive surveys of HMCs and comets will a consistent, coherent picture emerge regarding the basic chemistry involved in the evolution of HMCs and comets.

Most observations of the molecular composition of cometary gas and dust have taken place in the visible, ultraviolet (see e.g. Hutsemekers et al. 2005 and references therein) or infrared parts of the spectrum (see e.g. Helbert et al. 2005; Delbo Russo et al. 2001). Also, a vast inventory of interstellar ices and volatiles was discovered by satellites passing through the coma of Comet Halley (see e.g. Mitchell et al. 1992). However, there have been several successful detections of the rotational transitions of molecular species, including organic compounds, at millimeter wavelengths (see e.g. Crovisier et al. 2004a,b; Biver et al. 2002). Many of these species are important in prebiotic organic chemistry (e.g. H_2O , HCN , CH_3OH , aldehydes, nitriles) and are believed to be parent molecules (originating from the comet nucleus) rather than the products of photodissociation or gas phase chemistry in the cometary coma. Two prominent examples of parent molecules of prebiotic importance are hydrogen cyanide (HCN) (see e.g. Friedel et al. 2005) and methanol (CH_3OH) (see e.g. Ikeda et al. 2002).

In the Spring of 2004, there was a rare opportunity to observe two dynamically new Oort cloud comets passing into the inner solar system and within 0.3 AU of the Earth: Comet LINEAR (C/2002 T₇) (hereafter, T₇ LINEAR) and Comet NEAT (C/2001 Q₄) (hereafter, Q₄ NEAT). This paper presents the results of an effort to further investigate the astrochemistry of comets by observing several large molecular species in Comets T₇ LINEAR and Q₄ NEAT with the Berkeley-Illinois-Maryland Association (BIMA) array¹¹; these species include methanol (CH_3OH), methyl cyanide (CH_3CN), ethyl cyanide ($\text{CH}_3\text{CH}_2\text{CN}$), ethanol ($\text{CH}_3\text{CH}_2\text{OH}$), and methyl formate (CH_3OCHO). In addition, we also searched for transitions of the simpler molecules CS, SiO, HNC, HN^{13}C and ^{13}CO . We were successful in obtaining single-beam images, cross-correlation spectra, and production rates for cometary methanol (CH_3OH) in both comets and CS in Comet T₇ LINEAR. Upper limits were found for the remaining species in both comets.

¹¹Operated by the University of California, Berkeley, the University of Illinois, and the University of Maryland with support from the National Science Foundation.

2. OBSERVATIONS AND RESULTS

The observations were conducted using the BIMA array near Hat Creek, California¹² in D-con gruration (baselines from 6m to 35m), cross-correlation mode toward Comets T7 LINEAR and Q4 NEAT during their 2004 apparitions.

The observations of Comet T7 LINEAR, using JPL reference orbit 69, were taken 2004 May 11-15 near (J2000) = 01^h40^m, (J2000) = 09 30^{01.3}. The comet was at a heliocentric distance of 0.73-0.77 AU and a geocentric distance of 0.32-0.44 AU ($1^0 = 319$ km at 0.44 AU). W3(OH) was used as the flux density calibrator for these observations. The quasar 0108+015 was used to calibrate the antenna based gains. The observations of Comet Q4 NEAT, using JPL reference orbit 123, were taken 2004 May 20-24 near (J2000) = 09^h07^m, (J2000) = 34 30^{01.1}. The comet was at a heliocentric distance of 0.97 AU and a geocentric distance of 0.55-0.61 AU ($1^0 = 442$ km at 0.61 AU). Mars was used as the flux density calibrator and 0927+390 was used to calibrate the antenna based gains. The absolute amplitude calibration is accurate to within 20%. The channel spacing of all observations was 0.391 MHz, except for CS which had a spacing of 0.098 MHz. The data were reduced, combined, and imaged using the MIRIAD software package (Sault et al. 1995).

Table 1 lists the molecular line parameters of the searched species in Comets T7 LINEAR and Q4 NEAT. The first column lists the molecular species, the second lists the transition, the third lists the transition frequency (MHz), the fourth lists the upper state energy level of the transition (K), the fifth lists the line strength multiplied by the square of the electronic dipole moment (D^2), and the last column lists the quiet sun photodissociation rate for each species (s^{-1}). Table 2 lists the observational parameters. For each comet, the heliocentric distance (r_H), geocentric distance (Δ) in astronomical units (AU), and the synthesized beam size (θ_{syn}) are listed.

Figure 1 (a-c) display the map and spectra of CH₃OH around Comet T7 LINEAR. Figure 1a shows the contour image of the 3_{1,3} - 4_{0,4}A+ transition of CH₃OH starting at 3. Figure 1b shows the cross-correlation spectrum of this transition. The dashed line corresponds to the rest frequency of the 3_{1,3} - 4_{0,4}A+ line for a cometocentric rest velocity of 0 km s⁻¹. The 1 mJy noise level is indicated at the left of the panel. The CH₃OH line for Comet T7 LINEAR was fit with a Gaussian by a least-squares method which gives a peak intensity of 0.44 (0.06) Jy beam⁻¹ and a FWHM of 1.74 (0.31) km s⁻¹. Figure 1c shows the cross-

¹²121°28'00.0 West, 40°49'40.1 North; altitude 1033 m

¹³Both comets moved substantially across the sky during the observations, thus only approximate coordinates are given.

correlation spectrum (Hanning smoothed over three channels) of this transition. Figure 1(d-f) display the map and spectra of CH_3OH around Comet Q4 NEAT. Figure 1d shows the contour image of CH_3OH emission starting at 3 . Figure 1e is the cross-correlation spectrum and the dashed line is similar to Figure 1b for a cometocentric rest velocity of 0 km s^{-1} . The least-squares Gaussian line fit for Comet Q4 NEAT gives a peak intensity of 0.31 (0.13) Jy beam^{-1} and a FWHM of 0.72 (0.3) km s^{-1} . Figure 1f is the Hanning smoothed cross-correlation spectrum.

Figure 2(a-c) display the map and spectra of CS around Comet T7 LINEAR. Figure 2a shows the contour image of the $J = 2 - 1$ transition of CS starting at 2 . Figure 2b is the cross-correlation spectrum of this transition. The least-squares Gaussian line fit for Comet T7 LINEAR gives a peak intensity of 0.21 (0.05) Jy beam^{-1} and a FWHM of 2.45 (0.60) km s^{-1} . Figure 2c is the Hanning smoothed cross-correlation spectrum.

All other molecular transitions in Table 1, including CS in Comet Q4 NEAT, were not detected above 3 . Thus, all column density and production rate upper limits were calculated using the 1 m s noise level of the window containing the molecular line emission and a line width of one channel (1.3 km s^{-1}). Tables 3 and 4 list either the measured intensities (Jy beam^{-1}) and line widths (km s^{-1}) or the upper limits of the intensity and line width for each species in columns 2 and 3. Column 4 lists the total beam-averaged molecular column density (cm^{-2}) and column 5, the production rate (s^{-1}). The calculated production rates were determined using the variable temperature and outflow velocity (VTOV) model (Friedel et al. 2005). Finally, columns 6 and 7 list the production rate ratios with respect to H_2O and CN as will be discussed in §3.2.

3. DISCUSSION

3.1. Column Densities and Production Rates

As described in Friedel et al. (2005), the VTOV model calculates the total column density and production rate of a cometary species assuming optically thin emission in LTE and that the temperature and outflow velocity within the coma vary with cometocentric distance. The VTOV model calculations to determine the total column density and production rate are discussed in detail in Friedel et al. (2005) Appendix A. Here, we discuss the implication of the production rates and column densities. Using the VTOV model, tables 3 and 4 list the total beam-averaged column densities and production rates of each molecular species observed toward Comets T7 LINEAR and Q4 NEAT. The VTOV model was also used to determine the total beam-averaged column density and production rate of CS even though

it is believed to be the daughter species. The parent molecule of CS may be CS₂, which has a very short lifetime of $\sim 10^3$ s or less (Snyder et al. 2001). This short lifetime suggests that CS is formed in the inner coma and the measured line width of CS in Comet LINEAR is similar to HCN and CH₃OH, which are known parent species. Thus, at the spatial resolution of our observations, CS can be treated as though it were a parent species.

3.2. Relative Production Rates of X to H₂O and CN

3.2.1. Comet LINEAR

Based upon the H₂O production rates as measured by Schleicher et al. (2005, private communication) at different heliocentric distances, Friedel et al. (2005) determined the H₂O production rate to be 2×10^{29} s⁻¹ during our observing period. The relative production rates of each molecular species relative to H₂O are listed in column 6 of Table 3. The relative production rate ratios of HCN (Friedel et al. 2005), CH₃OH, and CS with respect to H₂O for Comet LINEAR are most similar to those observed around Comet Hale-Bopp (C/1995 O1) ($2.1 \pm 2.6 \times 10^{-3}$, 4×10^{-2} , and $(3.7 \pm 4.3) \times 10^{-3}$, respectively) (Snyder et al. 2001, Remijan et al. 2004). Schleicher et al. (2005) estimated the CN production rate to be 1.4×10^{26} s⁻¹. We also attempted to determine the relative production rates of each nitrogen bearing molecular species relative to CN. However in this work, no species observed toward Comet LINEAR with a CN bond was detected above the 3 σ detection limit. The upper limits to the production rate ratios are listed in column 7 of Table 3.

3.2.2. Comet NEAT

Schleicher et al. (2005) also determined the H₂O and CN production rates for comet NEAT during our observing period (as reported by Friedel et al. 2005). These are very similar to water production rates measured from Comet Hyakutake at similar heliocentric distances ($\sim 1 \times 10^{29}$ s⁻¹ Lis et al. 1997). The relative production rates of each molecular species relative to H₂O are listed in column 6 of Table 4. The relative production rate ratios of HCN (Friedel et al. 2005) and CH₃OH relative to H₂O are most similar to those measured around Comet Hyakutake (C/1996 B2) (1.2×10^{-3} and 1.7×10^{-2} , respectively) (Biver et al. 1999). The average CN production rate is $Q(\text{CN}) = 2.6 \times 10^{26}$ s⁻¹. The relative production rates of each nitrogen bearing molecular species relative to CN are listed in column 7 of Table 4, however the only species observed toward Comet NEAT with a CN bond was CH₃CN which was not detected above the 3 σ detection limit. The upper limit to the production rate ratio

was $< 4.0 \times 10^{-2}$.

3.3. Classifying Comets T7 LINEAR and Q4 NEAT

Comets are diverse objects, as has been noted by many researchers in the past. For example, A'Hearn et al. (1995) reported an optical study of 85 comets, Biver et al. (2002) reached this conclusion based upon a radio survey of 24 comets, while Mumma et al. (2005) reached the same conclusion from the hypervolatile gases CO and CH₄ from observations at IR wavelengths. Many comet observers have searched for differences between comets from the Oort cloud and the short-period (Jupiter family) comets to attempt to find clear differences in chemical composition depending on their place of formation. As more observations become available, it becomes harder to differentiate between the various comet classes, in particular after the Deep Impact mission (e.g., Mumma et al., 2005).

We can now include Comets T7 LINEAR and Q4 NEAT in the chemical comparison from the detections of both CH₃OH and HCN (Friedele et al. 2005). Both comets T7 LINEAR and Q4 NEAT are believed to be Oort cloud comets given their measured orbital parameters and our observations show that both comets are chemically very similar. However, from the Biver et al. (2002) 30 m survey, there appears to be a distinction between comets rich in HCN (HCN/H₂O > 0.2%) including comets Hale-Bopp (C/1995 O1), 109P/Swift-Tuttle, 1P/Halley and 9P/Tempel 1 (Mumma et al. 2005) compared to comets with an HCN/H₂O abundance ratio < 0.1% which include, among several others, comets Hyakutake (C/1996 B2), Austin (C/1989 X1) and Levy (C/1990 K1). It is unclear, however, due to such a small sample size, whether in fact there are two distinct classes of comets (i.e. Hale-Bopp and Hyakutake class) or if comets have a range of different molecular production rate ratios. Given the abundance ratios of Comet T7 LINEAR, it is more similar to Comet Hale-Bopp that was both HCN and CH₃OH rich. Comet Q4 NEAT on the other hand, has column densities of HCN and CH₃OH more similar to Comet Hyakutake. Furthermore, this distinction appears across both Oort cloud and Kuiper-Edgeworth belt comets as illustrated by recent Deep Impact observations of Comet 9P/Tempel 1 (Mumma et al. 2005). Thus, while our observations still tend to support a chemical distinction between Hale-Bopp and Hyakutake class comets due to the abundance ratios of CH₃OH/H₂O and HCN/H₂O, more observations of both long and short period comets are necessary to increase the sample size. With better statistics, it may be possible to reliably determine the origins of both Oort cloud and Kuiper-Edgeworth belt comets in the pre-solar nebula.

4. SUMMARY

In this paper, we have presented the results of an interferometric search for several large molecules including, methanol (CH_3OH), methyl cyanide (CH_3CN), ethyl cyanide ($\text{CH}_3\text{CH}_2\text{CN}$), ethanol ($\text{CH}_3\text{CH}_2\text{OH}$), and methyl formate (CH_3OCHO) in Comets T7 LINEAR and Q4 NEAT. In addition, we also searched for transitions of the simpler molecules CS, SiO, HNC, HN^{13}C and ^{13}CO . Of the 10 species listed in Table 1, we detected transitions of CH_3OH and CS around Comet T7 LINEAR and one transition of CH_3OH around Comet Q4 NEAT. Using the variable temperature and outflow velocity (VTOV) model described by Friedel et al. (2005), we determined the total beam-averaged column densities and production rates of each molecular species observed toward Comets T7 LINEAR and Q4 NEAT. Based on the molecular production rate ratios with respect to water it appears that Comet T7 LINEAR is more similar to Comet Hale-Bopp while Comet Q4 NEAT is more similar to Comet Hyakutake. However, due to such a small sample size, it is unclear whether there are two distinct classes of comet or if there is a continuous range of classes with numerous and different molecular production rate ratios. More observations with higher sensitivity interferometers such as CARM A or ALMA of both long and short period comets are necessary to increase the sample size, hopefully allowing a determination of the origins of both Oort cloud and Kuiper-Edgeworth belt comets in the pre-solar nebula.

We thank J. R. Dickel for assisting with the observations, an anonymous referee for many helpful comments, and D. G. Schleicher for providing H_2O and CN production rates. This work was partially funded by: NSF AST 02-28953, AST 02-28963, AST 02-28974 and AST 02-28955; and the Universities of Illinois, Maryland, and California, Berkeley.

REFERENCES

- A'Hearn, M. F., Millis, R. L., Schleicher, D. G., Osip, D. J., & Birch, P. V. 1995, *Icarus*, 118, 223.
- Biver, N., et al. 1999, *AJ*, 118, 1850
- Biver, N., Bockelée-Morvan, D., Crovisier, J., Colom, P., Henry, F., Moreno, R., Paubert, G., Despois, D., & Lis, D. C. 2002, *Earth, Moon, and Planets*, 90, 323
- Crovisier, J., 1994, *Journal of Geophysical Research*, 99, 3777
- Crovisier, J., Bockelée-Morvan, D., Colom, P., Biver, N., Despois, D., Lis, D. C., & the Team for target-of-opportunity radio observations of comets 2004a, *A & A*, 418, 1141

- Crovisier, J., Bockelée-Morvan, D., Biver, N., Colom, P., Despois, D., & Lis, D.C. 2004b, *A & A*, 418, L35
- Dello Russo, N., Muma, M.J., DiSanti, M.A., & Magee-Sauer, K. 2001, *BAAS*, 33, 1076
- Friedel, D.N., Remijan, Anthony J., Snyder, L.E., A'Hearn, M.F., Blake, Geoffrey A., de Pater, Imke, Dickel, H.R., Forster, J.R., Hogerheijde, M.R., Kraybill, C., Looney, L.W., Palmer, Patrick, & Wright, M.C.H. 2005, *ApJ*, 630, 623
- Helbert, J., Rauer, H., Boice, D.C., & Huebner, W.F. 2005, *A & A*, 442, 1107
- Hutsemekers, D., Manfroid, J., Jehin, E., Arpigny, C., Cochran, A., Schulz, R., Stuwe, J.A., & Zucconi, J.-M. 2005, *A & A*, 440, L21
- Ikeda, M., Kawaguchi, K., Takakuwa, S., Sakamoto, A., Sunada, K., & Fuse, T. 2002, *A & A*, 390, 363
- Jewitt, D.C. 2004 in *Comets II*, M.C. Festou, H.U. Keller, & H.A. Weaver (eds.), University of Arizona Press, Tucson, p.659–676
- Lis, D.C., Keene, J., Young, K., Phillips, T.G., Bockelée-Morvan, D., Crovisier, J., Schilke, P., Goldsmith, P.F., & Bergin, E.A. 1997, *Icarus*, 130, 355
- Mitchell, D.L., Lin, R.P., Carlson, C.W., Korth, A., Reme, H., & Mendis, D.A. 1992, *Icarus*, 98, 125 Authors:
- Muma, M.J., DiSanti, M.A., dello Russo, N., Magee-Sauer, K., Gibb, E., & Novak, R. 2002, *Proceedings of Asteroids, Comets, Meteors*, Berlin, Germany. Ed. Barbara Warmbein, 753
- Muma, M.J., DiSanti, M.A., Magee-Sauer, K., Bonev, B.P., Villanueva, G.L., Kawakita, H., dello Russo, N., Gibb, E.L., Blake, G.A., Lyke, J.E., Campbell, R.D., Aycock, J., Conrad, A., & Hill, G.M. 2005, *Science*, 310, 270
- Pickett, H.M., Poynter, R.L., Cohen, E.A., Delitsky, M.L., Pearson, J.C., & Muller, H.S.P. 1998, *J. Quant. Spectrosc. & Rad. Transfer*, 60, 883
- Remijan, A., Snyder, L.E., Friedel, D.N., Veal, J.M., Palmer, Patrick, Woodney, L.M., A'Hearn, Michael F., Forster, J.R., Wright, M.C.H., & de Pater, I. 2004, *BAAS*, 36, # 33.02
- Sault, R.J., Teuben, P.J., & Wright, M.C.H. 1995, *ASP Conf. Ser. 77: Astronomical Data Analysis Software and Systems IV*, 77, 433

Snyder, L.E., Veal, J.M., Woodney, L.M., Wright, M.C.H., Palmer, Patrick, A'Hearn,
M.F., Kuan, Y.-J., de Pater, I., & Forster, J.R. 2001, AJ, 121, 1147

Weissman, P.R. 1999, Space Science Reviews, 90, 301

Table 1. Molecular Line Parameters

Species	Quantum Numbers		Frequency (MHz)	E_u (K)	S^2 (D^2)	(1 AU) ^a (s^{-1})	
CH ₃ OH	3 _{1,3}	4 _{0,4} A+	107,013.770 (13)	28.3	3.1	1.3	10 ⁵
CS	2	1	97,980.985 (15)	7.7	7.1	6.7	10 ⁶
¹³ CO	1	0	110,201.353 (1)	5.3	0.01	1.2	10 ^{6b}
CH ₃ CN	6 _K	5 _K	110,383.502 (2) ^c	18.5	91.9	2.9	10 ⁶
CH ₃ CH ₂ CN	10 _{1,10}	9 _{1,9}	86,819.846 (0)	24.1	146.7		10 ⁵
SiO	2	1	86,846.960 (50)	6.3	19.2	6.7	10 ⁶
HN ¹³ C	1	0	87,090.850 (50)	4.2	7.3	1.5	10 ^{5d}
CH ₃ CH ₂ OH	4 _{1,4}	3 _{0,3}	90,117.576 (2)	9.4	5.4	1.8	10 ⁵
CH ₃ OCHO	8 _{0,8}	7 _{0,7} E	90,227.595 (13)	20.1	21.0	4.7	10 ⁵
	8 _{0,8}	7 _{0,7} A	90,229.647 (14)	20.1	21.0	4.7	10 ⁵
HNC	1	0	90,663.572 (4)	4.4	9.3	1.5	10 ⁵

Note. | All molecular line parameters taken from Pickett et al. (1998). Errors in the rest frequency are 2 .

^aTaken or estimated from Crovisier et al. (1994).

^bAssumed to be the same as for CO .

^cOnly the frequency of the $K = 0$ transition is given.

^dAssumed to be the same as for HNC .

Table 2. Observational Parameters

Species	LINEAR				NEAT			
	r_H	(AU)	a	b	r_H	(AU)	a	b
	(AU)		($^{\circ}$)	($^{\circ}$)	(AU)		($^{\circ}$)	($^{\circ}$)
CH ₃ OH	0.75	0.38	22.3	16.9	0.97	0.55	16.7	14.3
CS	0.74	0.41	23.7	18.9	0.97	0.55	18.8	15.4
¹³ CO	0.75	0.38	21.8	16.1	0.97	0.55	16.0	14.1
CH ₃ CN	0.75	0.38	21.6	16.4	0.97	0.55	16.2	14.0
CH ₃ CH ₂ CN	0.77	0.32	28.5	21.3				
SiO	0.77	0.32	28.5	21.3				
HN ¹³ C	0.77	0.32	28.7	20.0				
CH ₃ CH ₂ OH	0.77	0.32	28.2	19.6				
CH ₃ OCHO	0.77	0.32	27.7	19.3				
HNC	0.77	0.32	27.7	20.4				

Table 3. Variable Temperature and Out flow Velocity (VTOV) Results for Comet LINEAR

Species	I_0 (Jy bm^{-1})	v (km s^{-1})	$\ln T_i$ (cm^{-2})	Q (s^{-1})	$\frac{Q(X)}{Q(\text{H}_2\text{O})}$	$\frac{Q(X)}{Q(\text{CN})}^a$
CH ₃ OH	0.44 (0.06)	1.74 (0.31)	1.4 (0.3) 10^{14}	7.5 (1.5) 10^{27}	3.8 (0.8) 10^2	
CS	0.21 (0.05)	2.45 (0.60)	1.4 (0.4) 10^{12}	1.5 (0.5) 10^{27}	7.5 (25) 10^3	
CH ₃ CN	< 0.13	1.06	< 4.6 10^{11}	< 2.3 10^{25}	< 1.2 10^4	< 0.16
¹³ CO	< 0.12	1.06	< 1.5 10^{14}	< 7.7 10^{27}	< 4.0 10^2	
CH ₃ OCHO	< 0.09	1.30	< 2.5 10^{13}	< 2.2 10^{27}	< 1.0 10^2	
HNC	< 0.08	1.35	< 1.8 10^{11}	< 1.4 10^{25}	< 7.5 10^5	< 0.10
HN ¹³ C	< 0.08	1.35	< 2.8 10^{11}	< 2.2 10^{25}	< 1.1 10^4	< 0.16
SiO	< 0.08	1.35	< 2.2 10^{11}	< 1.7 10^{25}	< 8.5 10^5	
CH ₃ CH ₂ CN	< 0.08	1.30	< 2.3 10^{12}	< 1.8 10^{26}	< 9.0 10^4	< 1.3
CH ₃ CH ₂ OH	< 0.08	1.35	< 1.9 10^{13}	< 1.4 10^{27}	< 7.0 10^3	

^aThe ratio is given only for those species which contain a C-N bond.

Table 4. Variable Temperature and Out flow Velocity (VTOV) Results for Comet NEAT

Species	I_0 (Jy bm^{-1})	v (km s^{-1})	$\text{hN}_{\text{T i}}$ (cm^{-2})	Q (s^{-1})	$\frac{Q(\text{X})}{Q(\text{H}_2\text{O})}$	$\frac{Q(\text{X})^a}{Q(\text{CN})}$
CH_3OH	0.31 (0.13)	0.72 (0.30)	4.6 (2.7) 10^{13}	2.9 (1.7) 10^{27}	2.4 (1.4) 10^{-2}	
CS	< 0.14	0.60	< 4.0 10^{11}	< 1.4 10^{25}	< 1.4 10^{-3}	
CH_3CN	< 0.06	1.06	< 2.0 10^{11}	< 1.1 10^{25}	< 9.2 10^{-5}	< 4.0 10^{-2}
^{13}CO	< 0.06	1.06	< 9.8 10^{13}	< 8.3 10^{27}	< 7.0 10^{-2}	

^aThe ratio is given only for those species which contain a C-N bond.

Fig. 1. | Comet LINEAR (C/2002 T7) and NEAT (C/2001 Q4) single field CH_3OH images and spectra. (a) Comet T7 emission contours from the $J = 3_{1,3} - 4_{0,4}A+$ transition of CH_3OH at 107.013 GHz. The contour levels are -0.2, 0.3, 0.4, 0.5, and 0.6 Jy/beam (1 spacing, starting at 3). Image coordinates are arcseconds offsets relative to the predicted position of the nucleus. The synthesized beam is in the lower left, and the line segment shows the solar direction. (b) CH_3OH cross-correlation spectra, Ordinate is flux density per beam, I, in Jy/beam; 0.1 Jy/beam (c) CH_3OH cross-correlation spectra (Hanning smoothed over 3 channels), labels are the same as in (b). (d) Comet Q4 emission contours from CH_3OH . Contours indicate the CH_3OH emission near its peak centered at a cometocentric velocity of 0 km/s. The contour levels are -0.1, 0.15, 0.2, and 0.25 Jy/beam (1 spacing, starting at 3). Image coordinates and labels are the same as in (a). (e) CH_3OH cross-correlation spectra; labels are the same as in (b), 0.05 Jy/beam. (f) CH_3OH cross-correlation spectra (Hanning smoothed over 3 channels), abscissa and ordinate are the same as in (b).

Fig. 2. | Comet LINEAR (C/2002 T7) single field CS image and spectra. (a) Comet T7 emission contours from the $J = 2 - 1$ transition of CS at 97.980 GHz. Contours indicate the CS emission near its peak centered at a cometocentric velocity of 0 km/s. The contour levels are -0.15, 0.15, 0.225, and 0.3 Jy/beam (1 spacing, starting at 2) Image coordinates and labels are the same as in Figure 1. (b) CS cross-correlation spectra; labels are the same as in Figure 1 (b), 0.075 Jy/beam. (c) CS cross-correlation spectra (Hanning smoothed over 3 channels); labels are the same as in (b).




Figure 1.




Figure 2.