Lagrangian Form ulation of a Solution to the Cosm ological Constant Problem

J.W .Mo at

Department of Physics, University of Toronto, Toronto, Ontario, Canada M 5S 1A 7 (April 14, 2024)

Abstract

A covariant Lagrangian formulation of a solution to the cosm obgical constant problem, based on vizualising the uctuations of the vacuum energy as a nonequilibrium process with stochastic behaviour, is presented. The variational principle yields equations of motion for the cosm obgical \constant", treated as a dynamical eld, together with an equation for a Lagrange multiplier eld , and the standard Einstein eld equations with a variable cosm obgical constant term. A stochastic model of yields a natural explanation for the sm allness or zero value of the constant in the present epoch and its large value in an era of in ation in the early universe.

Typeset using REVT_EX

A recent model for solving the cosm obgical constant problem has been proposed [1], in which the vacuum energy is treated as a uctuating environment with stochastic behaviour. In the following, we shall present a covariant formulation based on a Lagrangian density, which yields classical equations incorporating Einstein's gravitational eld equations, upon which a stochastic treatment using a W iener process can be developed.

A lithough the model uses methods of critical phenomena and non-equilibrium statistical mechanics to model the vacuum energy, it can be considered as a phenomenological description of the kinds of behaviour that could be expected in a more fundamental quantum gravity theory. It is generally agreed that the cosmological constant problem cannot be solved within the context of a purely classical theory of gravity. However, no satisfactory quantum gravity theory has been formulated, so it is hoped that our model can shed light on the solution to the problem without the full apparatus of such a theory.

The Lagrangian density is given by

$$L = L_R + L + L_M;$$
(1)

where

$$L_{R} = \frac{p}{gg} R ; \qquad (2a)$$

$$L = 2^{p} - \frac{g}{g} [+(, u + 2)];$$
 (2b)

and = (x) is the variable cosm ological \constant", treated as a dynamical eld, is a Lagrange multiplier eld, u = dx =d is an observer's four-velocity along a world line in spacetime, is a constant and L_M is a matter Lagrangian density. A variation of L with respect to and yields the equations of motion:

$$; u + {}^{2} = 0;$$
 (3a)

Varying L with respect to g and using (3a) gives

$$R = \frac{1}{2}g R + g = 8 GT :$$
 (4)

The kinematical variable u is the four-velocity of a uid element along a world line in spacetime, associated with a uid with density and pressure p, so we do not vary L with respect to u.

The cosm ological constant enters through the vacuum energy density:

$$T_V = _V g = \frac{_V}{8 \ G} g$$
: (5)

Today, has the small value, $< 10^{46} \text{ GeV}^4$, whereas generic in ation models require that has a relatively large value during the in ationary epoch. This is the source of the cosm ological constant problem.

The line element in the Friedmann-Robertson-Walkermodel is

$$d^{2} = dt^{2} R^{2} (t) \frac{dr^{2}}{1 kr^{2}} + r^{2} d^{2} + r^{2} \sin^{2} d^{2}$$
; (6)

where k = 1;0;+1 and we have used comoving coordinates with u = (0;0;0;1). Then, Eqs.(3a), (3b) and (4) become

$$-+\frac{3R}{R}+(2)$$
 1=0; (7b)

H²
$$\frac{R}{R}^2 = \frac{8 G_M}{3} + \frac{1}{3} \frac{k}{R^2};$$
 (7c)

where -= d = dt, M denotes the mass density, H is the Hubble constant, and in the following H₀ and t₀ denote the present values of H and t, respectively. We de ne

tot
$$M + = 1$$
 k; (8)

where $= 8 \text{ G} = 3 \text{H}^2$.

We shall treat the vacuum energy as a uctuating environment and consider as a variable characterizing the state of this system. The parameter in Eq.(7a) corresponds to the dimensioned between the growth and decline of particle-antiparticle annihilation in the vacuum, while the second term is a self-restriction term which limits the growth of .

In ref.(1), we considered the situation in which the vacuum uctuations are rapid compared with $m_{aCTO} = 1$, which de ness the macroscopic scale of time evolution. We assumed that the parameter can be written as t = +t, in which is the average value, t is G aussian noise and measures the intensity of the vacuum uctuations. Let us write Eq.(7a) as

$$d_{t} = (t_{t})^{2} dt + t_{t} dW_{t} = f(t_{t}) + g(t_{t}) dW_{t};$$
(9)

where dW_t is a W iener process. The probability density p(x;t), which describes the distribution, satis as the Fokker-P lanck equation:

$$\mathcal{Q}_{t}p(x;t) = \mathcal{Q}_{x}[(x x^{2})p(x;t)] + \frac{2}{2}\mathcal{Q}_{xx}(x^{2}p(x;t)):$$
(10)

The di usion process is restricted to the positive real half line and 0 and 1 are intrinsic boundaries, because g(0) = 0 and f(1) = 1. The probability of the di usion process reaching in nity as t! 1 is zero, since in nity is a natural boundary. Moreover, zero is a natural boundary if > 2 =2, so neither boundary is accessible and no boundary conditions need be in posed on the Fokker-P lanck equation. For < 2 =2, it can be shown that zero is an attracting boundary.

The stationary-state solution for the probability density, $p_s(x)$, of Eq.(10) is given by [2]

$$p_{s}(x) = N x^{(2 = 2) 2} exp = \frac{2x}{2}$$
 (11)

The norm alization constant N is

$$N^{-1} = \frac{2}{2} \frac{2(2^{-2}) 1 1}{2} \frac{2}{2} \frac{1}{2} \frac{1}{2}$$

where denotes the -function. If p(x;t) is integrable between 0 and 1, then a stationary state solution exists when $> {}^{2}=2$. If it does not exist, then the probability density will be concentrated at zero, i.e., p() = () for $< {}^{2}=2$.

For $0 < 2^{2}=2$, the vacuum uctuations dom in the over the growth or decline of , although the value zero is still the most probable value for , since the distribution function

has a vertical slope at = 0. Because we are using a continuous variable, never reaches the boundary zero in a nite time.

When $> {}^{2}=2$, the growth of dominates the in uence of the vacuum uctuations, and in the neighborhood of zero the probability of = 0 drops to zero. For the stochastic model there are two transition points described by di erent order parameters. At $= {}^{2}=2$ real growth of becomes possible corresponding to a change from a degenerate random variable for steady-state behavior to a stochastic variable; the boundary at = 0 switches from attracting to natural. Secondly, there is the transition point $= {}^{2}$ which corresponds to a qualitative change in the stochastic variable with no change in the nature of the boundary. The probability of = 0 drops abruptly to zero.

The following scenario can be deduced from ourmodel. In the in-ation era, the intensity of vacuum uctuations is large and > 2 , causing a second-order phase transition and a maximum in not near zero. This corresponds to the large vacuum energy needed to drive in ation [3]. As the universe expands the intensity of vacuum uctuations decreases and for $0 < 2^{2}=2$ or $2^{2}=2 < 2^{2}$ the probability density is largest when is non-vanishing and small, which can lead to a current value of $_{0}$ that can be used to t the observational data. If the stationary probability density p_{s} does not exist for $2^{2}=2$, then = 0 is a stationary point; the drift and di usion vanish simultaneously for = 0 and p() = (). This corresponds to the case when is vanishingly small.

Thus, our model provides a natural explanation, in terms of non-equilibrium stochastic processes in an expanding universe, for the behavior of required to t observational data and still be consistent with in ationary models.

O ngoing searches [4] for Type Ia supernovae show that < 0.47 (at 95% condence level for spatially at models). Moreover, for $\neq 0$ a larger fraction of Q SO s would be gravitationally lensed and Q SO surveys give 0.7 [5]. Cold dark matter models (CDM) for large scale stucture formation, which include a cosmological constant, yield a better t to the shape of the observed spectrum of galaxy clustering than does the standard $_{\rm M} = 1$ CDM models, using h $H_0 = (100 \text{km} = \sec M \text{ pc}) = 0.7$, = 0.6 and a baryon density with

5

 $_{\rm B}$ = 0:0255, consistent with prim ordial nucleosynthesis [6]. However, the am plitude for the CDM models is too high compared to the data, a problem that persists at all scales.

The problem of the age of the universe is also alleviated in models. An analysis of the cosm ological data showed that for $= 0.65 \quad 0.1;_{M} = 1$ and a small tilt: 0.8 < n < 1.2, models exist which are consistent with the available data and an in ationary spatially at universe [7].

M odels based on treated as a scalar dynam ical eld [8] have been found to partially resolve observational problem s. The observations of gravitationally lensed Q SO s yield a less restrictive upper bound on H_0t_0 in these m odels [9]. They may also provide a solution to the size of the am plitude problem, since although the shape of the spectrum is the sam e as that of the CDM m odel with = 0.6, the dynam ical m odel yields a lower am plitude and therefore gives a better t to the galaxy clustering data [9].

ACKNOW LEDGM ENTS

I thank M.A.C layton for helpful discussions. This work was supported by the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- [1] J.W .M o at, astro-ph/9606071 (1996).
- [2] W. Horsthem ke and R. Lefever, Noise-Induced Transitions, Theory and Applications in Physics, Chemistry, and Biology, Springer-Verlag, Berlin, Heidelberg, New York, 1984.
- [3] A.D.Linde, Particle Physics and In ationary Cosmology, Gordon and Breach, New York, 1990.
- [4] S. Perlm utter et al., astro-ph/9602122 (1996).
- [5] C.S.Kochanek, astro-ph/9510077, A strophys. J., in press (1996); A strophys. J. 419, 12 (1993).
- [6] J.A. Peacock and S.J. Dodds, MNRAS, 267, 1020 (1994).
- [7] J.P.O stricker and P.J.Steinhardt, Nature 377, 600 (1995); L.K rauss and M.S.Turner, Gen.Rel.Grav. 27, 1137 (1995).
- [8] J.W. Mo at, Phys. Lett. B 357, 526 (1995). This article contains further references to dynamical scalar models. See also, J.A. Friem an et al., Phys. Rev. Lett. 75, 2077 (1995).
- [9] K. Coble, S. Dodelson, and J. A. Friem an, astro-ph/9608122 (1996).