

# Resolving the Structure of Cold Dark Matter Halos

Moore B.<sup>1</sup>, Governato, F.<sup>1</sup>, Quinn T.<sup>2</sup>, Stadel J.<sup>2</sup> & Lake G.<sup>2</sup>

<sup>1</sup> Department of Physics, University of Durham, Durham City, DH1 3LE, UK

<sup>2</sup> Department of Astronomy, University of Washington, Seattle, WA 98195, USA

## ABSTRACT

We examine the effects of mass resolution and force softening on the density profiles of cold dark matter halos that form within cosmological N-body simulations. As we increase the mass and force resolution, we resolve progenitor halos that collapse at higher redshifts and have very high densities. At our highest resolution we have nearly 3 million particles within the virial radius, several orders of magnitude more than previously used and we can resolve more than one thousand surviving dark matter halos within this single virialised system. The halo profiles become steeper in the central regions and we may not have achieved convergence to a unique slope within the inner 10% of the virialised region. Results from two very high resolution halo simulations yield steep inner density profiles,  $\rho(r) \sim r^{-1.4}$ . The abundance and properties of arcs formed within this potential will be different from calculations based on lower resolution simulations. The kinematics of disks within such a steep potential may prove problematic for the CDM model when compared with the observed properties of halos on galactic scales.

*Subject headings:* cosmology: theory - dark matter - galaxies: halos - clusters - methods: numerical

## 1. Introduction

The cold dark matter (CDM) model is a highly successful and well motivated cosmological model. The basic premises are an inflationary universe dominated by a dark matter particle, such as the axion, that leads to “bottom up” hierarchical structure formation. Small dense halos collapse at high redshifts and merge successively into the large virialised systems that contain the galaxies that we observe today (*e.g.* Davis *et al.* 1985 and references therein). On scales  $\gtrsim$  several Mpc, variants from the standard model can successfully reproduce the observed clustering pattern of galaxies together with the abundances of clusters – two fundamental statistics that are sensitive to both the shape and amplitude of the power spectrum (Eke *et al.* 1996). Comparisons between the model and data on smaller scales test a different aspect of the CDM paradigm. For example, the internal structure of halos can be compared with the dynamics of dwarf galaxies. This test shows that the dark matter is not as concentrated as numerical simulations of CDM indicate, perhaps indicating that the dark matter is not as “cold” as CDM (Moore 1996, Flores *et al.* 1996).

The properties of dark matter halos in the CDM model have been extensively investigated by numerous researchers since the early 80’s, however, only in the past decade have computational facilities and software improved to the point such that the central properties of dark halos can be compared directly with the observations. The first results of Zurek *et al.* (1986) and Frenk *et al.* (1988) did not have the resolution necessary to probe the “observed” region of dark halos, but did show isothermal structures in the outer regions. Higher resolution simulations by Dubinski & Carlberg (1991), Warren *et al.* (1992), Carlberg (1994), Crone *et al.* (1994) showed evidence for density profiles with slopes flatter than isothermal in the central regions, with densities varying as  $r^{-1}$  in the inner 10% of the halos. Navarro, Frenk & White (1996, hereafter NFW) made a systematic study of CDM halo structure over a range of mass scales. They found that the density profiles of halos follow a universal form, uniquely determined by their mass and virial radius; varying from  $r^{-1}$  in the central regions, smoothly rolling over to  $r^{-3}$  at the virial radii. (The virial radius,  $r_{vir}$ , is defined as the radius of a sphere containing a mean mass over-density of 200 with respect to the global value). The NFW halos typically contained 5,000 – 10,000 particles, a number that was claimed to be sufficient to resolve the density profile of halos beyond a distance  $\sim 1\%$  of the virial radius.

In this paper we investigate the inner structure of CDM halos using N-body simulations with much higher mass and force resolution than used previously. We shall test separately the effects of softening and particle number on the final density profiles and the issue of convergence to a unique profile. Similar tests were performed by Tormen *et al.* (1996) and Craig (1997). Although the scale of their tests were smaller than carried out here, their results suggested a similar dependence on numerical resolution. We begin with a brief discussion of the numerical techniques and simulation parameters. In Section 3 we show visually and graphically how the structure and substructure of dark halos changes with numerical resolution and we attempt to separate the affects of force softening and mass resolution on the density profiles. Finally, we discuss the role of resolution in

determining halo density profiles and the possible origin of the steep inner density cusp that we find.

## 2. Techniques and initial conditions

Simulating the formation of a dark matter halo requires an accurate treatment of the cosmological tidal field from a large volume of the universe, whilst obtaining high resolution throughout the infall and turnaround regions that will constitute the final halo. A candidate halo is initially identified from a large cosmological volume that has been simulated at lower resolution. The particles within the selected halo are traced back to the initial conditions to identify the region that will be re-simulated at higher resolution. The power spectrum is extrapolated down to smaller scales, matched at the boundaries such that both the power and waves of the new density field are identical in the region of overlap, then this region is populated with a new subset of less massive particles. Beyond the high resolution region the mass resolution is decreased in a series of shells such that the external tidal field is modelled correctly in a cosmological context. The starting redshift is increased such that the initial fluctuations are less than one percent of the mean density and we then re-run the simulation to the present epoch.

The particle distribution is evolved using a new high performance parallel treecode, “PKDGRAV”, that has accurate periodic boundaries (previous studies using treecodes had vacuum boundary conditions), and a variable timestep criteria based upon the local acceleration. The code uses a co-moving spline softening length such that the force is completely Newtonian at twice our quoted softening lengths. In terms of where the force is 50% of the Newtonian force, the equivalent Plummer softening length would be 0.67 times the spline softening length. We resimulate two halos, both extracted from standard CDM simulations normalised such that  $\sigma_8 = 0.7$  and the shape parameter  $\Gamma = 0.5$  ( $H=50\text{km s}^{-1} \text{ Mpc}^{-1}$  is adopted throughout). The “Virgo” halo has a virial radius of 2 Mpc and was extracted from an initial simulation with a box length of 100 Mpc. The “Coma” halo has a virial radius of 3.4 Mpc and was extracted from a larger initial simulation that had a volume of one  $\text{Gpc}^3$ . We performed several simulations of each halo; at our highest resolution the particle mass in the highest resolution regions was  $8 \times 10^8 M_\odot$  and we used a 5 kpc softening length.

PKDGRAV has been extensively tested using a variety of methods, for example, comparing forces with those calculated using a direct  $N^2$  code on a clustered particle distribution or comparing the growth of waves against linear predictions. We have made one additional test by simulating a halo with PKDGRAV and a  $P^3M$  code (Couchman 1991). We used the “cluster comparison” data generated by (Frenk *et al.* 1998). This is constrained realization of a standard  $\Omega = 1$  cold dark matter universe such that a rich cluster forms at the center of a 64 Mpc box. The cluster contains over  $10^5$  particles within  $r_{vir}$  at the final time and both simulations use a

force softening with a resolution of  $0.5\%r_{vir}$ . The global evolution was virtually identical in each simulation; the final virial radii are within 0.3% of each other and the density profiles agreed extremely well on all scales.

### 3. Results

Colour Plate 1 shows the final density distribution of our highest resolution simulation of the “Coma” cluster. The colour scale in both panels represents the local over-density at the position of each particle and reflects a variation from  $10^1$  to  $10^6$  times the mean density. The upper panel shows the entire 1000 Mpc box and illustrates the nested refinement zones, although plotted on this scale the central cluster is not resolved. In the lower panel we have extracted a sphere of radius  $3.4 \text{ Mpc} = r_{vir}$  around the cluster. A wealth of substructure and “halos within halos” can be seen within this cluster.

The left and right hand panel of Colour Plate 2 shows the physical densities and the phase space densities of the Coma halo simulated at three different resolutions. These quantities are plotted at the position of each particle using a colour scale spanning 5 orders of magnitude and they are measured by smoothing over the nearest 64 particles. The upper plot shows the cluster simulated at a resolution similar to that used by NFW. There are about 13,000 particles within the virial region. The central and lower plots show the same cluster simulated at higher resolution such that there are  $1 \times 10^5$  and  $2.7 \times 10^6$  particles respectively within the virial radius. The visual difference is remarkable. As we increase the mass resolution we begin to resolve more and more substructure; at the highest resolution there are  $\sim 1500$  halos within  $r_{vir}$ . A detailed analysis of the properties and dynamics of the dark matter substructure within the “Virgo” cluster simulation can be found in Ghigna *et al.* (1998).<sup>1</sup>

In Figure 1 we show the density profile (calculated in spherical shells) of the “Coma” cluster simulated with different mass and force resolutions. The halo centers are identified using both the center of mass of the inner halo and using the most bound particle and both methods give very similar results. We find that at higher resolutions, the central halo density profiles becomes steeper and as we increase the mass resolution by a factor of 200, the density at 1% of the virial radius increases by 250%.

We have performed 5 simulations of this cluster in which we systematically increased the mass resolution such that we have a total of 92, 1450, 13000, 100000 and 2700000 particles within  $r_{vir}$ . In each simulation we set the softening parameter to be 1/50'th of the mean inter-particle

---

<sup>1</sup>We note that the inner 10% of even the highest resolution cluster is entirely smooth. The overmerging problem is still inherent in dark matter simulations as a result of halo-halo collisions and the global tidal field destroying halos that pass close to the center (Moore, Katz & Lake 1996).

separation in the highest resolution region; in the highest resolution run, the softening was equal to 5 kpc. For each simulation we determine the radius at which the density profile falls by 20% with respect to the highest resolution simulation and define this to be a “reliable” radius beyond which we can trust the results. We find that this radius decreases systematically from 13%, 4.5%, 2.8%, 1.9% of  $r_{vir}$  and the total number of particles that lie within these radii are 9, 35, 160, 700 respectively. We conclude from this experiment that setting a minimum particle number is an insufficient criterion for defining the radius outside of which to “believe” the density profile. In fact, the reliable radius defined in this way is very close to a scale equal to the 0.5 times the mean inter-particle separation of those particles within  $r_{vir}$ .

These convergence tests show that a factor of 8 in mass resolution leads to a factor of 4 increase in the number of particles needed to define the reliable radius. We therefore estimate that the highest resolution run is correct to a radius that contains about 5600 particles which is 0.8% of  $r_{vir}$ . At this point the slope of the density profile has converged to a value of  $r^{-1.4}$ . It is possible that further increases in resolution will trace out this asymptotic gradient to higher densities but not to significantly steeper slopes.

### 3.1. Force softening and particle number

We are attempting to model a collisionless system by sampling phase space with a small number of particles. This sampling introduces numerical noise that is ameliorated by softening the gravitational force on small scales that can effect the halo properties. The number of particles defines the smallest halo that can collapse at high redshifts and thus introduces a maximum phase space density that can be resolved. The effect of softening is to introduce a “soft” core at the centers of all the halos approximately the size of the softening length. This causes halos to be disrupted more easily from tidal forces – the mechanism responsible for the over-merging problem (Moore, Katz & Lake 1996).

At a fixed mass resolution, decreasing the softening will lead to higher central densities, but there will be a point at which smaller softening lengths will not change the inner profile. At this point the density profile turns over to a constant value that is set by the maximum density that can be resolved by the first collapsing halos – unless artificial relaxation occurs. Figure 2 shows the effects of softening on the “Virgo” cluster profile. This halo has  $\sim 20,000$  particles within the virial radius of 2 Mpc and we simulate it using values of the softening length ranging from 1 kpc to 1 Mpc. The density profiles are plotted on scales smaller than the adopted softening lengths to indicate the behaviour in the inner regions, but results on scales less than the softening length have no physical meaning. Having said this, we note how well the density profile agrees on all scales slightly larger than the softening length. As the softening length is decreased we converge upon the same profile found by NFW, with a slope tending towards  $r^{-1}$  in the inner regions.

Eventually we run out of particles to resolve the inner regions and decreasing the softening from 20 kpc to 10 kpc has very little effect on the halo profile.<sup>2</sup>

The solid lines in Figure 2 show the same cluster simulated with 10 kpc softening, but with 20 times as many particles. This leads to a significantly steeper density profile than the same halo simulated with the same softening but using fewer particles. At a radius equal to 1% of the virial radius, just over two softening lengths, the density increases by 70%. Within this radius the low resolution run contains over 50 particles. The two solid lines in Figure 2 show the density profiles of the same halo at redshifts  $z=0$  and  $z=0.25$ , several billion years before the final output. This halo contains almost half a million particles within the virial radius and over 90% of the mass is in place at a redshift  $z=0.25$ . The density profiles normalised to  $r_{vir}$  are almost identical at both epochs, demonstrating that relaxation at late times or other numerical artifacts are not affecting our results even in the central regions.

### 3.2. Fundamental scatter in the density profiles

Although a great deal of substructure is evident in these simulations, the bulk of the mass distribution in these halos lies within a smooth background of particles that were tidally stripped from the infalling halos. A single subclump with velocity dispersion  $\sigma_{sub}$  will introduce a fluctuation in the density profile  $\delta\rho/\rho \sim \delta M_{sub}/\delta M_{clus} \sim (\sigma_{sub}/\sigma_{clus})^2$ . Therefore, even the largest clumps of substructure with circular velocity  $\sim 300 \text{ km s}^{-1}$  will only produce fluctuations of order 5% in a rich, virialised, galaxy cluster. The global profiles may also vary from cluster to cluster, perhaps as a result of different formation histories.

The residuals from the best fit NFW profiles to these halos show a characteristic “S” shape with deviations of order 20%. Within 10% of the virial radius a power law of slope ( $\sim r^{-1.4}$ ) is an excellent fit. A simple modification to the NFW profile from  $\rho(R) \propto (R(1 + R^2))^{-1}$  to  $\rho(R) \propto (R^{1.4}(1 + R^{1.4}))^{-1}$  fits our high resolution simulations very well. (Here  $R$  is the radius from the cluster center expressed in terms of the virial radius divided by a scale factor that is equal to  $r_{vir} \times 0.18$  for the halos simulated here). We emphasize the need to perform many more halo simulations at this resolution to study the scatter in the inner and outer slopes and the concentration parameter.

---

<sup>2</sup>Although not plotted here, values of the co-moving softening below 5 kpc lead to visible signs of relaxation in the density profile - the central region begins to expand. This evidence for collisionality occurs for values of the softening smaller than 1/200'th of the mean inter-particle separation within the cosmological volume simulated and less than 1/20'th the mean inter-particle separation within the virial radius.

#### 4. Achieving convergence in halo properties – what’s going on?

What are the possible numerical or physical reasons behind the shape of CDM density profiles and the dependence upon resolution that we have investigated here?

The simplest interpretation is that the gravitational softening is affecting our results on scales less than about 4 or 5 softening lengths; all of the halo density profiles we have calculated agree very well beyond this scale. Thus, at our highest resolution we have converged upon a unique solution and we can resolve the density profiles to  $\sim 1\%$  of  $R_{vir}$  with  $10^6$  particles and softening  $0.2\%$   $R_{vir}$ . However, we have demonstrated that changing the softening whilst keeping the mass resolution fixed leads to agreement between profiles on scales just larger than one softening length.

Splinter *et al.* (1997 and references within) emphasize that many statistics are not accurately reproduced using N-body codes that are intrinsically collisional. In comparison with PM codes, they argue that only scales larger than the mean inter-particle separation should be considered. This would correspond to a scale about  $10\%$   $r_{vir}$  within our highest resolution simulation - an order of magnitude larger than our convergence tests suggest.

Evans & Colett (1997) demonstrated that a density profile with a slope of  $-4/3$  is a stable solution to the Fokker-Planck and the collisional Boltzmann equations. However, we note that the trend in Figure 1 is the opposite as expected from collisional effects. Furthermore, we do not observe a significant inwards energy transfer as expected from the Evans & Colett model.

The inner density profiles may hold clues as to the initial power spectrum. For example, Subramanian & Ostriker (1998) extended Bertschinger’s (1985) self-similar solutions to the spherical collapse model to include angular-momentum. They find that the slope of the profiles are directly related to the degree of translational motion and the spectrum of density fluctuations.

Steeper profiles may arise from substructure halos that are transferring material at high phase space density to the center of the cluster (Syer & White 1997). As we move to higher resolution we can resolve structure collapsing at earlier redshifts. This material at the highest physical density, invariably ends up at the center of the cluster by  $z = 0$ . Because it is denser it is more robust to tidal disruption by the cluster, therefore has the potential of carrying more mass to the central regions.

We demonstrate this latter effect by tracing back the particles that lie within the final virial radius to their positions at  $z=5$ . Figure 3 shows the fractional mass of the cluster within a given overdensity contributed by the halos that have collapsed at  $z=5$ . In the high resolution simulation 14% of the total mass of the final halo has already collapsed within halos at  $z=5$ . This material ends up constituting 75% of the mass within  $0.01r_{vir}$ . Within our medium resolution simulation of the same halo we find that only 2% of the mass of the final halo has collapsed by  $z=5$  and this ends up making just 20% of the halo mass within  $0.01r_{vir}$ . The low resolution simulation does not contain any virialised halos at  $z = 5$ .

The power spectrum of CDM type models asymptote towards a slope  $n = -3$  and provides power on all scales relevant to cosmological numerical simulations. Therefore, most of the material in the CDM universe and must lie in collapsed virialised clumps at high redshifts. If the simulations had even better mass resolution, we would have expected to observe halos collapsing at much earlier epochs and at higher phase space densities. This material would end up at the center of the final cluster, perhaps altering the density profiles from what we find at the current resolution.

## 5. Summary

We have performed the highest resolution simulations of cold dark matter halos to date. Our force resolution is 0.2% of the virial radii and mass resolution is such that we can resolve halos 1/50,000th of the mass of the final system. The wealth of substructure within the final systems is phenomenal and will be the focus of a future paper. We can identify approximately 1500 “halos within halo” at the final time. Some of the halos within the cluster’s virial radius contain their own substructure.

Both particle number and softening play a role in shaping the final density profiles; as we increase the resolution the halo profiles become steeper and denser in the centers. With the same force softening, but with 10 times as many particles, an individual halo will be roughly twice as dense at  $0.01r_{vir}$  and increasing the mass resolution by another order of magnitude leads to densities that are a further 50% higher at the same radius. Even with 3 million particles per halo we might not have converged upon a unique density profile for cold dark matter halos. Of order  $10^6$  halo particles must be used in order to compare the dynamics of the inner few percent of the cluster mass distribution with observations.

Our convergence tests suggest that our highest resolution simulation can be used to determine the density profile to a scale just less than  $\sim 1\%$  of the virial radius. At this point the profile has reached an asymptotic slope of  $r^{-1.4}$ . It will be interesting to study the statistics of gravitational arcs within clusters using steeper density profiles (*c.f.* Bartelmann 1996). Since galaxy halos will have steeper profiles at a fixed scale length compared to the cluster sized halos simulated here, the problem of reconciling cold dark matter halos with observations of galaxy rotation curves will be considerably exacerbated by these results.

**Acknowledgments** We would like to thank John Dubinski, Gus Evrard, Carlos Frenk, Neal Katz, Adrian Melott, Julio Navarro and Simon White for stimulating discussions on halo formation. We also thank the referee for comments that helped clarify the paper. Adrian Jenkins kindly providing the IC’s for the cluster comparison run. This research required a great deal of computational resources and we are indebted to NASA’s High Performance Computing and



Communications program and the Virgo consortium. Computations were performed on the Pittsburgh T3E and the UK CCC Origin 2000. BM is a Royal Society University Research Fellow, FG is supported by the European Network grant “Galaxy Formation and Evolution”.

## References

- Bartelmann M. 1996, *A.A.*, **313**, 697.
- Bertschinger E. 1985, *Ap.J.Supp.*, **58**, 39.
- Carlberg R. 1994, *Ap.J.*, **433**, 468.
- Couchman H.M.P., 1991, *Ap.J.Lett.*, **368**, 23L.
- Craig M. 1997, Ph.D. thesis, University of California, Berkeley.
- Crone M.M., Evrard A.E. & Richstone D.O. 1994, *Ap.J.*, **434**, 402.
- Davis M., Efstathiou G., Frenk C.S. & White S.D.M. 1985, *Ap.J.*, **292**, 371.
- Dubinski J. & Carlberg R. 1991, *Ap.J.*, **378**, 496.
- Eke V.R., Cole S. & Frenk C.S. 1996, *M.N.R.A.S.*, **282**, 263.
- Evans W.N. & Collet 1997, *Ap.J.Lett.*, in press.
- Flores R.A. & Primack J.R. 1994, *Ap.J.Lett.*, **457**, L5.
- Frenk C.S., White S.D.M., Davis M. & Efstathiou G. 1988, *Ap.J.*, **327**, 507.
- Frenk C.S., White S.D.M., Jenkins A.R., Pierce F. & Evrard A. 1998, in preparation.
- Ghigna S., Moore B., Governato F., Lake G., Quinn T. & Stadel J. 1998, *M.N.R.A.S.*, submitted.
- Moore B., 1994, *Nature*, **370**, 620.
- Moore B., Katz N. & Lake G. 1996, *Ap.J.*, **457**, 455.
- Navarro J.F., Frenk C.S. & White S.D.M. 1996, *Ap.J.*, **462**, 563.
- Quinn P.J., Salmon J.K. & Zurek W.H. 1986, *Nature*, **322**, 329.
- Subramanian K. & Ostriker J. in preparation
- Splinter R.J., Melott A.L. & Shandarin S.F. 1998, *Ap.J.*, in press.
- Syer D. & White S.D.M. 1997, *Astro-ph*, 9611065
- Tormen G., Bouchet F.R. and White S.D.M. 1996, *M.N.R.A.S.*, **286**, 865.
- Warren S.W., Quinn P.J., Salmon J.K. & Zurek H.W. 1992, *Ap.J.*, **399**, 405.

### Figure Captions


**Colour Plate 1.** The upper panel shows the particle distribution in the full 1000 Mpc box from the resimulation of the Coma cluster. This highlights the scale of the simulation and the seven refinement zones that focus in on the highest resolution region. In the lower panel we show a sphere of radius  $r_{vir}$  that contains the cluster at  $z=0$ . All the visible halos and substructure in this plot lie within the virial radius and the colour scale spans a factor of  $10^5$  in local density.

**Colour Plate 2.** We show the projected particle distributions for the Coma cluster at three different resolutions. In each case, we have extracted a sphere of radius equal to  $r_{vir} = 3.4$  Mpc. On the left hand side we plot the logarithm of the local density covering 5 orders of magnitude, from  $\rho/\rho_o = 10^1 - 10^6$ . On the right hand side we plot the local phase space density,  $\rho/\sigma^3$ , where  $\sigma$  is the local velocity dispersion calculated by smoothing over the nearest 64 neighbours.

**Figure 1.** The density profiles of the Coma cluster simulated at four different resolutions. The curves begin at the spline softening lengths that were used and the number of particles within the final virial radii are indicated.


**Figure 2.** The broken lines show the Virgo halo simulated at the same mass resolution but varying only the softening parameter. This halo has a virial radius of 2 Mpc and contains 20,000 particles within  $r_{vir}$ . The values of the softening used are indicated next to each curve. The solid curves show the same cluster resimulated with a mass resolution 20 times higher, but keeping the force softening fixed at 10kpc. To demonstrate that relaxation is not affecting our results one of the solid curves shows the profile at a redshift  $z=0.25$ .

**Figure 3.** We plot the mass fraction of the final halo that resides in material identified at  $z=5$  with  $\rho/\rho_o > 200$ , against the overdensity that that material lies at  $z=0$ . The solid curve denotes our highest resolution simulation of the Coma cluster and the dashed curve denotes the intermediate resolution simulation. (The low resolution run did not contain any virialised halos at  $z=5$ .)



This figure "plate1\_lores.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/astro-ph/9709051v2>



This figure "plate2\_lores.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/astro-ph/9709051v2>

