

## WHY IS THE CMB FLUCTUATION LEVEL $10^{-5}$ ?

Max Tegmark<sup>a,b</sup> and Martin J. Rees<sup>c,1</sup>

<sup>a</sup>Institute for Advanced Study, Princeton, NJ 08540; max@ias.edu

<sup>b</sup>Hubble Fellow

<sup>c</sup>Institute of Astronomy, University of Cambridge, Cambridge CB3 0HA, UK; mjr@ast.cam.ac.uk

ApJ, in press. Submitted September 8, 1997, accepted December 8.

### ABSTRACT

We explore the qualitative changes that would occur if the amplitude  $Q \approx 10^{-5}$  of cosmological density fluctuations were different. If  $Q < 10^{-6}$ , the cosmological objects that form would have so low virial temperatures that they may be unable to cool and form stars, and would be so loosely bound that even if they could produce a supernova explosion, they might be unable to retain the heavy elements necessary for planetary life. If  $Q > 10^{-4}$ , dense supermassive galaxies would form, and biological evolution could be marred by short disruption timescales for planetary orbits. If  $Q$  were still larger, most bound systems would collapse directly to supermassive black holes. These constraints on  $Q$  can be expressed in terms of fundamental constants alone, and depend only on the electromagnetic and gravitational coupling constants, the electron-proton mass ratio and the matter-to-photon ratio. We discuss the implications for inflation and defect models, and note that the recent anthropic upper bounds on the cosmological constant would be invalid if both  $Q$  and  $\Omega$  could vary and there were no anthropic constraints on  $Q$ . The same applies to anthropic bounds on the curvature parameter  $\Omega_k$ .

### 1. INTRODUCTION

A key parameter in the standard adiabatic cold-dark matter-based models of structure formation is the amplitude that fluctuations in the gravitational potential have when they enter the horizon. This number, which we will denote  $Q$ , has been measured by the COBE satellite to be of order  $10^{-5}$  (Smeets et al. 1992; Bennett et al. 1996), and is assumed to be virtually independent of scale in the most popular models. Why  $10^{-5}$ ? The answers proposed by theorists fall into two categories:

1.  $Q \approx 10^{-5}$  can be computed from first principles using some (still undiscovered) fundamental theory.
2.  $Q \approx 10^{-5}$  cannot be computed from first principles, since the correct fundamental theory merely predicts an ensemble of superhorizon-sized spatial regions with a wide range of  $Q$ , forcing us to treat  $Q$  as random number subject to various anthropic selection effects.

The purpose of this paper is to consider such selection effects, by studying how the physical processes of structure formation depend on  $Q$ . Our motivation for this is threefold:

1. It affects which inflation/defect models should be considered natural as opposed to fine-tuned.
2. It is related to a crucial loophole in the recent arguments for an anthropic upper bound on  $\Omega$ .
3. It poses useful test problems for comparing cosmological simulations.

The structure of our Universe is fixed by a rather small number of physical parameters. The electron mass, the

neutron mass and the low energy coupling constants of the four basic forces determine the physical properties of most objects on scales ranging from the atomic to the galactic (see e.g. Carr & Rees 1979; Davies 1982; Barrow & Tipler 1986), and these parameters can in turn be computed from the roughly 20 free parameters of the standard model of particle physics. A number of additional parameters are often thought of as initial data laid down in the early Universe: the baryon-to-photon ratio, the relative abundances of various dark-matter candidates, the vacuum density contributed by a cosmological constant  $\Lambda$ , the spatial curvature (related to  $\Omega_k$ ) and the amplitude  $Q$  of cosmological density fluctuations, although it is not implausible that abundances such as  $\Lambda$  can ultimately be derived from other particle physics constants. Together with the basic laws, these parameters determine when cosmic structures first emerge and how they evolve. Although the detailed outcome in any one locality, and what complex systems evolve there, depends on local accidents, these parameters nonetheless determine the statistical properties.

Will it ever be possible to compute the values of all these parameters from first principles, within the framework of some yet to be discovered fundamental theory? The answer is a resounding no within some variants of inflationary cosmology (e.g., Linde 1983, 1987, 1990, 1995; Linde & Zelnikov 1988; Coleman 1988; Albrecht 1994; Vilenkin 1995abcd; Vilenkin & Winitzki 1997), where the spatial region that we conventionally call "our Universe", itself perhaps extending far beyond the present observational horizon, is just one element in an ensemble whose members have widely disparate properties. Some physical parameters may take a range of different values throughout this ensemble of exponentially large and causally disconnected regions. The predictions of such theories therefore take the form of probability distributions for the parameters in

question, and these must be computed in Bayesian fashion taking into account the selection effect that observers are not equally likely to inhabit all parts of the ensemble. For instance, just as we expect low surface brightness galaxies to be underrepresented in many surveys, we might expect O-stars to be underrepresented in solar systems containing planet-based extraterrestrial civilizations and, as we shall see, spacetime regions with  $Q \sim 10^{20}$  to be underrepresented in the set of regions that contain observers. The importance of such anthropic selection effects was stressed by Carter (1974), and is discussed in great detail in books by, e.g., Davies (1982) and Barrow & Tipler (1986). More recent reviews can be found in, e.g., Balashov (1991) and Tegmark (1997).

### 1.1. Inflationary predictions

Many inflationary models predict an ensemble of exponentially large space-time regions, each with a different value of  $Q$  (see e.g. Linde 1990; Vilenkin 1995abcd and references therein). Although the cosmological literature abounds with remarks on the "unnaturally" high potential required to produce  $Q \sim 10^5$  in our own Hubble volume, often as a motivation to study defect models, one can just as well argue that it is unnatural that the potential is not even flatter, since super at potentials make inflation last longer and hence dominate the ensemble by volume (Vilenkin 1995a). This dispute cannot be resolved without taking the inevitable anthropic selection effects into account: if these turn out to place a firm upper limit on  $Q$  near the observed value, then inflation models predicting ensembles peaked at high  $Q$  clearly require no fine tuning to explain why we observe  $Q \sim 10^5$ . Conversely, if these selection effects give a firm lower limit on  $Q$  near  $10^5$ , then inflation models predicting ensembles peaked at low  $Q$  require no fine tuning.

### 1.2. The cosmological constant puzzle

Another hotly debated parameter is  $\Lambda$ , the cosmological constant. Although one might expect the most "natural" value of the vacuum density to be of order the Planck density, the observational upper limits on  $\Lambda$  are a striking factor of  $10^{123}$  smaller. This has led to fine tuning criticism of cosmological models with  $\Lambda \neq 0$ , the argument being that they were ruled out at high condence, since such a small value of  $\Lambda$  was extremely unlikely (see Dolgov 1997 for an up-to-date review). As was pointed out by Barrow & Tipler (1986), Weinberg (1987, 1989) and Efstathiou (1995), there is a flaw in this argument, since it neglects a powerful anthropic selection effect. If  $\Lambda$  is too large, then the Universe becomes vacuum dominated before the density fluctuations have grown enough to form non-linear structures. Hence the fluctuations stop growing, and neither galaxies nor observers will ever form. It is therefore no surprise that we find ourselves in a region where  $\Lambda$  is small. A calculation of the probability distribution for  $\Lambda$  given our existence shows that values of order of the current limits are in fact rather typical (Efstathiou 1995), and more accurate calculations (Weinberg 1996; Martel et al. 1997) have confirmed this conclusion.

Unfortunately, there is a loophole in this argument (Rees 1997). As described in more detail in Section 5, increasing  $\Lambda$  by some factor  $f$  can be completely offset by increasing  $Q$  by a factor  $f^{1/3}$  as far as this argument is concerned.

Whether this is really a loophole thus depends crucially on the topic of the present paper, specifically on whether observers could exist if  $Q \sim 10^5$ . The analogous potential loophole exists for anthropic lower bounds on  $\Lambda$  (cf. Barrow 1982; Vilenkin & Linde 1997).

### 1.3. Simulation-testing

A third and entirely different motivation for exploring counterfactual values of parameters such as  $Q$  is that it provides a challenging and bias-free test of cosmological simulation techniques. State-of-the-art simulations including hydrodynamics (which breaks the degeneracy between  $Q$  and  $t$  in pure gravity simulations), gas chemistry and star formation often achieve a good fit to our actual universe (see e.g. Kang et al. 1994 and references therein), but only after tweaking a number of parameters empirically. It is therefore unclear to what extent the agreement between different groups is due to realistic modeling as opposed to simply living in (and parameter-tuning to) the same Universe. It would be far more convincing if two groups could obtain indistinguishable results for hypothetical universes with other values of  $Q$ , where the answer would not be known beforehand.

In Section 2, we outline how  $Q$  affects structure formation in a universe with  $\Omega = 1$  and  $\Lambda = 0$ . We discuss the effects of lowering and raising  $Q$  in Sections 3 and 4, respectively, and the effects of changing  $\Omega$  and  $\Lambda$  in Section 5.

## 2. GALAXY FORMATION AND COOLING

### 2.1. Notation

We will find it convenient to work in Planck units where  $\hbar = c = G = k = 1$ , and the fundamental units of length, time, mass and temperature are  $t_{pl} = (\hbar G/c^3)^{1/2} \approx 5.4 \times 10^{-44}$  s,  $m_{pl} = (\hbar c/G)^{1/2} \approx 2.2 \times 10^{-8}$  kg and  $T_{pl} = (\hbar c^5/G)^{1/2} \approx 1.2 \times 10^{32}$  K, respectively. Important dimensionless constants that will recur frequently are the electromagnetic coupling constant  $\alpha = e^2/4\pi \approx 1/137$ , the gravitational coupling constant  $\alpha_g = m_p^2/4\pi \approx 6 \times 10^{-39}$ , the electron-proton mass ratio  $m_e/m_p \approx 1/1836$ , the baryon-to-photon ratio  $\eta \approx 10^9$ , the baryon fraction  $\Omega_b \approx 0.04$  of the nonrelativistic matter density (which we take to equal the critical value that gives a spatially flat Universe) and the matter-to-photon ratio

$$m_p \frac{\Omega_b}{\Omega_g} = \frac{1}{g} \frac{\Omega_b}{\Omega_g} \approx 10^{27} \quad (1)$$

This constant is simply the amount of nonrelativistic matter per photon,  $\Omega_b/\Omega_g \approx n_m/n_\gamma$ , measured in Planck masses. As our goal is to highlight the main physical effects rather than to make detailed numerical calculations, we will frequently use the symbol  $\Omega_b/\Omega_g$ , which we take to mean that numerical factors of other unity ( $\alpha$  and the like) have been omitted. For instance, the hydrogen binding energy (1 Rydberg), the Bohr radius and the Thomson cross section are given by  $Ry = \frac{1}{2} \frac{1}{g} \approx 13.6$  eV,  $a_0 = \frac{1}{g} \approx 0.38$  Å and  $\sigma_T = \frac{2}{3} \frac{1}{g} \approx 6.65$  Å<sup>2</sup>, respectively.

The reader may find it unfamiliar to see almost no reference below to familiar quantities such as the redshift  $z$ ,

the current CMB temperature  $T_0 = 2.726\text{K}$ , the current Hubble constant  $H_0$  and the current density parameter  $\Omega_0$ . This is because we strive to highlight how structure formation depends on fundamental parameters, and these quantities are not fundamental since they have meaning only once the epoch at which we happen to be living has been specified. Indeed, for the the open Universe case,  $T_0$ ,  $H_0$  and  $\Omega_0$  can be thought of as merely alternative time variables, since they all decrease monotonically with  $t$ . For instance, we are not interested in examining what  $Q$ -values allow galaxies to form by the present epoch  $t_0 = 10^{10}$  years, but what  $Q$ -values allow them to form at all.

## 2.2. When non-linear structures form

The rising curves in Figure 1 show when different mass scales go non-linear, defined as the time when linear perturbation theory predicts an overdensity of 1.69 in a top hat sphere containing the mass  $M$  (Press & Schechter 1974). The curves were computed for the cold dark matter (CDM) power spectrum of Bond & Efstathiou (1984) with  $h = 0.5$ , "shape parameter"  $n = 0.25$ , and an  $h^{-1}\text{Mpc}$  normalization  $\sigma_8 = 0.7$  ( $Q = 10^5$ ). We assume a standard spatially flat Universe ( $\Omega = 1$ ,  $\Lambda = 0$ ) everywhere in this paper<sup>1</sup> except in Section 5. Since fluctuations cannot grow before the matter-radiation equality epoch<sup>2</sup>  $t_{\text{eq}} = 2 \times 10^{11}\text{s}$  (the vertical line in the figure), all scales below the horizon mass at this epoch,

$$M_{\text{eq}} = \frac{4\pi}{3} \rho_{\text{eq}} t_{\text{eq}}^3 \approx 2 \times 10^6 M_{\odot}; \quad (2)$$

have similar fluctuation levels, and are seen to virialize roughly simultaneously (up to a logarithmic factor), at

$$t_{\text{vir}} = t_{\text{eq}} Q^{3/2} f_{\text{vir}} \approx 2 Q^{3/2} f_{\text{vir}}; \quad (3)$$

(The origin of the " $3/2$ " is that, during the matter-dominated epoch, fluctuations grow as the scale factor  $a$  and  $a / t^{2/3}$ .) Since the figure shows that the actual curves approach vertical only for very small mass scales, we have included a factor  $f_{\text{vir}}$  in equation (3) which depends weakly on mass.  $f_{\text{vir}} = 1$  for  $M = M_{\text{eq}}$ , with the value for typical galactic scales  $M = 10^{12} M_{\odot}$  being  $f_{\text{vir}} = 0.03$ . Far above this mass scale,  $P(k) / k$  (we assume the standard spectral index  $n = 1$ ), which means that  $M = M_{\text{hor}} Q^{3/2}$ , where the horizon mass is  $M_{\text{hor}} = \frac{4\pi}{3} \rho t^3$  (straight solid line). Thus the curves all have the same shape, and their left and right asymptotes lie about a factor  $Q^{3/2}$  to the right of the two heavy straight lines in the figure, giving the following broad-brush picture. Mass scales  $M < M_{\text{eq}}$  virialize roughly simultaneously, at  $t = t_{\text{vir}}$ . As time progresses, ever larger scales keep virializing, the non-linear mass scale always being a fraction  $Q^{3/2}$  of the horizon mass scale (a fraction  $Q^{1/2}$  in radius). Note that the the number  $10^{16}$  occurring in this crucial mass  $M_{\text{eq}}$  is simply  $\frac{4\pi}{3} \rho_{\text{eq}} t_{\text{eq}}^3$  | the well-known result that a stellar mass  $M = \frac{4\pi}{3} \rho_{\text{eq}} t_{\text{eq}}^3$  (Dyson 1971) was used in equation (2).

<sup>1</sup>We assume a standard scale-invariant Harrison-Zeldovich primordial power spectrum throughout this paper. More general primordial spectra would correspond to a scale-dependent  $Q$ , thus requiring more than a single number for their parameterization.

<sup>2</sup>At  $t_{\text{eq}}$ , the radiation energy per proton,  $T_{\text{eq}} = \frac{4}{3} \rho_{\text{eq}} / n_p$ , equals the dark matter energy per proton,  $m_p = m_b$ , so  $T_{\text{eq}} = m_p = m_b$ . Since the energy density is  $\rho_{\text{eq}} = \frac{4}{3} H^2 t_{\text{eq}}^2$ , the Friedmann equation gives the Hubble expansion rate  $H = \frac{1}{2} \frac{1}{t_{\text{eq}}}$ , and so the age of Universe at this time is  $t_{\text{eq}} = 2 H^{-1} = \frac{2}{H}$ .

## 2.3. Their virial temperature

When an overdensity has collapsed, the resulting virial halo will have a typical density that exceeds the background density by a collapse factor  $f = 18^2$ , i.e.,

$$\rho_{\text{vir}} = \rho_{\text{eq}} \frac{t_{\text{vir}}^2}{t_{\text{eq}}^2} = f^4 f_{\text{vir}}^2 f Q^3; \quad (4)$$

For a CDM halo of mass  $M$ , this corresponds to a characteristic size  $R = (M / \rho_{\text{vir}})^{1/3}$ , velocity  $v_{\text{vir}} = (M G / R)^{1/2} = (M^2 / \rho_{\text{vir}} G^3)^{1/6}$  and virial temperature

$$T_{\text{vir}} = m_p v_{\text{vir}}^2 = \frac{1}{2} \frac{4\pi}{3} \rho_{\text{vir}} R^3 = \frac{1}{2} f_{\text{vir}}^2 f^{1/3} M^{2/3} Q; \quad (5)$$

A number of isotherms are plotted in Figure 1, and we see that as time progresses and ever larger halos form, the virial temperature stops increasing around the characteristic time  $t_{\text{vir}} / Q^{3/2}$  and approaches a maximum value  $T_{\text{max}} = m_p c^2 Q$ , corresponding to a maximum virial velocity  $v = Q^{1/2} c$ . Thus for our  $Q = 10^5$  universe, typical cluster temperatures are  $10\text{keV}$ , about  $10^5$  times the proton rest energy, and characteristic cluster velocity dispersions are  $1000\text{km/s}$ , about  $10^{5/2}$  times the speed of light.

## 3. WHAT IF $Q = 10^5$ ?

This direct link between  $Q$  and halo temperatures immediately indicates why lowering  $Q$  can cause qualitatively different structure formation scenarios. Unless  $m_p c^2 Q$  exceeds typical atomic energy scales  $\approx 1\text{Ry}$ , which corresponds to

$$Q > \frac{\text{Ry}}{k T_{\text{max}}} = \frac{m_e c^2}{m_p c^2} = \frac{1}{1836} \approx 5 \times 10^{-4}; \quad (6)$$

it will be difficult for the gas in these halos to dissipate their energy to collapse and form stars. Hydrogen line cooling freezes out at about  $\text{Ry} = 15\text{eV}$ , for instance, corresponding to  $Q = 10^9$ . We will now discuss cooling constraints in more detail, and see that these cause qualitative changes even for much smaller departures from  $Q = 10^5$ .

The fate of the baryons in a virialized halo depends crucially on the ratio of the cooling timescale  $t_{\text{cool}} = \frac{3}{2} \frac{M}{\dot{M}}$  to the gravitational collapse timescale  $t_{\text{grav}} = (v_{\text{vir}} G)^{-1/2}$  (see e.g. Binney 1977, Rees & Ostriker 1977; Silk 1977; White and Rees 1978). If  $M$  and  $t$  are such that  $t_{\text{cool}} > t_{\text{grav}}$  (the dark-shaded region in Figure 1), the cloud cannot promptly commence free-fall collapse and fragment into stars, but will remain pressure supported for at least a local Hubble time. For the halo formation curve corresponding to  $Q = 10^5$ , the part of the  $t_{\text{cool}} = t_{\text{grav}}$  curve setting the upper limit on galaxy mass is seen to have a logarithmic slope around  $-2$  (because Bremsstrahlung, with  $t_{\text{cool}} / t^{1/2} = \text{const}$ , is the dominant cooling process), corresponding to  $M \propto t^{-2}$  and a constant radius  $R = \frac{3}{2} \frac{M}{\rho_{\text{vir}}} = \frac{3}{2} \frac{M}{f_{\text{vir}}^2 f Q^3} = 50\text{kpc}$  (Carr & Rees 1979). The

corresponding mass scale is seen to be  $M \sim 10^{12} M_\odot$ . For slightly lower  $Q$ , the upper limit is dominated by line cooling in neutral Hydrogen (rightmost bump), Helium (second bump) and any heavier elements released by early stars (not included here). The lower mass limit is set by the  $T \sim 10^4 \text{K}$  isotherm, below which there are essentially no free electrons and both line cooling and Bremsstrahlung become ineffective. Molecular cooling can potentially lower this mass limit slightly (cf. Haiman et al. 1996; Abel et al. 1997; Gnedin & Ostriker 1997; Tegmark et al. 1997), but is ignored in the figure for the same reason as heavy elements: it is irrelevant to our  $Q$ -constraints, which depend only how far the cooling region extends to the right, not on the vertical extent.

What happens if we start lowering  $Q$ ? The first change is that the upper limit becomes set not by Bremsstrahlung but by line cooling. Figure 1 indicates that as we keep lowering  $Q$ , the range of galactic masses narrows down and finally vanishes completely for  $Q < Q_{\text{min}} \sim 10^6$ . Let us express this critical value  $Q_{\text{min}}$  in fundamental constants. The figure shows that it is determined by the "Hydrogen bump" in the cooling function, which is caused by free electrons collisionally exciting neutral Hydrogen atoms into their first excited state, which is immediately followed by emission of a Ly photon. This gives a cooling timescale (e.g., Dalgarno & McCray 1972)

$$t_{\text{cool}} \sim \frac{m_e^2 c}{h^2} \frac{3^2 e^3}{x(1-x)n}; \quad (7)$$

where  $Ry = kT$ ,  $n$  is the total (bound and free) proton number density, and  $x$  is the ionization fraction. In thermal equilibrium, this is given by (e.g. Tegmark, Silk & Evrard 1993)

$$x \sim [1 + 3^{7/6} e^{-Ry}]^{-1}; \quad (8)$$

Substituting equation (8) into equation (7) gives

$$t_{\text{cool}} \sim \frac{m_e^2 c}{h^2} \frac{h}{n} \frac{1}{e^{-4} (1 + 3^{7/6} e^{-Ry})^2}; \quad (9)$$

where the dimensionless quantity in square brackets is minimized for  $\ln[ \dots ] \sim 10$ , corresponding to  $T \sim Ry = 10^{15}/15,000 \text{K}$ . This minimum value is  $e^{-8/3} \sim e^{-2.67} \sim 0.07$ . Equating this minimal cooling timescale with  $t_{\text{vir}} \sim (G \rho)^{-1/2}$  using  $n \sim \rho/m_p$  finally tells us that the latest time at which line cooling can be efficient is

$$t_{\text{max}} \sim \frac{3^2 \ln[ \dots ]^{8/3}}{g^{3/2} \rho^{1/2}} \sim 10^9 \text{s}; \quad (10)$$

Equating this with  $t_{\text{vir}}$  from equation (3) thus tells us that efficient cooling occurs when

$$Q > \frac{1}{\ln[ \dots ]^{16/9}} \frac{4^{3/2}}{g^{4/3}} \frac{4^{3/2} f_{\text{vir}}^{2/3}}{b^{2/3}} \sim 10^6; \quad (11)$$

If  $Q < 10^6$ , then what is the ultimate fate of the quasi-static pressure supported gas clouds? It is plausible that

they will become increasingly rare as their dark matter halos eventually merge into larger (and less dense) halos, thereby never entering a phase of runaway cooling, fragmentation and star formation. However, even in the arguably contrived case where such a cloud escaped any further collisions, and eventually managed to cool after a (perhaps exponentially) long time, perhaps through some exotic mechanism such as 21 cm cooling, and developed a dense, self-gravitating core which fragmented into stars, there would still be reason to doubt whether it could produce intelligent observers. Since the binding energy of the halo is so low (of order  $T_{\text{vir}}$ ), the first supernova explosion might well eject all the gas from the halo, thereby precluding the production of population II stars and planets containing heavy elements.

#### 4. WHAT IF $Q > 10^5$ ?

What happens if we start increasing  $Q$  instead? The allowed mass range for galaxies keeps broadening at a steady rate until Compton cooling suddenly eliminates the upper mass limit altogether. This is because the time scale on which cosmic microwave background (CMB) photons at temperature  $T$  cool an ionized plasma,

$$t_{\text{comp}} \sim \frac{h^3 c^4 m_e}{t (kT)^4 x}; \quad (12)$$

is independent of both its density and temperature (assuming that  $T > 15,000 \text{K}$ , so that  $x \sim 1$ ). Since  $T \sim T_{\text{eq}} (t = t_{\text{eq}})^{-2/3} \sim 10^{-2} t^{-2/3}$ , this timescale  $t_{\text{comp}}$  equals the age of the universe  $t$  at a characteristic time

$$t_{\text{comp}} \sim \frac{6^{5/3}}{g^{13/10}} \frac{9^{5/3}}{4^{5/3}} \sim 10^{16} \text{s}; \quad (13)$$

Setting  $t_{\text{comp}} = t_{\text{vir}}$ , we find that the upper limit to galaxy masses persists only for

$$Q < \frac{4^{5/3}}{g^{3/5}} \frac{6^{5/3}}{9^{5/3}} \frac{4^{5/3}}{f_{\text{vir}}^{2/3}} \sim 10^{4.5}; \quad (14)$$

For larger  $Q$ -values, all mass scales can cool efficiently, so the characteristic mass for the first generation of galaxies will simply be  $M_{\text{eq}} \sim 10^6 M_\odot$ , given by equation (2). This corresponds to a characteristic size  $R \sim t_{\text{eq}} Q / t_{\text{eq}} t^{2/3}$  for newly formed galaxies, which is constant in comoving coordinates (rather than in absolute coordinates, as the above-mentioned cooling scale  $R \sim 50 \text{kpc}$ ). It would plainly need detailed simulations to determine the mix of discs and spheroids, and the effects of subsequent mergers. However, the galaxies could well have a broader luminosity function than in our actual universe (as well as a much higher characteristic mass); and clustering would also extend up to a larger fraction of the Hubble radius.

#### 4.1. Disruption of planetary orbits

Would this qualitative change affect the number of habitable planets produced? Let us first consider the stability of planetary orbits.<sup>3</sup>

Lightman (1984) has shown that if the planetary surface temperature is to be compatible with organic life, the

<sup>3</sup>After submission, it was brought to the authors' attention that Vilenkin discussed orbit disruption caused by high  $Q$ -values at the 1995 Tokyo RESCEU symposium in Tokyo. Orbit disruption constraints on galaxy densities have also been discussed in the context of axion physics (Linde 1998).

orbit around the central star should be fairly circular and have a radius of order

$$r_{\text{au}} \approx 5 \frac{g^{3=4}}{g^2} 10^{11} \text{m}; \quad (15)$$

roughly our terrestrial astronomical unit, precessing one radian in its orbit on a timescale

$$t_{\text{orb}} \approx 15=2 \frac{g^{5=8}}{g^3} 0.1 \text{ year}; \quad (16)$$

An encounter with another star with impact parameter  $r < r_{\text{au}}$  has the potential to throw the planet into a highly eccentric orbit or even unbind it from its parent star. This happens on a timescale  $t_{\text{enc}} = n_{\text{g}} v_{\text{vir}}^2 r_{\text{au}}^2$ , where  $n_{\text{g}}$  and  $v_{\text{vir}}$  denote the typical stellar density and stellar velocity in a galaxy, respectively. Writing  $n_{\text{g}} = f_{\text{g}} \rho_{\text{vir}} / M_{\text{g}}$ , where  $M_{\text{g}} = M_{\text{g}}^1$  and  $f_{\text{g}}$  is the additional factor by which the dissipating baryons collapse relative to the dark matter before fragmenting into stars, the Milky Way is empirically ruled out by  $f_{\text{g}} \approx 10^4$ . For Earth, this gives  $t_{\text{enc}} \approx 10^{12} \text{s}$ , orders of magnitude above its present age. Moreover, the distant encounters that we have experienced in the past have had a completely negligible effect since they were adiabatic. This means that the impact duration  $\tau = v_{\text{orb}} t_{\text{orb}}$ , so that the solar system returned to its unperturbed state once the encounter was over. For hypothetical galaxies forming before  $t_{\text{comp}}$ , on the other hand,  $M_{\text{eq}} = M_{\text{eq}}$ , so the time between non-adiabatic encounters is

$$t_{\text{adiab}} \approx \frac{1}{n_{\text{g}} v_{\text{orb}}^2} \frac{15 \frac{1=4}{g} 6 f_{\text{vir}}^3}{4 Q^{9=2} f^{3=2} f_{\text{g}}} 10^5 \text{ years} \quad (17)$$

for  $Q = 10^4$  and dropping as  $Q^{-9=2}$  if we increase  $Q$  further. In other words, non-adiabatic encounters are frequent events for  $Q > 10^4$ , occurring often during the geological timescales required for a planet to form, cool and ultimately evolve life. In the conservative approximation of ignoring gravitational focusing (assuming that the flyby speed  $v$  exceeds the orbital speed), the typical time interval between  $r < r_{\text{au}}$  encounters is

$$t_{\text{enc}} \approx \frac{1}{n_{\text{g}} v_{\text{au}}^2} \frac{10 \frac{1=2}{g} 4 f_{\text{vir}}^{7=3}}{4 f^{7=6} f_{\text{g}} Q^{7=2}} 10^7 \text{ years} \quad (18)$$

for  $Q = 10^3$ . Requiring this to exceed some geological or evolutionary timescale  $t_{\text{min}}$  thus gives an upper limit  $Q / t_{\text{min}}^{2=7}$ . Although it is far from clear what is an appropriate  $t_{\text{min}}$  to use, the smallness of the exponent  $2=7$  implies that it makes only a minimal difference whether we choose  $10^6$  or  $10^{10}$  years. Taking  $t_{\text{min}} = 10^9$  years  $\frac{2}{g} \frac{3=2}{g^2}$ , the lifetime of a bright star (Carr & Rees 1979), we obtain the limit

$$Q < 16=7 \frac{4=7}{g} \frac{12=7}{g^2} \frac{8=7}{g^3} f_{\text{vir}}^{2=3} f^{1=3} f_{\text{g}}^{2=7} 10^4; \quad (19)$$

This upper limit appears more uncertain than the lower limit from cooling. The momentum kick given to the planet scales as  $v^2$ , so an impact with  $r \approx r_{\text{au}}$  would not necessarily cause a catastrophic disturbance of the planetary orbit — the event rate for this grows only as  $Q^{5=2}$ , or as  $Q^3$  if the galactic stars settle into a disk where  $v$  is roughly independent of  $Q$ . On the other hand, a very close encounter (especially with an O-star) might cause disastrous heating of the planet. In view of this uncertainty, as well as the uncertainty regarding  $f_{\text{g}}$  and  $f_{\text{min}}$ , we now consider two additional effects of raising  $Q$ .

## 4.2. Black hole domination

For much greater  $Q$ -values, of order unity, typical fluctuations would be of black-hole magnitude already by the time they entered the horizon, converting some substantial fraction  $f$  of the radiation energy into black holes already shortly after  $t_{\text{infl}}$ , the end of the inflationary era. At  $t_{\text{eq}}$ , the universe has expanded by a factor  $a = (t_{\text{eq}}/t_{\text{infl}})^{1=2}$ , and the energy densities in black holes and photons have dropped by factors of  $a^3$  and  $a^4$ , respectively. The black hole density will therefore completely dwarf the density of cold dark matter and baryons if  $f \approx a^{-1}$ . Thus even if  $Q \approx 1$ , extremely rare fluctuations that are  $Q^{-1}$  standard deviations out in the Gaussian tail can cause black hole domination if  $(Q^{-1}) \approx a^{-1}$ , where  $(x)^{-1} = \int_0^x \frac{1}{u^2} \exp[-u^2] du$ . This gives the upper limit

$$Q^{-1} \approx \frac{1}{a} \approx 10^{-27} \approx 10^{-1}; \quad (20)$$

where we have simply assumed that  $t_{\text{infl}}$  is within a few orders of magnitude of unity (the Planck time) since this affects the result only logarithmically. As opposed to the previous constraints, this one depends strongly on whether the power spectrum is strictly scale invariant or not — increasing the spectral index  $n$  from its scale-invariant value  $n = 1$  to  $n = 1.3$  causes primordial black hole domination even if  $Q$  is as low as  $10^{-5}$  (Green et al. 1997).

Even if  $Q$  were low enough to avoid black hole formation in the early radiation dominated phase (say in the range  $10^{-3}$  to  $10^{-2}$ ), rampant black-hole formation may still occur in the matter-dominated era. At times of order  $10^6$  years, i.e., shortly after recombination, clumps of order  $M_{\text{eq}}$  will collapse. If dissipation leads to enough reionization to make their Thomson optical depth larger than  $c/v$  (itself of order  $Q^{-1=2}$ ), then they will trap the background radiation and collapse like supermassive stars, without being able to fragment. The dominant structures in such a universe would then be supermassive black holes, and it is unclear whether any galaxies and stars would be able to form. Even if they could, they would be hurtling around at speeds of order a tenth of the speed of light, and it is far from clear how anthropically favorable such an environment would be!

## 5. WHAT IF $\Omega$ AND $\Lambda$ WERE DIFFERENT?

Our discussion above applied to a flat FRW universe with  $\Omega = 1$  and  $\Lambda = 0$ . As we will now describe, anthropic limits on these two parameters are intimately linked with  $Q$ . In Planck units, the Friedmann equation that governs the time evolution of the radius of curvature of the Universe,  $a$ , is conveniently written as

$$\frac{\dot{a}^2}{a^2} = \frac{8}{3} (\rho_{\text{rad}} + \rho_{\text{m}} + \rho_{\text{c}} + \rho_{\text{v}}); \quad (21)$$

where  $\rho_{\text{rad}}$ ,  $\rho_{\text{m}}$  and  $\rho_{\text{v}}$  are the energy densities corresponding to radiation, nonrelativistic matter and vacuum energy (a cosmological constant), respectively.  $\rho_{\text{c}} = 3=8 a^2$  is

the contribution from spatial curvature (the sign is positive if  $\Omega < 1$  and negative if  $\Omega > 1$  for the flat case  $\Omega = 1$ , the radius of curvature is infinite and  $a$  must be redefined). The first three of these densities evolve as

$$\rho_{\text{m}} \propto a^{-4}; \quad (22)$$

$$\rho_{\text{c}} \propto a^{-3}; \quad (23)$$

$$\rho_{\text{v}} \propto a^{-2}; \quad (24)$$

and  $\rho_{\text{r}}$  does not evolve at all. The constant  $\Omega_0$  is defined as the curvature that the Universe would have had at the Planck time if there was no inflationary epoch, and can be evaluated at any time in the post-inflationary radiation-dominated epoch as  $\Omega_0 = c^2 t^2 / a^2$ , during which this quantity is time independent. We have introduced  $\Omega_0$  simply because we need a constant that quantifies the curvature, and the more familiar  $\Omega$  is unusable since it changes with time. The epochs of matter domination  $a_{\text{md}}$ , curvature domination  $a_{\text{cd}}$  and vacuum domination  $a_{\text{vd}}$  are given by  $\rho_{\text{m}} = \rho_{\text{c}} = \rho_{\text{v}}$  and  $\rho_{\text{m}} = \rho_{\text{v}}$ , respectively, i.e.,

$$a_{\text{md}} = 1 / \Omega_0^{1/2}; \quad (25)$$

$$a_{\text{cd}} = 1 / \Omega_0^{3/2}; \quad (26)$$

$$a_{\text{vd}} = 1 / \Omega_0^{1/3} = 1 / \Omega_0^{1/2} \Omega_0^{1/3}; \quad (27)$$

It is well-known that sub-horizon fluctuations can only grow during the matter-dominated epoch, where they grow at the same rate as the scale factor  $a$ . As we saw in Section 2.2, the first non-linear structures therefore form at  $a_{\text{vir}} = a_{\text{md}} Q^{-1}$  providing that the Universe remains matter-dominated until this epoch ( $a_{\text{cd}} > a_{\text{vir}}$  and  $a_{\text{vd}} > a_{\text{vir}}$ ) or otherwise no nonlinear structures will ever form. We thus obtain the two anthropic constraints

$$\Omega_0 < 2 Q^{-1} 10^{59}; \quad (28)$$

$$\Omega_0 < 4 Q^{-3} 10^{124}. \quad (29)$$

Although we tacitly assumed that  $\Omega_0 < 1$  here, the closed case gives essentially the same constraints – indeed, if no non-linear structures have formed at the epoch  $a_{\text{cd}}$  in a closed universe, time is literally running out for not yet evolved life forms, since the Big Crunch is imminent!

In comparison, the current observational limits are (very conservatively)  $\Omega_0 < 10^{123}$  and  $0.1 < \Omega_0 < 2$ , which corresponds to  $a_{\text{vd}} > 10^3 a_{\text{md}}$  and  $\Omega_0 < 10^{57}$ . The conclusion is that although the anthropic upper limits superficially appear quite strong on both curvature and vacuum density, these constraints are only strong if the two variables on the right-hand side ( $\Omega_0$  and  $Q$ ) are independently constrained – which was one of our motivations for studying upper limits on  $Q$ . The parameter  $Q$  probably deserves more attention than it has received in this context so far (e.g. Rees 1979), and the effects of varying the baryon/photon ratio and introducing a non-zero neutrino mass would also warrant further study. We note in passing that we can obtain crude  $Q$ -independent limits on  $Q$  by requiring that our lower limits on  $Q$  not exceed our upper limits. For instance, the virialization epoch of equation (3) will occur too late for cooling to be efficient (after  $t_{\text{max}}$  of equation (10)) unless  $\Omega_0 > 10^{32} Q^{-3/4}$ . Thus the white region in the figure disappears completely if  $\Omega_0 < 10^{32}$ , and

the conservative limit  $Q < 10^3$  gives the (rather weak) constraint  $\Omega_0 > 10^{30}$ . Conversely, the planetary disruption constraint of equation (19) gets stronger if we increase  $\Omega_0$ , and conflicts with the  $\Omega_0$ -independent limit of equation (6) unless  $\Omega_0 < 10^{23}$ . In addition, there are of course separate limits on the baryon fraction  $b$ , in that if there are too few baryons, the cooling becomes less efficient – see equation (10). Lowering  $b$  may also impede galaxy and star formation, since a gas cloud must collapse by a larger factor before it becomes self-gravitating.

For the reader preferring to think in terms of  $\Omega_0$  and redshift  $z$ , the above argument can be re-expressed as follows. If the current matter density is  $\rho_{\text{m}}$ , then vacuum domination occurs at the epoch  $(1+z_{\text{vd}}) = (\rho_{\text{m}})^{1/3}$ . If  $\Omega_0 < 1$ , then the Universe became curvature dominated at a redshift given by  $(1+z_{\text{cd}}) = \Omega_0^{1/2}$ . Since the first structures form at an epoch  $(1+z_{\text{vir}}) / Q$ , the upper limits on  $\Omega_0$  and  $\Omega_0^{1/2}$  thus scale as  $1/Q^3$  and  $1/Q^1$  for the  $\Omega_0 < 1$  case. For instance, maintaining spatial flatness but making a million times larger than the current observational limits could correspond to  $Q = (10^6)^{1/3} 10^5 = 10^3$ , with galaxy formation about ten expansion times after recombination. When the Universe had reached its current age of  $10^{10}$  years, it would have expanded by a further factor  $e^{100}$ , and ours would be the only galaxy in the local Hubble volume – alas, a drab and dreary place for extragalactic astronomers, but not ruled out by the above-mentioned arguments alone – although perhaps by the  $Q$ -arguments that we have presented.

## 6. DISCUSSION

We have explored counterfactual cosmological scenarios with  $Q$  shifted away from its observed value  $10^5$ . We found that qualitative changes occur if we either increase or decrease  $Q$  by about an order of magnitude. If  $Q < 10^6$ , efficient cooling becomes impossible for gas in virialized halos. If  $Q > 10^4$ , Compton scattering against CMB photons enables efficient cooling in arbitrarily massive halos, and the higher stellar densities and velocities may lead to planetary orbits being disrupted before observers have had time to evolve.

Needless to say, this does not preclude that some form of life might evolve in a Universe with a more extreme  $Q$ -value due to lucky circumstances, for instance around a red star that was ejected from its giant host galaxy in a  $Q = 10^3$  scenario. However, as stressed by Vilenkin (1995a), the key feature of anthropic selection effects is not what the rock-solid extreme limits are on a parameter, but which is the most favorable value for producing observers. This point is also emphasized by e.g. Garcia-Bellido & Linde (1995). To predict a probability distribution for the observed value of  $Q$  from some inflationary model (to potentially rule the model out), its a priori probability distribution for  $Q$  (of quantum origin, say) must be multiplied by some Bayesian selection function such as the number of observers or civilizations corresponding to each  $Q$ -value. It seems plausible that much more stars with habitable planets are formed for  $Q = 10^5$  (where perhaps 1%–10% of all baryons are in stars) than in a  $Q = 10^6$  universe where 1000 times lower densities make cooling difficult. Likewise, it appears likely that  $Q = 10^4$  gives fewer planets in favorable stable orbits than  $Q = 10^5$ ,

where close encounters are completely negligible for most stars. In conclusion, it is possible that the anthropic selection function peaks at  $Q \sim 10^5$ . If this is the case, then what Vilenkin terms "the principle of mediocrity" would imply that since we are most likely to be a typical civilization, this is what we should expect to observe.

The authors wish to thank Tom Abel, Andreas Albrecht,

Ted Bunn, Wayne Hu, Robert Kirshner, Andrei Linde, Avi Loeb, Scott Tremaine, Alexander Vilenkin and Ned Wright for helpful suggestions. This work was supported by NASA grant NAG 5-6034 and Hubble Fellowship # HF-01084.01-96A, awarded by the Space Telescope Science Institute, which is operated by AURA, Inc. under NASA contract NAS5-26555.

#### REFERENCES

- T., Anninos, P., Zhang, Y., & Norman, M. L. 1997, *New Astr.*, 2, 181
- Albrecht, A. 1994, in *The Birth of the Universe and Fundamental Forces*, ed. Occionero, F. (Springer: Berlin)
- Bennett, C. L. et al. 1996, *ApJ*, 464, L1
- Balashov, Y. V. 1991, *Am. J. Phys.*, 59, 1069
- Barrow, J. D., & Tipler, F. J. 1986, *The Anthropic Cosmological Principle* (Clarendon Press: Oxford)
- Barrow, J. D. 1982, *Quart. J. R. A. S.*, 23, 344
- Bond, J. R., & Efstathiou, G. 1984, *ApJ*, 285, L45
- Carr, B., & Rees, M. J. 1979, *Nature*, 278, 605
- Carter, B. 1974, in *IAU Symposium 63*, ed. Longair, S. (Reidel: Dordrecht)
- Coleman, S. 1988, *Nucl. Phys. B*, 307, 867
- Dalgarno, A., & McCray, R. A. 1972, *ARA & A*, 10, 375
- Davies, P. C. W. 1982, *The Accidental Universe* (Cambridge U.P.: Cambridge)
- Dolgov, A. D. 1997, preprint astro-ph/9708045
- Dyson, F. J. 1971, *Sci. Am*, 225, 50
- Efstathiou, G. 1995, *MNRAS*, 274, L73
- García-Bellido, J., & Linde, A. D. 1995, *Phys. Rev. D*, 251, 429
- Gnedin, N. Y., & Ostriker, J. P. 1997, *ApJ*, 486, 581
- Green, A. M., Liddle, A. R., & Riotto, A. 1997, preprint astro-ph/9705166
- Haiman, Z., Thoul, A. A., & Loeb, A. 1996, *ApJ*, 464, 523
- Kang, H., Ostriker, J. P., Cen, R., Ryu, D., Hemquist, L., Evrard, A. E., Bryan, G. L., & Norman, M. L. 1994, *ApJ*, 430, 83
- Lightman, A. P. 1984, *Am. J. Phys.*, 52, 211
- Linde, A. D. 1983, *Phys. Lett.*, 129B, 177
- Linde, A. D. 1987, *Physics Today*, 40, 61
- Linde, A. D. 1988, *Phys. Lett. B*, 201, 437
- Linde, A. D., & Zelnikov, M. I. 1988, *Phys. Lett. B*, 215, 59
- Linde, A. D. 1990, *Particle Physics and Inflationary Cosmology* (Harwood: Chur, Switzerland)
- Linde, A. D. 1995, *Phys. Lett. B*, 351, 99
- Martel, H., Shapiro, P. R., & Weinberg, S. 1997, preprint astro-ph/9701099
- Press, W. H., & Schechter, P. 1974, *ApJ*, 187, 425
- Rees, M. J., & Ostriker, J. P. 1977, *MNRAS*, 179, 541
- Rees, M. J. 1979, *Physica Scripta*, 21, 614
- Rees, M. J. 1997, *Complexity*, 3,
- Silk, J. 1977, *ApJ*, 211, 638
- Smoot, G. F. et al. 1992, *ApJ*, 396, L1
- Tegmark, M., Silk, J., & Evrard, A. 1993, *ApJ*, 417, 54
- Tegmark, M., Silk, J., Rees, M. J., Blanchard, A., Abel, T., & Palla, F. 1997, *ApJ*, 474, 1
- Tegmark, M. 1997, preprint gr-qc/9704009
- Vilenkin, A. 1995a, *Phys. Rev. Lett.*, 74, 846
- Vilenkin, A. 1995b, *Phys. Rev. D*, 52, 3365
- Vilenkin, A. 1995c, preprint gr-qc/9507018
- Vilenkin, A. 1995d, preprint gr-qc/9512031
- Vilenkin, A., & Winitzki, S. 1997, *Phys. Rev. D*, 55, 548
- Weinberg, S. 1987, *Phys. Rev. Lett.*, 59, 2607
- Weinberg, S. 1989, *Rev. Mod. Phys.*, 61, 1
- Weinberg, S. 1996, preprint astro-ph/9610044
- White, S. D. M., & Rees, M. J. 1978, *MNRAS*, 183, 341

Tegmark &amp; Rees 1997




Fig. 1. The nine rising curves show the largest virialized mass scale as a function of time for different values of  $Q$ . Structures with  $M < M_{\text{eq}}$  (horizontal line) are seen to all virialize about a factor  $Q^{3/2}$  after the end of the radiation-dominated epoch (shaded, left), whereas for later times, the virialized mass scale asymptotes to about  $Q^{3/2}$  times the horizon mass (shaded, upper left). Cooling is inefficient in the remaining shaded region (right). The star corresponds to the Milky Way.