## The Axis Ratio Distribution of Local and Distant Galaxies.<sup>1</sup>

Stephen C.O dewahn<sup>2</sup>, David Burstein, and Rogier A.W indhorst

Dept. of Physics & Astronom y, Arizona State University, Box 871504, Tempe, AZ 85287-1504

Em ail: sco@ poseidon caltech edu, raw@ cosm os.la asu edu, burstein@ sam uri.la asu edu

Send preprint requests to \scol poseidon caltech edu"

D ate of this version: M arch 21, 2024

Received \_

accepted

<sup>1</sup>Based on observations with the NASA/ESA Hubble Space Telescope obtained at the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA Contract NAS 5-26555.

<sup>&</sup>lt;sup>2</sup>Currently at Department of Astronomy, California Institute of Technology

### ABSTRACT

Surface photom etry from 16 H ST /W FPC2 elds in the  $I_{e14}$  -band liter is used to derive the distribution of apparent axis ratios for galaxies in progressively fainter magnitude intervals for  $I_{814}$  25. We assess the system atic and accidental errors in ellipticity measurements as a function of image resolution and signal-to-noise ratio, and statistically correct for the ellipticity of cosm ological surface brightness dimming on our isophotal measurements. The axis ratio distribution for the local galaxy population was computed using  $\log R_{25}$  measurements for 1569 RC3 galaxies with  $B_T$  13 mag. Nonparametric tests are used to show that our distant samples, in the redshift range  $0.1 \le z \le 1.0 \{1.5, are not statistically dillocation and signal the evolution and signal the evolution and signal the evolution and statistically dillocation and explanent$ magnitude ranges and discuss the evolutionary consequences of the lack of astrong dill erence between the ellipticity distributions in near and far data sets.

Subject headings: galaxies: elliptical | galaxies: spiral | galaxies: irregular | galaxies: evolution | galaxies: fundam ental param eters

#### { 2 {

#### 1. IN TRODUCTION

The apparent isophotal axis ratio, b=a, of a galaxy is related to its intrinsic threedimensional structure. A statistical recovery of the frequency distribution of intrinsic shapes using apparent axis ratios is dependent on the condition that galaxies are random ly oriented in space with respect to the observer. It is also dependent on the condition that known biases in the galaxy sample, such as those of M alm quist-type, or m easurem ent system atics, and dust, etc., can be adequately m odeled.

The rst detailed study of the axial ratios of galaxies was made by Hubble (1926), who established the basic m ethodology used in later papers by de Vaucouleurs (1959), Sandage, Freem an & Stokes (1970), and de Vaucouleurs & Pence (1973). Subsequent work by Noerdlinger (1979) and Binney & de Vaucouleurs (1981) used mathematical inversion techniques to transform the observed f (b=a) to the intrinsic distribution. O dewahn (1989) applied the modeling methods of the earlier papers to the much improved axis ratios and diameters that refer to a standard isophotal level in the RC3 (cf. de Vaucouleurs et al. 1991) to study intrinsic galaxy shapes as a function of morphological type. More extensive studies of this type, employing larger samples of photographic and CCD galaxy surface photom etry are presented by Ferguson & Sandage (1989), Fasano & Vio (1991), Lambas et al. (1992), and Ryden & Temdrup (1994).

The principal aim of these papers was to determ ine the distribution of intrinsic shapes for galaxies drawn from di erent population samples, e.g., from di erent morphological types or surface brightness classes. The intrinsic shape of the optical component of a galaxy is likely to be linked to the physical state of the dark matter halo and hence may present in portant information about the history and dynamical state of the system. Yet, any statistical approach to determ ine true axial ratios from observed ellipticities is valid only if two simplifying assumptions are made, and various systematic elects are taken into account.

#### { 3 {

These two simplifying assumptions are: a) that galaxies are random ly oriented in space with respect to the observer; and b) that all galaxies in a given sample have the same intrinsic three-dimensional form. Systematic e ects include surface-brightness selection e ects, M alm quist-bias types, sky subtraction errors, internal dust properties, and triaxiality of early-type (elliptical) galaxies. Hence, in practice it is very di cult to derive the 3-D shapes of galaxies. Lam bas et al. (1992) nd strong evidence for triaxiality among the early-type ellipticals, a situation which complicates the derivation of intrinsic shapes using the apparent axis ratio distribution. Our present purpose is not to derive the intrinsic shapes of galaxies as a function of Hubble type. Rather, we wish to use the observed ellipticity distribution as tool to search for evidence of dynam ical galaxy evolution over a reasonably large fraction of the age of the Universe. We want to investigate if di erent galaxy populations with di erent ellipticity distributions (i.e., diskor bulge-dom inated populations) are present at di erent ux levels (i.e., di erent lookback times). In particular, we want to investigate if the epoch of the onset of disk form ation can be delineated from the axis-ratio distribution of galaxies observed with the Hubble Space Telescope (HST).

Recently, Im etal. (1995) used the form of f (b=a) to make inferences about the faint galaxy populations surveyed with HST. The  $0^{\circ}$ I FW HM resolution provided by HST'sW ide Field P lanetary C am era 2 (W FPC2) allows the determ ination of the sub-kpc m orphology and isophotal shapes of distant galaxies | as well as the population fractions of di erent galaxy types, over a wide range of epochs. The m orphological properties of the faint galaxy population have recently been studied with HST by D river et al. (1995a, D 95a; 1995b, D 95b), G lazebrook et al. (1995), C ow is et al. (1995), O dewahn et al. (1996), and A braham et al. (1996). These m orphology-based studies found that a wider range of peculiar or disturbed m orphologies are observed at faint m agnitudes (  $20^{\circ}$  I  $^{\circ}$   $2^{\circ}$  1(2), com pared to what is seen in the local U niverse. U ncertainties arising from cosm ologicale ects such as bandpass shifting and (1 + z)<sup>4</sup> surface brightness (SB) dim m ing com plicate the interpretation of m orphological classi cation systems for studying galaxy

evolution (Burg et al. 1997). The goal of the present study is therefore to compare the properties of nearby galaxies to those seen at high redshift with HST without regard to predicted Hubble type.

These tests using the galaxy axial ratio distributions are a useful counterpart to m orphological studies to understand how galaxies at high z com pare to those near us. In etal. (1995) presented the axis ratio distributions for di erent H ST galaxy types, as determ ined by bulge- and disk-decom position of the H ST in ages. This, how ever, assume es that the galaxy types can be determ ined reliably and independently from the axis ratios. It is not a priori clear that this can be done, and that the axis ratios therefore are independent of the galaxy type. Strictly speaking, we do not know which galaxy types seen at high redshift evolve into which kind of local counterparts in a universe if hierarchical clustering dom inates, and therefore we prefer not to separate the axis ratios as function of galaxy type. Therefore, in this paper, we will com pute the axial distributions for all galaxies detected on the deep H ST in ages, irrespective of their predicted Hubble type. W hile we forego quantitative inform ation of the axial ratio distributions of speci c H ubble types, we bene t from inclusion of all galaxies, whether or not easily classi ed in a m orphological type based on present-day galaxies (e.g., linear or \chain" galaxies).

In x2, we discuss the observations and their reduction using the autom ated in age analysis system MORPHO. In x3, we discuss the system atic and accidental errors in axis ratio measurements in the regime of low resolution and/or low S/N ratio. We also discuss the errors that exist in the axial ratio distribution of the local galaxy sample from the RC3. In x4 we present our determination of the frequency distributions of apparent axis ratio in progressively fainter magnitude intervals. In x5 we discuss how our results can be related to existing morphological tests of galaxy evolution in terms of the hierarchical clustering/merger paradigm for the form ation of galaxies.

## 2. OBSERVATIONS and REDUCTION

{ 5 {

#### 2.1. Observations

The HST data archive was accessed via STEIS to obtain a number of random W FPC2 elds with F814W in ages. The F814W liter was chosen to make a consistent m easurement of axial ratios in the redshifted passband which | for  $0.5 \le z \le 1.0$  | comes closest to the B passband for local galaxies. The deepest eld analyzed in this work was the Hubble Deep Field (HDF; W illiam s et al. 1996), which for the present purposes has 49 exposures in F814W, yielding a 1 surface brightness (SB) sensitivity of  $I_{614}$  28 m ag arcsec <sup>2</sup>, and a 5 point-source sensitivity of  $I_{614}$  ' 29.0 m ag (O dewahn et al. 1996).

A deep W FPC2 eld surrounding the weak radio galaxy 53W 002 at z = 2.39, was in aged in Cycles 4{5 using the F814W lter for 12 orbits (see e.g., D river et al. 1995a). These dithered in ages reach a 1 SB sensitivity of I<sub>814</sub> 26:7 m ag arcsec <sup>2</sup>, and a 5 point source sensitivity of I<sub>814</sub> ' 28.0 m ag (W indhorst et al. 1997). M oreover, the dark H ST sky at 53W 002's location near the N orth E cliptic Pole and the sm all galaxy scale-lengths (O dewahn et al. 1996) allow detection of com pact galaxies in this eld down to F 27 m ag. M orphological studies of these in ages (hereafter referred to as the \W 02" data set) are presented by D river et al. (1995a) and O dewahn et al. (1996).

Two other sets of observations go less deep than HDF and W 02: (1) the eight W FPC2 anking elds surrounding the HDF (W illiams et al. 1996) have mean integration times from 2500 sec to 5300 sec (hereafter referred to as the \HFF" data set). Six M edium D eep Survey Fields have mean integration times from 866 sec to 2100 sec (D river, W indhorst, & G ri ths et al. 1995b; the \DW G " data set). Together, these 14 elds cover a sky area of 0.0168 deg<sup>2</sup>. Based on their total F814W galaxy number counts, we estimate that these shallower in ages have a 90% completeness limit of  $I_{814} \leq 24.0$  m ag for galaxies of average SB, although our use of these elds, listed in Table 1, will be restricted to the higher S/N in ages with total ux  $I_{814} \leq 22$  m ag.

{ 6 {

#### 2.2. Data processing and catalog generation

Initial photom etric catalogs of galaxies in each W FPC2 eld were made using the SExtractor software (Bertin & A mouts, 1996). This package is designed to locate discrete galaxy in ages and extract photom etric parameters for extended sources in crowded galaxy elds. Candidates are de ned to have at least 8 connected pixels with ux 2 above the local median sky. In each case, a 3 3 gaussian smoothing liter is applied prior to detection. A dequate S/N is required to determ ine reliable isophotal axis ratios, so we cut our nal catalogs well above the form al limiting magnitudes of each sam ple: i.e.,  $I_{814}$  22.0 m ag for the HFF and DW G elds,  $I_{814}$  25.0 m ag for the W 02 eld, and  $I_{814}$  26.0 m ag for the HDF.Photom etric zeropoints for each W FPC2 in age were computed from the mean W FPC2 integration times and the synthetic zero points listed in Table 9 of Holtzm ann etal. (1995).

A utom ated surface photom etry of all cataloged objects was performed with the software package MORPHO (O dewahn 1995,1997; O dewahn et al. 1996), designed for neural network-based morphological classication of galaxies. The initial galaxy catalogs generated by SE xtractor are used to cut postage stamp in ages around each detected galaxy. Local sky values are substituted for faint galaxy in ages close to each given object in terms of elliptical areas whose size, shape and orientation are established in a previous iteration by SE xtractor. Local sky estimates are made by the MORPHO package by iteratively tting a sloped plane to a surrounding empty sky-box (see W indhorst et al. 1991), excluding the area of all other objects found by SE xtractor in that sky-box. We used for the rst step an iterative rejection algorithm that determines the best local sky mode.

We compared the sky values from this automated surface photom etry to those for 50 faint objects in the same W 02 images measured by Pascarelle et al. (1996) with a completely independent interactive package, that otherwise similarly used a sloped plane-t to a surrounding visually empty sky-box. Both methods produce sky-estimates consistent within 0.07% (and consistent with M onte C arlo tests of the accuracy of this interactive sky

#### { 7 {

determ ination by W indhorst et al. 1991; W 91). The two m ethods also give total aperture m agnitudes mutually consistent within 0.05 m ag, with rm s' 0.22 m ag for each algorithm at the faint ux levels. Hence, we believe that our sky-errors are dom inated by random (tting) errors, but not by large-scale system atics.

W e also compare the total magnitudes from the MORPHO  $(I_s)$  and SExtractor  $(I_M)$  packages for about 1200 galaxies. U sing galaxies with  $I_{314}$  22, a mean zeropoint o set of  $hI_s$   $I_M$  i = 0.01 was derived (negligibly small). A 3% scale error, signi cant at the 3 level, was evident between the MORPHO and SExtractor magnitude systems. This result was anticipated since we compare the \K ron" magnitudes of SExtractor (magnitudes extrapolated to near-total using the second moment of the light distribution) to the isophotal magnitudes of MORPHO (integrated magnitude measured within an isophotal aperture sized to match a ducial isophotal level). As expected, the SExtractor magnitudes are consistently brighter than the corresponding MORPHO values. For I 24 mag, this system atic di erence is described wellby:

$$I_{\rm S} = I_{\rm M} = 0.01 \quad 0.035 (I_{\rm S} = 22)$$
 (1)

We have adopted the MORPHO magnitudes for use in this paper, but give the following relation to bring the isophotal MORPHO magnitudes onto the total SExtractor system:

$$I_{\rm S} = 0.966 I_{\rm M} + 0.73$$
 (2)

Over the range of m agnitudes to be investigated here,  $19 < I_{814} < 25$ , this system atic di erence will result in a change of at most 0.25 m agnitudes, or 12% of the width of the binning interval used in this work to divide our galaxy samples.

Surface photom etry is performed on each of the patched postage stamp in ages using the MORPHO in age analysis package. This provides ellipse to over a range of isophotal levels (based on the local sky-noise) and elliptically averaged surface brightness pro les, as well as sets of photom etric param eters known to show signi cant dependence with m orphological type form ore nearby galaxies as well as HST galaxies (O dewahn & A ldering 1995, O dewahn et al. 1996).

As discussed in x4, we expect that a signi cant portion of our faint galaxy sample ( $\Gamma$  24) will have redshifts of  $z^{>}$  0.5, and hence will su er signi cant cosm ological SB dimming. For this reason, an ellipse tting package was added to MORPHO that uses axis ratio measurements over a range of isophotal levels and that uses an estimate of the redshift, z, based on the expected median redshift vs.  $I_{814}$  magnitude relation. For this, we used the measured redshift-magnitude relation from Lilly et al. (1995) for I 22 mag, and photom etric redshift estimates by Sawicki et al. (1996) for I 25 mag, to estimate the proper restfram e surface brightness level for measuring the apparent axis ratio b=a.

We cleaned our nalgalaxy sample using an interactive MORPHO package which plots m osaics of the selected galaxy postage stamps with the isophotal ellipse ts overplotted. In some cases, in age merging due to nearby galaxies or di raction spikes from nearby stars made it in possible to compute reasonable ellipse ts. Such cases, comprising less than 5% of the cataloged in ages, were interactively rejected from subsequent analysis. These rejected galaxies were isolated and their f (b=a) distribution was compared to that of the bona de galaxy sample to assure that no strong signature in the apparent axis ratio distribution was being rem oved.

### 3. SYSTEMATIC ERRORS IN AXIS RATIO MEASUREMENTS

G iven the fundam ental di erences between the HST in ages and those from di erent ground-based telescopes, it is inevitable that system atic errors will exist for any com parison of axial ratios of galaxies observed with these instrum ents. As a result, we should expect that com parisons between photom etric properties of local and high redshift galaxies will likewise su er from system atic measurement errors. We note that such errors occur at both low redshift and at high redshifts. For example, even with the superbresolution of HST, images of high redshift galaxies ( $z^{>}$  1) su er from poor linear resolution and low S/N . As we go fainter with the HST images, our ability to measure meaningful isophotal parameters such as size and shape becomes progressively impaired. Separately, existing axial ratio data for nearby galaxies su er either from being a mix of direct measurements from images and eye estimates (such as in the RC3), and/or from problems of zero point calibration and sky subtraction issues (e.g., purely CCD-based surveys).

We adopt the point of view that there is no one \correct" measurement of the axial ratios of galaxies, but rather a series of dierent kinds of measurements that need to be brought onto the same internal system. As such, it is imperative to understand how our apparent axis ratio measures are a ected by these systematics as we progress to fainter magnitude intervals. Similarly, we need to test how consistent are our ellipticity data for nearby galaxies.

#### 3.1. Im age Simulations of the Distant Galaxy Sample

The MORPHO program allows us to simulate the appearance of a galaxy of known redshift at some user-speci ed higher redshift. This package is designed primarily to simulate the elects of cosm obgical SB dimming at high redshift predicted by the equations given in W eedman (1986), with K-corrections of Lilly et al. (1995). With the z relation used by Bohlin et al. (1991), Hillet al. (1992), and M utz et al. (1994) for H<sub>0</sub> = 75 km/s/M pc and  $q_0 = 0.5$ , in ages of local galaxies with high resolution and S/N were repixelated to yield the resolution that would have been observed at redshift z with the W FPC2 pixel size of 0%0996. Sets of high redshift in ages were simulated using two sets of input in ages: (1) 63 B<sub>J</sub> in ages of local galaxies (z 0.05) from the Frei et al. (1996) sample, and (2) a subset of 223 galaxies observed with W FPC2 in I<sub>814</sub> with I 22 m ag from D river et al. (1995a, 1995b). In both cases, we want to simulate the appearance of galaxies observed at high

redshift ( $z^{>}$  0.5-1) in the I<sub>814</sub> bandpass with W FPC2.

For the  $B_J$  galaxy sample of Freietal. (1996) with  $B_T$  13 m ag, we assume a mean redshift of hzi = 0:007, based on the mean of 1569 galaxies with measured  $V_{qsr}$  km/s ( redshift relative to the galactic standard of rest) in the RC3 for  $B_T$ 13 m ag. The Frei sample was used to get a rest-frame bandpass as close as possible to that of our HST  $I_{814}$ observations of high redshift galaxies, therefore accounting for the K-correction to rst order. We note that B $_{\rm J}$  , centered at 4500A, may strictly only be used to simulate the  $I_{\rm 814}$ -band appearance of galaxies in the range  $0.6^{\circ}$  z  $^{\circ}$  1, and that higher redshift m odeling will require the use of large sam ples of U in ages. However, as noted by O dewahn et al. (1996), the system atic changes across the  $U_{300}$ ,  $B_{450}$ ,  $V_{606}$ , and  $I_{814}$  lters in elective radius and isophotal axis ratio are sm all (less than 15%) for galaxies in the HDF and 53W 002 elds (see also gure 1 of Burg et al. 1997). Hence, we used the FreiB<sub>J</sub> in ages to simulate galaxy in ages to z 1:6, where the HST  $I_{814}$  lter samples around rest-fram e 3000A . On the basis of z m easurem ents from the CFRS (Lilly et al. 1995) and the photom etric redshift estimates in the HDF (Sawicki et al. 1997), we adopt a median redshift for our sample of 223  $I_{814}$  galaxy in ages with  $I_{814}$  22 m ag of  $z_{m ed} < 0.5$ . The  $B_{450}$  and  $V_{606}$  in ages of these lower redshift galaxies are used to simulate the appearance of more distant galaxies ( $z^{<}$  1.2) observed in I<sub>814</sub> .

Each of the 63 Frei B<sub>J</sub> galaxy in ages is used to simulate galaxies at redshifts of z = 0.3, 0.6, 0.9, and 1.6. Each of the 223 galaxies in the bright I<sub>814</sub> sam ple ( $z^{<}$  0.5) is used to simulate galaxies at redshifts of z = 0.6, 0.8, and 1.2. These simulated images are reduced with exactly the sam e MORPHO routines as used in x 2 for our actual W FPC2 in ages. These simulated measurements are then used to study the change in measured axis ratio as a function of S/N ratio and linear resolution for every type of galaxy, using their known \true" b=a from the high resolution and high S/N ground-based images. The S/N ratio of the galaxy image in the distant galaxy simulation is computed using the integrated signal within an optim ally selected elliptical aperture (measured at an approximate surface

brightness of  $_{\rm I} = 24.5$  m ag arcsec  $^2$  ) and the standard deviation of the local sky, whose mode was determined after a few iterations of outlier rejection. As a measure of in age resolution, R, we use the number of pixel elements along the major axis of this optim al isophotal ellipse. Axis ratio changes are measured with residuals computed as  $(b=a)_{o}$   $(b=a)_{z}$ , where  $(b=a)_{o}$  is the isophotal shape measured in the original local galaxy in age, and  $(b=a)_{z}$  is the value measured in the simulated high redshift in age. For subsets of  $(b=a)_{o}$ , we search for system atic trends with S/N and resolution to assure that the local comparison set is as much as possible bias-free.

The results of the  $I_{814}$  in age experiments for galaxies with  $(b=a)_0$  0.5 are summarized in Figure 1. Two major points are clear from these diagrams: (1) b=a measurements are dominated by accidental errors, as evidenced by the large scatter in both panels of Figure 1, and (2) in ages with intrinsically small axis ratios are measured systematically more round (large b=a) as the resolution and/or S/N are decreased. In Figure 1, we plot for 176 simulated high z galaxy in ages a linear least squares regression line t for  $(b=a)_0$   $(b=a)_z$  vs.  $\log(S/N)$ . A lineagh dominated by a large scatter, the measured slope of = 0.049 0.014 is statistically signing cant at the 3.5 level, and predicts residuals in the sense expected: faint, low S/N in ages are measured to be systematically too round.

In Fig. 1b we show the same t relative to  $\log(R)$  with a tted regression of slope = 0.067 0.019, again signi cant at the 3.5 level, and with again the same e ect: faint compact in ages tend to be measured too round. The residual axis intercepts of these ts predict o sets of  $(b=a)_{0}$   $(b=a)_{z} = 0.05$  at S/N = 100 and R = 32 respectively. Sim ilar ts using in age simulations of galaxies with  $(b=a)_{0}$  0.5 showed no statistically signi cant trends with S/N and R. Sum marizing both results, intrinsically round galaxies at high redshift are not measured to be system atically too at, but intrinsically at galaxies are measured too round. Results obtained with in age simulations using the Frei B<sub>J</sub> galaxies were consistent with our I<sub>814</sub> -band experiment. The measured slopes had the same magnitude and sense as for I<sub>814</sub> simulations. These results were signi cant at only the 2 level due to a smaller number of input galaxies with (b=a) 0 0.5 (only 18 such cases).

As expected, S/N and R are highly correlated (correlation coe cient of 0.86), and hence it is di cult to determ ine if one in age property really dom inates the e ect observed in Figure 1. Independent of this, the important result of these experiments is that we can now predict system atic b=a measurement errors on the basis of easily measured properties of images. The bulk of the errors resulting from this e ect are accidental, but important system atic trends are derived from the data in Figure 1 of the form :

$$(b=a)_{\circ}$$
  $(b=a)_{z} = 0.067 \log R \quad 0.151$  (3)

or

$$(b=a)_{\circ}$$
  $(b=a)_{z} = 0.049 \log S = N 0.148$  (4)

A lthough relatively small, we must consider this systematic e ect in x4 when we compare local galaxies (observed with high resolution and large S/N) and high redshift galaxies (poor resolution and low S/N).

#### 3.2. Consistency of N ear and D istant G alaxy Sam ples

Since we have chosen galaxies at high redshift on the basis of their apparent m agnitudes in  $I_{814}$ , we must choose present-day galaxies by the same criterion in the de-redshifted

Iter for the expected redshifts of our distant galaxies. As such, we choose a bright subset of the B-band data ( $B_T$  13.0 m ag) contained in the RC3 for our local sam ple of galaxies with m easured axial ratios. For each galaxy, the RC3 gives the decim al logarithm of the isophotal axis ratio,  $R_{25} = (D = d)_{25}$ , where D and d are the major and m inor axis lengths m easured at 25.0 m ag arcsec<sup>2</sup> in the  $B_{450}$  (note that the RC3 de nes  $D_{25}$  in units of 0.1 arcm in). As discussed by de Vaucouleurs et al. (1991), the majority of these axis ratios are based on visual estimates from a variety of sources, but have been transformed to a self-consistent system via comparisons with detailed surface photom etry for a sample of 608 galaxies. For galaxies fainter than  $B_T > 13 \text{ m ag}$  the nom inal limit of the original Shapley (Am es (1932) Catalog | the RC3 galaxies represent a heterogeneous sample which su ers from a variety of selection e ects.

The 1569 galaxies in the RC 3 with  $B_T$  13.0 m ag have other parameters in the range: 0.0 log R<sub>25</sub> 1.2, log D<sub>25</sub> 0.5, and {7 T 10.0, where T is the morphological Type in the revised Hubble system. This galaxy sample is essentially a somewhat enlarged version of the original Shapely-Am es C atalog, and should be reasonably complete to  $B_T$  13.0 m ag except for dwarf or extrem ely low surface brightness galaxies.

To test the consistency of the RC3 eyeball axis ratios with our own MORPHO m achine m easurem ents of axis ratios, we compare the MORPHO m easurem ents to the RC3 m easurem ents for 145 galaxies: 63 galaxies observed in B<sub>J</sub> by Frei et al. (1996) and 82 UGC galaxies observed in B by de Jong (1994) using CCD in ages. The autom ated surface photom etry package in MORPHO was used to estimate isophotal axis ratios for these 145 galaxies at  $_{\rm B} = 25$  m ag arcsec  $^2$ . The resulting di erence, b/a (RC3) { b/a (MORPHO) is plotted versus b/a (RC3) in Figure 2. A linear least squares to only the 63 galaxies of the Frei et al. data gives a slope of 0.21 0.01 and an rm s scatter 0.07. The deJong sam ple does not cover a wide range of b/a (the galaxies were selected to have b/a 0.6), but their values of b/a (RC3) { b/a (MORPHO) are consistent with those of Frei et al. in the region where their ellipticities overlap. Hence, it appears that the eyeball RC3 b=a estimates are system atically atter for at objects, and system atically rounder for round objects.

The origin of this small, but statistically signi cant trend is unknown at present, but obviously important for the comparison with HST data. Based on this twe adopt the following correction formula to transform the 1569 RC3 axis ratios to our own MORPHO measurements, so that now for both the low and the high redshift galaxies we can measure a consistent ellipticity by machine:

$$(b=a)_c = 0.80 (b=a)_{RC3} + 0.12$$
 (5)

G iven the uncertain reason for making such a correction, we will in x4 exam ine how both the original RC 3 axial ratio distribution and the axial ratio distribution for our 1569 local galaxy sam ple | when transformed to the MORPHO system | compare to those of high-redshift galaxies.

It is in portant to note that the linear correction in plied by equation 5 is used for convenience, as the present data in this comparison are too few to derive a form ally correct relationship. The reader will note that a correction of this form will deplete the nearly circular apparent axis ratios (b=a = 0.92), as well as predict unrealistically elongated galaxies. This clearly would complicate the derivation of the 3-D axial ratio on the basis of f (b=a) (not our purpose here), especially in light of the triaxility question for early-type system s (Lam bas et al. 1992). A nonlinear expression is required for such a correction, however the lack of highly elongated and circular isophotal shapes in our present sam ple overlapping the RC 3 is to sm all for such a derivation. Since the statistical tests discussed in x4 2 are most sensitive to changes at more interm ediate axis ratios, particularly around the median, we feel justi ed in making this correction for the purposes of com paring low and high redshift sam ples.

#### 4. APPARENT AXIS RATIO DISTRIBUTIONS - NEAR AND FAR

#### 4.1. Subdividing the D istant Sam ple

The 16 W FPC2 elds listed in Table 1 were processed with MORPHO to assemble photom etric catalogs to the depths given in x2 on the basis of total integration time. U sing the  $I_{814}$  -band vs. z data of Lilly et al. (1995) for  $I_{814}$  22 and the photom etric redshift estimates of Saw icki et al. (1997) for  $I_{814}$  25, we specified a series of magnitude intervals for drawing galaxy samples with progressively higher median redshift and adequate statistical size. Unlike the local galaxy samples, thing isophotal shapes at a xed SB level is inadequate. Due to cosm obgical e ects, the change in observed SB is  $= 10 \log (1 + z)$ m ag arcsec <sup>2</sup>. Hence, thing ellipses at a xed isophotal level in a high redshift galaxy gives a measurement of isophotal shape in an intrinsically brighter region of the galaxy. In the case of an early-type, bulge-dom inated system where the higher surface brightness isophotes often have rounder shapes than the lower surface brightness outer disky isophotes, we may expect to see a trend in b=a with surface brightness. The same, or in fact the opposite, may be true for the faint Irr galaxies that dom inate the HST in ages, as we will address in the simulations below.

In order to avoid possible system atic errors in our current analysis, we ran MORPHO in a mode where multiple ellipse ts are made at progressively fainter surface brightness levels. These levels are chosen on the basis of S/N in the local sky-background. For each galaxy, we used the apparent total  $I_{814}$  magnitude to estimate a median z for the corresponding magnitude bin. The cosm ological correction to SB, , is then applied to the desired isophotal level (25.0 m ag arcsec <sup>2</sup> in B<sub>450</sub> was adopted here), and an interpolated value of the axis ratio at the correct intrinsic surface brightness is computed. W hile not a perfect way to compensate for redshift e ects, it is the best we can do until all the galaxies sam pled have individual redshifts measured for  $I_{814}$  <sup>4</sup> 26 m ag, which is a major challenge even for 8{10 m eter telescopes. The nalW FPC2 sam ples consists of:

- 1. a bright sample of 105 galaxies in the range  $I_{814}$  20 m ag with  $hI_{814}$  i = 19:9 m ag at an estimated median redshift of z 0:2 and a mean SB threshold of I = 23:9 m ag arcsec <sup>2</sup>;
- 2. 254 galaxies in the range 20  $I_{814}$  22 m ag with  $hI_{814}$  i = 21:3 m ag at an estim ated median redshift of z 0.5 and a mean SB threshold of I = 24:6 m ag arcsec <sup>2</sup>;
- 3. 210 galaxies in the range 22  $I_{814}$  24 m ag with  $hI_{814}$  i = 22.8 m ag at an estim ated median redshift of z 0.8 and a mean SB threshold of I = 25.5 m ag arcsec <sup>2</sup>; and

4. a faint sample of 296 galaxies in the range 23  $I_{814}$  25 m ag with  $hI_{814}$  i = 24.2 m ag at an estimated median redshift of z 1.0 and a mean SB threshold of  $_{I}$  = 26.0 m ag arcsec  $^{2}$ .

The dimensional frequency distribution of apparent axis ratios for each m agnitude selected sample was computed using xed bins of 0.05 in b=a. The results are plotted in Figures 3 through 5, where the W FPC2 samples are represented as open circles connected by thin dotted lines.

#### 4.2. Comparing RC3 and D istant Sam ples

In accounting for system atic errors in both the distant and nearby galaxy samples, we have chosen to model how RC3 galaxies would appear at the median redshift for each distant galaxy sample, including measured S/N and linear resolution e ects (x3.1). Various methods are used to simulate the corresponding f (b=a) distribution we expect to observe for the RC3 sample at a given median redshift if the galaxy population mix at that epoch were the same as our local RC3 sample. The apparent magnitude of a galaxy in the RC3 sample is estimated for H<sub>0</sub> = 75 km /sec/M pc and q<sub>0</sub> = 0.5 using:

$$m M = 42.38 5 \log h_0 + 5 \log z + 1.086 (1 q_0) z + m_K + m_L$$
(6)

where  $\log h_0 = H_0 = 100$ , and M is the absolute m agnitude of the galaxy computed using  $B_T$  and  $V_{GSR}$  (from RC3). Type dependent K-corrections,  $m_K$ , were computed from F igure 1 of Lilly et al. (1995) and applied to the distance m odulus of equation 6. D river et al. (1995) and Lilly et al. (1995), respectively, note that there is substantial evidence of strong lum inosity evolution since  $z^> 0.5$  for late-type galaxies (T > Sc) and blue galaxies, respectively. The m orphological num ber counts in  $B_{450}$  by 0 dewahn et al. (1996) also show that sm aller levels of lum inosity (and/or density) evolution have occurred in  $B_J$  for E+S0 and early-type spiral populations for  $z^< 1$ .

To account for these possible evolutionary e ects on our apparent m agnitude estimates for the redshifted RC3 galaxies, we applied a redshift-dependent apparent m agnitude increase,  $m_L$ , for E+SO, Sabc and Sd/Irr galaxies following the models of D river et al. (1995b):  $m_L = 0$  for all galaxies with z < 0.2;  $m_L = 0.5$ ; 1:0; 1:5 respectively for E+SO, Sabc and Sd/Irr galaxies with 0:2 z 0:75; and  $m_L = 1:0$ ; 1:5; 2:5 respectively for E+SO, Sabc and Sd/Irr galaxies with z > 0:75. These estimates are used only to investigate the possible e ects of luminosity (and/or in plied density) evolution on our predicted f (b=a) for galaxies out to z < 1.2, and we do not suggest that these are necessarily the exact am ounts of lum inosity evolution for each of these galaxy classes.

On the basis of the apparent m agnitude com puted in this manner, each redshifted RC3 galaxy was binned by apparent magnitude. The resulting magnitude-binned sets of b=a were used to compute the observed f (b=a) distributions for a local (i.e., RC3-like) population seen at high redshift. Each RC3 b=a estimate was corrected to the MORPHO system using Equation 5. These predicted distributions are plotted as solid squares connected by bold solid lines in Figure 3. On the basis of Figure 3, one might infer a depletion of highly attened axis ratios (b=a < 0.4) at progressively higher redshifts (I > 22 mag or z > 0.4) compared to our local population.

We have used two methods to compare the WFPC2 b=a distributions with those from the \redshifted" RC3 data sets. A Kolm ogorov-Sm into probability, P (KS), was computed for each set of distributions to assess the likelihood that the two distributions were drawn from the same parent population. The P (KS) statistic gives the probability that the largest deviation between the two cumulative distributions drawn from each data set is due to random uctuation, and hence a sm all value of P (KS) would indicate that the di erence between two data sets is statistically signi cant. We have used the conservative level of P (KS) 0.01 to con m that two sets are drawn from a di erent parent distribution. Values of P (KS) are tabulated in each b=a set compared in the panels of F igures 3 through 5. For the brightest and faintest magnitude ranges, the P (KS) probability just fails to

{ 18 {

con m that f (b=a) for the W FPC2 samples is di erent from that of the RC3 set. The two interm ediate ranges, covering 20 24, con m to better than a 99.9% con dence I<sub>814</sub> level that the two distributions are drawn from di erent parent populations. As noted by Press et al. (1986), the KS test has the advantage that it is a nonparam etric test that is independent of data binning, but since all cum ulative distributions are the sam e at the low and high ends, it is sensitive primarily to changes in the central region of the distribution around the median. We are primarily concerned with comparing the low axis ratio end of our high and low redshift samples, and hence use a second simple statistic for this case. For each b=a set, we com pute the ratio, (b=a), of the num ber of galaxies with axis ratio sm aller than b=a to the num ber of galaxies with axis ratio larger than b=a . Larger values of  $_{i}$  (b=a) indicate a higher fraction of at cases (low b=a). To compare each set of distributions, we compute the di erence between these values,  $FD = _{RC3}(0:4)$  $W = PC_2(0:4)$ , to form an index that measures the relative enhancement or depletion of at systems at high redshifts. When FD is large and positive then the WFPC2 samples must be relatively poor in at cases relative to RC3. The value of FD for each distribution comparison is also plotted in the panels of Figures 3 through 5. In Figure 3, this simple statistic con m s our KS test results in the sense that we do nd FD to be di erent and progressively larger in the faint m agnitude sam ples.

To see a more realistic distribution of axial ratios predicted for the RC3 at high redshift, we have to take into account system atic measurement e ects due to degradation of S/N and linear resolution with redshift, speci cally the results of equation 3 and equation 4 (x3.1). To do this, the apparent  $I_{814}$  magnitude estimated for each RC3 galaxy was used to estimate the expected R and S/N that would have been obtained had the galaxy been observed with an integration time appropriate to a given data set (i.e., HFF, HDF, etc...). We then applied equations 3 and 4 to the redshifted RC3 distributions based on these values of R and S/N .

Equations 3 and 4 correct the observed b=a values as a function of logR and S=N,

with the highest connections coming for the most-edge on galaxies. In the present case, this means that the most edge-on galaxies in the RC3 seen today will have their axial ratios measured rounder with decreasing values of S/N and R. However, within the errors of the calculations presented in x3.1, we were unable to detect a strong systematic change in measured b=a for galaxies with intrinsic b=a > 0.5. In order to avoid a discontinuous jump in the b=a properties of our connected high redshift sam ples, we have applied a sm oothing factor to the connection predicted by equation 3: the full connection is applied for b=a > 0.45 with the connection decreasing to 10% of that predicted by equation 3 at b=a = 0.8. The basic properties of the predicted high redshift f (b=a) distribution were found to be rather insensitive to the form of this sm oothing function. The systematic b=a changes estimated in this way were applied to all measurements composing the \redshifted RC3" catalog.

The bold-line f (b=a) distributions in F igures 4 are derived from these \resolutioncorrected" redshifted RC3 sam ples. It should be noted that we applied R and S/N corrections independently, and found the results to be unchanged. For sim plicity, we show only the R corrected estimates in our f (b=a) plots. The important point to be noted in F igure 4 is that our apparent dearth of attened systems at high redshift (F 22 m ag or  $z^{>} 0.4$ , see F igure 3) has generally disappeared after applying these corrections. W ith the exception of the 20 I<sub>814</sub> 22 interval, the P (K S) probability indicates no signi cant di erence between the sets of low and high redshift distributions. The large jump in F D that was observed in the high redshift sam ples in F igures 3, which indicated a 19% increase in the relative fraction of attened systems at low redshift, has now dropped to the <sup><</sup> 5% level.

Not only do the S/N and R corrections rem ove m any edge-on galaxies from the high redshift sam ples, the correction to m ake the eyeball RC 3 axial ratio distribution consistent with the MORPHO m achine m easurem ents does so as well. If we had m ade this study com paring just the local RC 3 axial ratio distribution (uncorrected) to that of the distant galaxy sam ples (uncorrected), we would have found a very large di erence in the num bers

#### { 20 {

of edge-on galaxies seen today versus those seen at high redshift, with possibly enroneous conclusions about morphological evolution of galaxies as a consequence. However, to do so would be not correct in our opinion. Rather, by correcting for detectable systematic e ects, we not the present-day axial ratio distributions of galaxies | when moved out to high redshift and accounting for all possible systematics as well as we can | to be very similar to those for HST galaxies with  $0.3^{<}$  z<sup><</sup> 1.0.

To what extent is the lack of detection of a change in ellipticity in the range  $0.3^{\circ}$  z<sup> $\circ$ </sup> 1.0 consistent with what we currently know about the morphologies of galaxies at these redshifts? As discussed in D river et al. (1995), 0 dewahn et al. (1996), and A braham et al. (1996), the galaxy population appears to be increasingly dom inated by late-type (types later than Sc) at high redshift (I<sup> $\circ$ </sup> 21 m ag, z <sup> $\circ$ </sup> 0.5). The f (b=a) distribution for E+S0 and early spiral populations are di erent am ong the local galaxies (B inney & de Vaucouleurs 1980; Lam bas et al. 1992). The E+S0 f (b=a) show a smooth increase in the frequency of galaxies with increasing b=a, while the spirals show a rather at distribution with a sharp drop in frequency around b=a 0.3. For the very latest types (later than Sd), the f (b=a) distributions com puted in O dewahn (1989) are even m ore strikingly di erent from the early- and m id-spiral sam ples, with a pronounced redistribution of objects at the high (= round) b=a end of the distribution. Hence, a population dom inated by late-type spiral system s at high redshifts could produce a relative overabundance of round objects in f (b=a) with b=a <sup>2</sup> 0.6.

As a test of this possibility, we altered the input catalog of RC3 galaxies to re ect a population that is dom inated by a late-type spiral and irregular galaxies, as in the deep HST in ages of D river et al. (1995a) and O dewahn et al. (1996). For our original sam ple of 1569 RC3 galaxies with  $B_T$  13, the type fractional percentages are f(E + S0,Sabc,Sd+ Irr) = (32%, 48%, 20%), respectively. We random ly sam pled this catalog to readjust these values to f(E + S0,Sabc,Sd+ Irr) = 10%, 30%, and 60%, to create a sam ple m ore in line with the population fractions in plied by the faint W FPC2 m orphological studies mentioned

{ 21 {

above. This spiral/Im-rich \local" catalog was shifted to the same high redshift intervals used in Figures 3 (with system atic corrections as in Figure 4), and is shown in Figure 5. As is evident from comparing Fig. 4 to Fig. 5, we see only small changes in the predicted f (p=a) functions when compared to the original local RC 3 type fractions. The tendency to produce a more peaked distribution at the high b=a end is only mildly apparent, and this happens only in the high redshift bins where the observed late-type fraction is increased due to strong lum inosity evolution in the late types (see D river et al. 1995a) and to the loss of early-type galaxies from the redshifted local sam ples due to their larger K -corrections. Hence, it would seem that only an unreasonably extrem e alteration of the type fractions (beyond those allowed by the studies of D river et al. 1995 and O dewahn et al. 1996) would produce a detectable change in our currently observed W FPC 2 sam ples.

Finally, we think that seeing is believing. Hence, we have put together montages of galaxies (Figures 6{ 8) with similar ranges in b/a including: representative Freietal. (1996) galaxies seen today in the B passband; the same Freietal. galaxies placed at z 0:6; and representative galaxies from our W FPC2 in ages, spanning a range of  $I_{814} = 14$  to 25 m ag. A s one can see, there is little di erence in morphology within each range of b/a. These

gures illustrate that our use of B-band in ages for present day galaxies and  $I_{0.14}$  in ages for distant galaxies can produce system atic e ects that are not quanti able with the present data sets. In particular, it should be noted that the appearance of the galaxies becomes m ore asymmetric in the fainter magnitude intervals. This is most evident in the low b=a m ontage of F igure 6. W hether such changes are due to genuine morphological variation, or some cosmological e ect is still open to question. As discussed in Bohlin et al. (1991), G iavalisco et al. (1996) and O dewahn et al. (1996), the appearance of some types of galaxies can change substantially as a given passband e ectively samples bluer or rest-frame UV emission at higher redshift. Even though we have used the  $I_{0.14}$  – liter for our HST data to get a better m atch to the rest-fram e B-band observations for our local sample, this m atch is not perfect. In the future, we will also make a comparison with a U-band survey in progress of a large number of UGC and NGC galaxies with the VATT telescope (see Burg

#### { 22 {

etal. 1997).

#### 5. DISCUSSION

#### 5.1. Comparison to other HST ndings

M orphological type fractions have been shown to change system atically at fainter m agnitudes or at higher redshift. The rather drastic increase in late-type systems at  $z^{>} 0.5$  observed by D river et al. (1995), G lazebrook et al. (1995), and O dewahn (1996) is interpreted as evolution among the galaxy populations at high redshift. From num erous papers referenced in our introduction, it is well known that early- and late-type systems display di erent apparent axis ratio distributions locally, and hence it was surprising that a large change in the fraction of Sd/Irr galaxies at high redshift would not manifest itself strongly in the observed axis ratio distributions. On the contrary, as we have shown in the previous section, only sm all deviations in the f (p=a) distribution for high redshift galaxy sam ples are present compared with local sam ples. In fact, these sm all deviations m ay be totally explained if we take into account system atic errors in isophotal param eters measured in in ages with low linear resolution and low signal to noise. The observed f (p=a) distribution for faint galaxy sam ples will also be a ected by two additional factors: weak lensing and the possible in uence of a population of linear "chain galaxies" at high redshift.

The shapes and orientations of distant galaxy in ages will be distorted from gravitational lensing by intervening mass distributions. Tyson et al. (1990) and K aiser and Squires (1993) have used this elect to probe the mass distributions in rich clusters in an elect to con m mass determinations derived from purely dynamical methods. The samples used in our current study were drawn from a variety of positions on the sky and are not expected to be profoundly in uenced by foreground rich clusters. The presence of rich clusters in our current samples would manifest them selves by increased number counts relative to the eld. On the contrary, the (morphological) number counts derived from the samples used here were found by D river et al. (1995) and O dewahn et al. (1996) to be very consistent.

However, the weak distortion of faint galaxy in ages caused by lensing from large scale structures (LSS) in the Universe must be considered. Large scale deviations in the mass distribution inferred from system atic galaxy redshift surveys by Huchra et al. (1988) and the measurement of uctuations in the cosm ic microwave background by COBE (Smoot et al. 1992) indicate that large m atter overdensities m ay be com m on throughout the U niverse, and that lensing resulting from these inhom ogeneities may produce ellipticity changes of up to 3% for galaxies as distant as z 1. A recent m easurem ent of this weak distortion by LSS is presented by M ould et al. (1994), in a ground-based study of 4363 galaxies to a lim it ofr 26 mag. They nd a weak lensing signal signi cant at the 1 level in the form of a mean position angle polarization of 1% over that expected for a random distribution. The detection of galaxy-galaxy lensing has been studied by Tyson et al. (1985) in a sample of 47000 galaxy in ages in the range 22.5 B<sub>J</sub> 23.5 with no statistically signi cant signal above that expected for a purely random distribution. More recent CCD-based studies by Brainerd et al. (1996) use 3202 galaxy pairs in the range 23 r 24 m ag to detect a lensing signal in the form of a position angle polarization of 1.1% 0.6%. They report that an ellipticity of approximately 2-3%, similar to the observed polarization e ect, will be induced in faint galaxy in ages with angular separations of 3<sup>00</sup> from bright galaxies. In the course of this ground-based study, they considered the ellipticity distribution of galaxies from their faint samples, and determ ine that the ellipticity distributions of these objects are consistent with f (b=a) drawn from brighter, more local sam ples. Finally, the presence of a large am ount of im age distortion by LSS should be accompanied by a large am ount of multiple in aging, an e ect which is not observed commonly in well-studied distant radio galaxy samples (Blandford, private communication). The overall conclusion from these studies is that weak gravitationally lensing may certainly in uence the observed f (b=a) of faint galaxies by \stretching" the isophotal in age contours. However, recent redshift surveys by Lilly et al. (1995) and Ellis et al. (1996) suggest that the median redshift of the galaxy sample used here is z<sup><</sup> 1, and thus the overall in age distortions due to lensing will

produce a rather sm all 1-2% system atic e ect.

Separately, a population of faint blue galaxies with very narrow shapes and lum py m orphologies was reported by C ow ie et al. (1996) and term ed \chain" galaxies, a possibly new m orphological class of galaxy comm on during younger epochs of the Universe. A truly linear structure, such as the chain galaxies are thought to be, will present an isophotal shape having a sm all value of b/a in nearly any spatial orientation. Hence, the presence of a large population of such galaxies in our faint sam ples could skew our observed f (b=a) towards the low b/a end. The presence of such a new m orphological population has been refuted by D alcanton and Shectm an (1996), who claim that these galaxies do indeed exist at low redshift and represent edge-on views of low surface brightness late-type ( $T \ge Sd$ ) disk galaxies. U sing a low-redshift sam ple of low surface brightness galaxies (central disk surface brightness 21.3 m ag arcsec <sup>2</sup> in the r band), D alcanton & Shectm an show that the f (b=a) of this chain galaxy sam ple is consistent with that of a population of thin disk galaxies viewed with random orientations. The galaxies with the the sm allest b=a display m orphological sin ilarities with the \chain" galaxies, especially in the far-UV.

The UIT images by O'C onnell et al. (1997) and M arcum et al. (1997) of ordinary nearby disk galaxies seen edge-on resemble those of high redshift \chain galaxies", so that the clum py rest-fram e UV -m orphology of such objects at high redshift m ay be classi ed as a di erent or new class of object, while in reality they m ay not be. Our observations are consistent with this latter view. However, P ascarelle et al. (1996) note that the C ow ie et al. (1996) chain galaxies are a clear m inority of the faint blue galaxy population in deep W FPC2 images. The faint chain galaxies offen do show clear \dum bbell" structure with m ost of the light in clum ps at both outer ends | unlike m any of the U IT in ages of nearby edge-on galaxies | so that these objects m ight be subgalactic sized clum ps seen directly after or during their rst m ergers, which would trigger subsequent starbursts and form disks from gaseous m aterial that settles into rotation during and after the m erger. H ence, they could be short-lived \chain-like" objects, but their fraction of the total population is

{ 25 {

not large enough to signi cantly a ect our b-a distributions.

# 5.2. Comparison to the H ierarchical C lustering/M erging G alaxy Form ation Scenario

Consider what we might expect to see at high redshift from galaxies that form via the hierarchical cluster/m erger hypothesis (e.g. G ott & Rees 1975). Peebles (1993) de nes lookback tim e to be the tim e m easured backwards from the present tim e, in other words, since z = 0. Hence, larger values of lookback tim e correspond to earlier cosm ic epochs (and larger redshift) in the Universe. The familiar galaxy types we see today are thought to rst start out in the form of  $10^7$   $10^8$  M clumps of gas and new ly form ed stars. We will rst see such pre-galactic clumps when they give o su cient radiation to be detected, at a lookback tim e that we call  $t_f$ . The Pascarelle et al. (1996) cluster of 18 subgalactic sized objects at z' 2.4 is an example of this process. We stress, however, that in hierarchical CDM m odels the form ation of these clumps does not have to be coeval, and will in fact likely happen over a wide range of epochs (z' 1{4}. When these clumps start to combine and form ever larger clumps, at a sm aller lookback tim e  $t_m$  (<  $t_f$ ), we will begin to see the merged clumps as galaxies of m orphologies that are familiar to us today: E/S0's, Sabc's, Sd/ Irr's.

At bokback times smaller than  $t_m$  (i.e., after the merging epoch began), hierarchical models predict that we will see galaxies with a range of morphological types not dissimilar to what we see today, but with a galaxy mix that is not necessarily equal to the local one. What we will not know | if hierarchical clustering is correct | is what a particular galaxy at a lookback time smaller than  $t_m$  will look like today. Will it simply passively evolve? Or will it merge with its nearest neighbors, turning (temporarily) into another type of galaxy altogether? In other words, in a hierarchical merging scenario, while the galaxy population mix may not change in a statistical manner for lookback times less than  $t_m$ , one cannot necessarily uniquely assign galaxy of type X at high redshift with galaxy of type X today. The lack of change in the distribution of edge-on system s with ellipticity that we observe is consistent with hierarchical clustering in the following sense. We know from the distribution of Hubble types today that bulge-dom inated galaxies comprise the minority of giant galaxies (less than 30%, see D ressler 1980). If all galaxies initially are formed as disk-type systems, with subsequent merging producing pronounced spiral bulges and E/S0 galaxies, then we would expect a steady migration from galaxies with low b/a to those with high b/a. W hile such a trend is suggested in the present comparison, the observed result is barely of one-sigm a signi cance. The problem is observational: It is di cult to accurately measure the ellipticities of galaxies with b/a > 0.85. E mores in photom etry will always scatter ellipticities away from b/a = 1.0 (since b=a cannot be larger than 1), always depopulating the highest bin in f (b=a). As discussed in the introduction and in x3.2, our system atic correction of the RC3 axis ratios, as well as uncertainties related to the triaxiality of early-type galaxies, makes treatment of the high b=a system s problem atic.

If there is one place in comparing the ellipticity distributions of nearby and distant galaxies to hide evolutionary e ects, it is an ong galaxies with b/a 0.85. Yet, this is where m any m erger products are likely to end up in terms of their intrinsic axial ratios. W hat we conclude is that nature apparently conspires to have system atic e ects (observational ones and those due to the redshift) on the ellipticity distributions to essentially cancel the e ects expected from the evolutionary change in galaxy m orphology. If the hierarchical clustering/m erging picture is correct, then the kind of ellipticity test that we have done in this paper is in essence a null test. If we had seen a di erence at high ellipticities, it would have been a surprising result. By seeing none, we are consistent with the standard hierarchical clustering/m erger m odel.

We believe that there is enough evidence now to give a reasonable estimate to the redshift corresponding to a lookback time of  $t_m$ . Pascarelle et al. (1996) have shown that at least in one direction at z=2.39, we still not the subclumps that have generally not yet merged into morphologies recognizable today. If we make the reasonable estimate that

#### { 27 {

the high redshift galaxies in our ellipticity sample are outnum bered by those at  $z \le 1$  (see Figure 2 of Sawicki et al. 1997), we conclude that whatever changes in morphology that have occurred since z = 1 m ust have preferentially occurred among round-appearing galaxies -ie., E, S0 and bulge-dom inated spirals. Mergers of system s we would recognize today as galaxies classi able by the Hubble scheme have therefore been taking place at least as far back as z' = 1, but likely no farther back than z' = 2.4. This would argue that  $1.0^{\circ} z_{t_m} \le 2.5$ , with the most likely time being  $1.5^{\circ} z_{t_m} \le 2.0$ , consistent with the peak in the star-form ation rate in the universe (M adau et al. 1996).

If this view of hierarchical clustering is correct, then understanding how galaxies evolve with time will not be a simple matter of estimating galaxy morphologies and colors at all lookback times. Rather, if what we see at lookback times smaller than  $t_m$ , including the present-day, is the current state of mergers, we will need more detailed information about how groupings of galaxies change with time. That information can only come with measurements of redshifts of very faint galaxies, which is a major challenge that the new generation of 8{10 meter telescopes face if we are to truly understand galaxy evolution.

W e adknow ledge support from HST grants GO .5308.01.93A, GO .5985.01.94A, GO .2684.03.94A, and AR .6385.01.95A. SCO wishes to adknow ledge a very useful conversation with R.B landford and D.Hogg.

#### REFERENCES

Abraham, R., Tanvir, N.R., Santiago, B., Ellis, R.S., Glazebrook, K.G.& van den Bergh, S., 1996, MNRAS, 279, L47

Bertin, E., & A mouts, K., 1996, in preparation

Bohlin, R.C., et al. 1991, ApJ, 368, 12

Brainerd, T., Blandford, R.D., & Smail, I. 1996, ApJ, 466, 623

Binney, J., & de Vaucouleurs, G. 1980, MNRAS, 194, 679

Burg, C., W indhorst, R., O dewahn, S., de Jong, R., & Frogel, J. 1997 in Proceedings of the M aryland W orkshop on \The U ltraviolet Universe at Low and High Redshift: Probing the Progress of Galaxy Evolution", Eds. M. Fanelli, & W. W aller, A IP Conf. Ser., in press

Cowie, L.L., Songaila, A., Hu, E.M., & Cohen, J.G. 1996, AJ, 112, 839

Dalcanton, J., & Shectman, S.A., 1996, ApJ, 465, L9

- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H.G., Buta, R., Paturel, G. & Fougue, P. 1991, The Third Reference Catalog of Bright Galaxies, Springer Verlag: New York, (RC3)
- de Vaucouleurs, G. & Pence 1973, BAAS, 5, 466

de Vaucouleurs, G. 1959, Handbuch der Physik, 53, 311

de Jong, R.S., & van der Kruit, P.C. 1994, A & A S, 106, 451.

- D ressler, A. 1980, ApJ, 236, 351
- Driver, S.P., Windhorst, R.A., Ostrander, E.J., Keel, W.C., Griths, R.E.& Ratnatunga, K.U. 1995a, ApJ, 449, L23 (D 95a)

Driver, S.P., W indhorst, R.A., & Gri ths, R.E. 1995b, ApJ, 453, 48 (D 95b)

Ellis, R., Colless, M., Broadhurst, M., Heyl, J., & Glazebrook, K. 1996, MNRAS, 280, 235

- Fasano, G. & Vio, R. 1991, MNRAS, 249, 629
- Ferguson, H.C. & Sandage, A. 1989, ApJ, 346, L53
- Frei, Z. et al. 1996, AJ, 111, 174

- Giavalisco, M., Livio, M., Bohlin, R., Macchetto, D.F., & Stecher, T.P. 1996, AJ, 112, 364
- G lazebrook, K., Ellis, R.E. Santiago, B., & Gri ths, R.E. 1995, MNRAS, 275, L19
- Gott, J.R., & Rees, M.R. 1975, A&A 15, 235
- Hill, J.K. et al. 1992, ApJ, 395, L37
- Holtzman, J.A., etal. 1995, PASP, 107, 1065
- Hubble, E. 1926, ApJ, 64, 321
- Huchra, J., Geller, M., De Lapparent, V., & Burg, R., 1988, IAU Symposium 130, Large-Scale Structures of the Universe, ed. J. Audouze, M. Pelleto, and A. Szalay, Kluwer, 105
- Im, M. etal. 1995, ApJ, 445, L15
- Kaiser, N. & Squires, G. 1993, ApJ, 404, 441
- Lambas, D.G., Maddox, S.J., & Loveday, J. 1992, MNRAS, 258, 404
- Lilly, S. J., Tresse, L., Hammer, F., Crampton, D., & LeFevre, O. 1995, ApJ, 455, 108
- Madau, P. et al. 1996, MNRAS, 283, 1388
- Marcum, P. et al. 1997 in Proceedings of the Maryland Workshop on \The Ultraviolet Universe at Low and High Redshift: Probing the Progress of Galaxy Evolution", Eds. M. Fanelli, & W. Waller, AIP Conf. Ser., in press
- Mould, J., Blandford, R., Villum sen, J., Brainerd, T., Smail, I., Small, T., & Kells, W. 1994, MNRAS, 271, 31

M utz, S.B., etal. 1994, ApJ, 434, L55

Noerdlinger, A. 1979, ApJ, 234, 802

- O'Connell, R. et al. 1997 in Proceedings of the Maryland Workshop on \The Ultraviolet Universe at Low and High Redshift: Probing the Progress of Galaxy Evolution", Eds. M. Fanelli, & W. Waller, AIP Conf. Ser., in press
- O dewahn, S.C. 1997, Nonlinear Signal and Im age Analysis, Annals of the New York A cademy of Sciences, 188, 184
- Odewahn, S.C., Windhorst, R.A., Driver, S.P., & Keel, W.C. 1996, ApJ, 472, L13
- Odewahn, S.C, & Aldering, G. 1995, AJ, 110, 2009
- Odewahn, S.C. 1995, PASP, 107, 770
- Odewahn, S.C. 1989, PhD. thesis, Univ. of Texas
- Pascarelle, S.M., Windhorst, R.A., Keel, W.C., & Odewahn, S.C. 1996, Nature, 383, 45
- Peebles, P.J.E. 1993, \Principles of Physical Cosm ology", (Princeton University Press), Princeton, New Jersey
- Press, W. H., Flannery, B. P., Teukolsky, S. A., and Vetterling, W. T. 1986, Numerical Recipes, (Cambridge University Press), 472
- Ryden, B.S. & Temdrup, D.M. 1994, ApJ, 425, 43
- Sandage, A., Freem an, K.C., & Stokes, N. 1970, ApJ, 160, 831
- Sawicki, M. J., Lin, H., & Yee, H. 1997, AJ, 113, 1
- Smoot, G.F., Bennett, C.L., Kogut, A., & Wright, E.L., 1992, ApJ, 396, L7
- Tyson, J.A. 1985, Nature, 316, 799
- Tyson, J.A., Valdes, F., & Wenk, R.A. 1990, ApJ, 349, L1
- W illiam s, R E. et al. 1996, AJ, 112, 1335
- W indhorst, R.A., Keel, W.C., & Pascarelle, S.M. 1997, ApJ, submitted

- W indhorst, R.A., Burstein, D., Mathis, D.F., Neuschaefer, L.W., Bertola, F., Buson, L., Matthews, K., Barthel, P.D., & Chambers, K.C. 1991, ApJ, 380, 362
- Weedman, D.W., 1986, in Quasar Astronomy (Cambridge: Cambridge University Press) 71

Fig. 1. Using 176 simulated high z in ages based on observed  $I_{314}$  -band W FPC2 in ages with I 22 (and assumed  $z^{<}$  0.5) and intrinsic axis ratio b=a 0.5, we plot the residual axis ratio, (b=a)<sub>o</sub> (b=a)<sub>z</sub>, as a function of signal-to-noise ratio (log (S=N) in panel a (left) and in age resolution log (R) in panel b (right). Both of these tests predict residuals in the sense expected: faint, low S/N compact in ages are measured to be system atically too round.

Fig. 2. Comparison between the eyeball RC3 b=a values derived from  $\log R_{25}$  and those derived from the autom ated MORPHO package. Shown are 63 galaxies observed in B<sub>J</sub> by Frei et al. (1996; open circles) and 82 galaxies observed in B<sub>CCD</sub> by de Jong (1996; open squares). A linear least squares t to the (eyeball{machine}) residuals gives a slope of 0.21 0.01 and an rm s. scatter about the best t (dashed) line of 0.07. The t was made only with the Frei data which cover an adequate range in b/a, but the de Jong sam ple (by selection has b=a 0.6), were plotted to show consistency with the derived t. Based on this t, we adopted the correction formula in the text (equation 5 to transform the 1569 RC3 (eyeball) axis ratios to the system de ned by the MORPHO autom ated ellipse tting software used to reduce all our W FPC2 faint galaxy in ages.

Fig. 3. The dimensional frequency distributions of apparent axis ratio, b=a, for four  $I_{014}$ -band ranges are plotted using open circles connected by thin dashed lines. The bcalRC3 galaxy b=a distribution is plotted in each panel using lied squares connected by bold lines. The RC3 data were redshifted assuming H<sub>0</sub> = 75 km/sec/M pc, q<sub>0</sub> = 0.5, plus K-corrections and lum inosity evolution estimates discussed in the text. No systematic corrections for R or S/N elects have been applied to the RC3 data in this gure.

Fig. 4. As in Figure 3, but system atic corrections for R esolution e ects have been applied to the RC3 b=a estimates. No statistically signi cant di erence is seen between the RC3 sample moved out of high redshift and the observed W FPC2  $I_{814}$  -band samples, even for galaxies with low b=a values.

Fig. 5. As in Figure 3, but the local Type fractions in the RC 3 for E + S0, Sabc, and Sd+ Irr galaxies have been altered from 32%, 48%, and 18%, respectively, to a spiral-rich sample of 10%, 30%, and 60% at z ' 1. System atic corrections for R e ects have been applied to the RC 3 b=a estimates, as discussed in the text. Note that with these corrections, the higher and lower redshift samples are virtually indistinguishable for  $I^{>} 22 \text{ m ag} (z^{>} 0.5)$ .

Fig. 6. A montage of galaxies with 0.0 b=a 0.4 selected at random from the groundbased and HST galaxy sam ples used in this work. The top row consists of 8 galaxies (observed in  $B_J$ ) by Frei et al. 1996, and the second shows the appearance of these sam e galaxies at z=0.6, as modeled by MORPHO. In the 1st row, in ages are shown at ground-based resolution with each postage stam p in age having a linear size of 167°. In all remaining rows, a W FPC2 resolution (0.0996°/pix) is used with each stam p having a linear dimension of 6.2°. The remaining 4 rows consist of W FPC2 in ages in I<sub>614</sub> roughly corresponding to the fourm agnitude ranges constituting our faint sam ples: 14 I<sub>614</sub> 20, 20 I<sub>614</sub> 22, 22 I<sub>614</sub> 24, 24 I<sub>614</sub> 25. Each in age is norm alized to a linear grey-scale from 3 to +20, where is the noise measured in the local background. M orphological features such as spiral arm s or strong dust lanes are not prom inently seen in the z 0.4 rows of W FPC2 in ages, although such features are still clearly visible in the B<sub>J</sub> in ages m odeled for z = 0.6. The high redshift system s (z 0.8) do appear to display a higher degree of in age asymmetry along the m a pr axis and m ore clum piness relative to the lower redshift system s.

Fig. 7. As in Figure 6, but for galaxies with 0:6 b=a 0:75. As in the b=a 0.4 cases of Figure 6, it appears that prom inent spiral arm s are not seen in the z 0:8 rows of W FPC2 in ages, and the degree of in age clum piness and asymmetry is higher compared to their low redshift counterparts.

Fig. 8. As in Figure 6, but for galaxies with 0.8 b=a 1.0. Again no prominent spiral structure is visible for z 0.8, but now the in pression of increased global in age asymmetry is not as strong for the rounder galaxies.

{ 34 {

N am e	RA (2000) DEC	l <b>,</b> b	AB	T (sec)	N <sub>im ages</sub>
HDF	12:36:49.59 + 62:12:54.0	125.9 + 54.8	0.00	123600	49
53W 002	17:14:15:47 + 50:15:28:3	77.0 + 35.8	0.03	1700	17
HFF_OW	12:36:21.16 + 62:14:27.2	126.0 + 54.8	0.00	2500	3
HFF_SW	12:36:33.91 + 62:10:54.2	125.9 + 54.9	0.00	2500	3
HFF_IW	12:36:35.77 + 62:13:40.9	125.9 + 54.8	0.00	5300	4
HFF_SE	12:36:47.77 + 62:10:19.3	125.9 + 54.9	0.00	2500	3
HFF_NW	12:36:50.81 + 62:15:41.9	125.9 + 54.8	0.00	2500	3
HFF_IE	12:37:03.86 + 62:12:15.2	125.8 + 54.8	0.00	5300	4
HFF_NE	12:37:05.85 + 62:15:06.7	125.8 + 54.8	0.00	2500	3
HFF_OE	12:37:17.92 + 62:11:33.4	125.8 + 54.9	0.00	3000	3
DWG_UEH0	00:53:23:57 + 12:33:50.3	123.7 -50.3	0.18	2100	3
DWG_UM0	03:55:31.91 + 09:43:38.5	179.8-32.1	0.46	866	6
DW G_UOP0	07:50:47.39 + 14:40:42.7	206.1 + 19.6	0.06	2100	2
DWG_UY40	14:34:48.54 + 25:08:11.1	033.9 + 66.8	0.04	1000	6
DWG_UX40	15:19:41:22 + 23:52:04.3	035.8 + 56.5	0.14	1875	4
DW G_USA2	17:12:23:17 + 33:35:48:5	056.7 + 34.3	0.12	2100	3

Table 1. W FPC2 Fields A nalyzed in F814W

Ξ

\_







Fig. 3



Fig. 4



Fig. 5

Frei Bj z ~ 0.007



Frei Bj z ~ 0.6



 $14 < I(F814W) < 20 z \sim 0.1$ 



 $20 < I(F814W) < 22 z \sim 0.4$ 



 $22 < I(F814W) < 24 z \sim 0.8$ 



 $24 \le I(F814W) \le 25 z \sim 1.1$ 



Galaxies with 0.0 < B/A < 0.4

Frei Bj z ~ 0.007



Frei Bj z ~ 0.6



 $14 < I(F814W) < 20 z \sim 0.1$ 



20 < I(F814W) < 22 z ~ 0.4



 $22 < I(F814W) < 24 z \sim 0.8$ 



24 < I(F814W) < 25 z ~ 1.1



Galaxies with 0.6 < B/A < 0.75

Frei Bj z ~ 0.007



Frei Bj z~0.6



 $14 \le I(F814W) \le 20 \ z \sim 0.1$ 



 $22 < I(F814W) < 20 z \sim 0.4$ 



 $22 < I(F814W) < 24 z \sim 0.8$ 



24 < I(F814W) < 25 z ~ 1.1



Galaxies with 0.8 < B/A < 1.0