# Them alRelaxation in One-D imensional Self-Gravitating Systems

Junichiro Makino
Department of Systems Science, College of Arts and Sciences,
University of Tokyo, Tokyo 153-8902

January 21, 2022

#### A bstract

In this paper, we study the therm alrelaxation in the one-dim ensional self-gravitating system , or the so-called sheet m odel. A coording to the standard argument, the therm al relaxation time of the system is around N  $t_{\rm c}$ , where N is the number of sheets and  $t_{\rm c}$  is the crossing time. It has been claimed that the system does not reach the thermal equilibrium in this thermal relaxation timescale, and that it takes much longer time for the system to reach true thermal equilibrium.

We demonstrate that this behavior is explained simply by the fact that the relaxation time is long. The relaxation time of sheets with average binding energy is 20N t<sub>c</sub>, and that of sheets with high energy can exceed 1000N t<sub>c</sub>. Thus, one needs to take the average over the relaxation time scale of high-energy sheets, if one wants to look at the thermal characteristic of these high energy sheets.

#### 1 Introduction

The one-dim ensional self-gravitating m any-body system was originally discussed mainly as a simple toy model to understand the violent relaxation, [4] because the thermal relaxation timescale of its discrete realization, the sheet model, was believed to be long. Until 1980s, it had been generally accepted that the thermal relaxation time of the system of N equal-mass sheets is of the order of N  $^2$ t<sub>c</sub>, where t<sub>c</sub> is the crossing time of the system.

However, by means of numerical simulation Luwel et al.[3] have demonstrated that the relaxation time is of the order of N  $t_c$ . Reideland Miller [6, 7] reached a similar conclusion, though they reported the presence of systems which apparently did not relax for much longer timescale.

In a series of papers, T suchiya et al. [10, 11, 12] have studied the therm alrelaxation process of one-dimensional self-gravitating systems in detail, by means of the numerical integration over very long time scale (some of their experiments covered 5  $10^8 t_c$ ). They claimed that the thermal relaxation of the sheet model proceeds in a highly complex manner. In the \microscopic relaxation timescale" of N  $t_c$ , each sheet forgets its initial condition, and the system is well mixed. However, according to them, the system does not really reach the thermal equilibrium in this timescale, and the distribution function remains different from that of the isothermal state. They called this state a quasiequilibrium

By pursuing the time integration for much longer timescale, T suchiya et al. [12] found that the system exhibits the transition from one quasiequilibrium to another, and they claimed that the thermal equilibrium is only realized by averaging over the timescale longer

than the tim escale of these transitions. Thus, they argued that there exists the tim escale for  $\mbox{m}$  acroscopic" relaxation, which is much longer than the usual therm alrelaxation (what they called  $\mbox{m}$  icroscopic relaxation").

In this paper, we try to exam ine the nature of this \m acroscopic" relaxation of the one-dimensional sheet model. In section 2, we describe the numerical model. In section 3, we present the result of the measurement of the relaxation time. It is shown that the relaxation time, dened as the timescale in which individual sheets change their energies, depends very strongly on the energy itself, and is very long for high energy sheets. This strong dependence of the relaxation timescale on the energy naturally explains the apparent \transient" phenomena observed by T suchiya et al.[12] Section 4 discusses the implication and relevance of our results.

### 2 The Model

#### 2.1 Sheet model

The Ham iltonian of the sheet model is given by

$$H = \frac{m}{2} \sum_{i=1}^{X^{N}} v_{i}^{2} + 2 G m^{2} \sum_{i < j} \dot{x}_{i} \quad x_{j} \dot{j}$$
 (1)

where  $x_i$  and  $v_i$  are the position and velocity of sheet i, m is the mass of the sheets, N is the number of the sheets and G is the gravitational constant. The crossing time is dened as

$$t_c = \frac{1}{4 \text{ GM}} \frac{\frac{4E}{4E}}{M}; \qquad (2)$$

where M = m N is the total mass of the system. Following T suchiya et al.[12] and others, we use the system of units in which M = 4E = 4 G = 1. In this system,  $t_c = 1$ .

A unique nature of the one-dimensional gravitational system is that there exists the therm all equilibrium, unlike its counterpart in three dimensions. Rybicki [8] obtained the distribution function

$$f(") = \frac{1}{8} \frac{1}{2} \frac{1}{2} \frac{3M}{2E} \exp \frac{3M}{2E}"$$
; (3)

where " is the speci c binding energy de ned as

$$" = \frac{v^2}{2} + (x)$$
 (0):

Here, (x) is the specie control energy. This distribution function satisfies the relation

$$\exp \quad \frac{3M}{2E} " = \operatorname{sech}^2 \frac{3x}{8E} : \tag{5}$$

We perform ed the time integration of the system with N = 16, 32, 64, 128 and 256. For all systems, the initial condition is a water-bag with the aspect ratio  $x_{m ax} = v_{m ax} = 2.5$ .

### 2.2 Num erical m ethod

The important character of the sheet model is that one can calculate the exact orbit of each sheet until two sheets cross each other. Thus, we can integrate the evolution of the system precisely (except for the round-o error). This may sound like a great advantage, compared

to the system s in higher dimensions whose orbits can be calculated only numerically. Instead of numerically integrating the orbit of each sheet, we can calculate the exact orbit for any sheet, until it collides with the neighboring sheet. Thus, by arranging the pairs using heap, we can handle each collision in log N calculation cost.

Note, however, that typically each sheet collides with all other sheets in one crossing time. Thus, the calculation cost is O (N  $^2$  log N ) per crossing time. Our simulation with N = 64 for 2  $10^7 t_{\rm c}$  took 8 hours on a VT-A lpha workstation with DEC A lpha 21164A CPU running at 533 MHz. For this run, the total energy of the system was conserved better than 3  $10^{\,12}$ .

### 3 Results

### 3.1 Approach to the therm alequilibrium

Figure 1 shows the time-averaged energy distribution function N ("), for different time periods and number of sheets. In all gures, the thin solid curve is the energy distribution of the isothermal distribution function of equation (3). What we see is quite clear. As we make the time interval longer, the time-averaged distribution function approaches to the isothermal distribution. Thus, the numerical result suggests the system is ergodic. However, it also shows that the time needed to populate the high-energy region is very long. The sampling time interval is 128 time units for N = 16, and 512 time units for N = 64 and 128. Thus, in the case of T =  $2^{18}$  and N = 16 (dash-dotted curve in gure 1a), total number of sample points is  $2^{15} = 32768$ .

If we can assume that the sample points are uncorrelated, the possibility that no sample exceeds energy level  $\mathbf{v}_0$  is given simply by

$$P("_0;N) = [1 P(" < "_0)]^n;$$
 (6)

w here

$$P (" < "_0) = \sum_{0}^{Z} N (")d";$$
 (7)

and n is the number of sample points. Figure 2 shows 1 P (") as a function of ". For " = 1.25, P (") = 0.996, and therefore the probability that none of 32768 samples does not exceed " = 1.25 is practically zero (< e  $^{100}$ ). In other words, the numerical result seems to suggest that the system is not in the thermally relaxed state even after 2  $^{105}$  crossing times.

Of course, this result is not surprising if the relaxation time is long. Samples taken with the time interval shorter than the relaxation time have a strong correlation, and therefore the elective number of freedom can be smaller than n.Roughly speaking, if the relaxation time is longer than  $10^4$ , our numerical result is consistent with the assumption that the system is in the thermal equilibrium. In the next subsection, we investigate the relaxation time itself.

## 3.2 Relaxation timescale

W e m easured the following quantities:

$$D_{1} = \frac{\langle "_{1}(t_{0}) \ "_{1}(t_{0} + t) \rangle}{t};$$
 (8)

$$D_{2} = \frac{\langle [\mathbf{I}_{1}(t_{0}) \quad \mathbf{I}_{1}(t_{0} + t)]^{2} \rangle}{t} :$$
 (9)




Figure 1: The time-averaged distribution function in the energy space N (E); (a) N = 16, (b) N = 64, (c) N = 128.




Figure 2: The compliment of the cumulative distribution function  $1\ P$  (e) for the thermal equilibrium.

These quantities correspond to the coe cients of the rst and second-order terms in the Fokker-Planck equation for the distribution function, and have been used as the measure of the relaxation in many studies (see, e.g., Hemquist and Bames, [1] Hemquist et al. [2]), for three-dimensional systems. However, to our know ledge this measure has not been used for the study of the sheet model.

In order to see the dependence of these di usion coe cients on the energy, we calculated them for intervals of "= 0:15. Figure 3 shows the results, for N = 16;64 and 256. The time interval t was taken equal to N  $t_c$ . We used smaller values for t and con rmed that the choice of thas negligible e ect if t is larger than 10 $t_c$  and smaller than 4N  $t_c$ . T im e average is taken over the whole simulation period. We can see that both the rstand second-order terms show very strong dependence on the energy of the sheets, and of the order of 1=100N for e 1:5. Figure 3 suggests that the relaxation timescale grows exponentially as energy grows. This behavior is independent of the value of N .

We can de nethe relaxation timescale as

$$t_r = ^{n^2} = D_2; \tag{10}$$

that is, the time scale in which energy changes signicantly. Figure 4 shows this relaxation time scale for dierent values of N and ". The relaxation time shows very strong dependence on the energy and the relaxation of high-energy sheets is much slower than that of sheets in lower energies. This is partly because of the dependence of  $t_r$  on "itself. However, as we can see in gure 3, the dependence of the diesion coes cient is the main reason.

This result resolves the apparent contradiction between the fact that the relaxation timescale is of the order of N  $t_c$  [5] and that the system reaches the true thermal equilibrium only in much longer timescale.[12] It is true that the relaxation timescale is O (N), but the coecient before N is quite large, in particular for sheets with high energies.

An important question is why the relaxation timescale depends so strongly on the energy. This is provably due to the fact that high-energy sheets have the orbital period signicantly longer than the crossing time. Typical sheets have the period comparable to the crossing time, and therefore they are in strong resonance with each other. However, a high-energy sheet has the period longer than the crossing time, and thus it is out of resonance with the




Figure 3: The di usion coe cients (a)  $D_1$  and (b)  $D_2$  plotted against the energy e for three values of N. Long-dashed, solid, and short-dashed curves are the results for N=16, 64 and 256, respectively.




Figure 4: The relaxation time in unit of N  $t_c$  plotted against the energy e for three values of N . Curves have the same meanings as in gure 3

rest of the system . Therefore, the coupling between high energy sheets and the rest of the system is much weaker than the coupling between sheets with average energy. This explains why the relaxation of high energy sheets is slow.

# 4 Summary and Discussion

In this paper, we studied the therm alrelaxation process in one-dimensional self-gravitating systems. We can med the result obtained by T suchiya et al.[12] that the thermal relaxation takes place in the timescale much longer than N t<sub>c</sub>. However, we found that this is simply because the thermal relaxation timescale is much longer than N t<sub>c</sub>. Even for typical sheets, the relaxation timescale is around 10N t<sub>c</sub>. In order to obtain good statistics, we need to take average over many relaxation times. Moreover, the relaxation time for sheets in the high-energy end of the distribution function is even longer, since the relaxation timescale grows exponentially as the energy grows. Thus, it is not surprising that we have to wait for more than  $10^4$ N t<sub>c</sub> to obtain good statistics.

Does this inding have any theoretical/practical relevance? Theoretically, there is nothing new in our result. What we found is simply that numerical simulation should cover the period much longer than the relaxation timescale to obtain statistical properties of the system, and that the relaxation timescale of a sheet depends on its energy. Both are obvious, but some of the previous studies neglected one or both of the above, and claimed to have found a complex behavior, which, in our view, is just a random walk.

Our nding of the long relaxation time by itself has rather little astrophysical signicance, since in the large N limit, the relaxation time is in nite anyway. However, since any numerical simulation su ers some form of numerical relaxation, it is rather important to understand how the relaxation elect changes the system. To illustrate this, we exam in the claims by T suchiya et al.[12] in some detail here.

They argued that the evolution of the mass sheet model proceeds in the following four steps: (1) viliarization, (2) dynamical equilibrium, (3) quasiequilibrium, and (4) them all equilibrium. A coording to them, the viliarization timescale is order of  $t_c$ , and the energy of each sheet is \conserved" in the dynamical equilibrium phase, which continues up to to N  $t_c$ . Then, \microscopic relaxation" takes place in the timescale of to N  $t_c$ , where the energy of each sheet is relaxed, but the whole system needs timescale much longer to reach the true equilibrium, because of some complex structure in the phase space.

Our num erical results are in good agreement with those of T suchiya et al.,[12] but our interpretation is much simpler: First system virializes, and then relaxation proceeds in the timescale of thermal relaxation, which depends on the energy of the individual sheets. Thus, the central region with short relaxation time relaxes to the distribution close to the thermal relaxation in less than 100N t<sub>c</sub>, but the distribution in the high-energy tail takes much longer to settle. In addition, the small number statistics in the high-energy region makes it necessary to average over many relaxation times to obtain good statistics. In other words, there are no distinction between the \microscopic" and \macroscopic" relaxation, and the evolution of the system is perfectly understood in terms of the standard thermal relaxation.

## A cknow ledgm ents

I would like to thank Yoko Funato, Toshiyuki Fukushige and Daiichiro Sugim oto for stimulating discussions, and Shunsuke Hozum i for comments on the manuscript. This work is supported in part by the Research for the Future Program of Japan Society for the Promotion of Science (JSPS-RFTP97P01102).

# R eferences

- [1] L.Hemquist and J.E.Bames, Astrophys. J. 349 (1990), 562.
- [2] L. Hemquist, P. Hut and J. Makino, Astrophys. J. Lett. 411 (1993), 248.
- [3] M. Luwel, G. Severne, and P. J. Rousseeuw, Astrophys. Space Sci. 100 (1984), 261.
- [4] D. Lynden-Bell, Mon. Not. Roy. Astron. Soc 136 (1967), 101.
- [5] G. Severne, M. Luwel, and P. J. Rousseeuw, Astron. Astrophys. 138 (1984), 365.
- [6] C.J.Reidl, Jr. and B.N.M iller, Astrophys. J. 318 (1987), 248.
- [7] C.J.Reidl, Jr. and B.N.M iller, Astrophys. J. 332, (1988) 619.
- [8] G.B.Rybicki, Astrophys. Space Sci. 14, (1971) 56.
- [9] L.Spitzer and M.H.Hart, Astrophys. J. 164, (1971) 399.
- [10] T. Tsuchiya, T. Konishi, and N. Gouda, Phys. Rev. E 50 (1994), 2607.
- [11] T. Tsuchiya, N. Gouda, and T. Konishi, Phys. Rev. E 53 (1996), 2210.
- [12] T. Tsuchiya, N. Gouda, and T. Konishi, Ap. SS. 257 (1997), 319.