Line Strengths and Line Strength G radients in Bulges along the Hubble Sequence

PaulG oudfrooij

Space Telescope Science Institute, 3700 San M artin D rive, Baltim ore, M D 21218, U nited States of Am erica (goudfroo@stsci.edu)

A $\;$ liated with the A strophysics D ivision, SSD , European Space Agency $\;$

Javier Gorgas

Departamento de Astrosica, Facultad de Ciencias Fisicas, Universidad Complutense de Madrid, E-28040 Madrid, Spain (fjg@astrax.fis.ucm.es)

Pascale Jablonka

DAEC (URA 173, O bærvatoire de Paris-M eudon, P lace Jules Janssen, M eudon, F-92195, France (pascale.jablonka@obspm.fr)

A bstract. We present rst results of a comprehensive survey of deep long-slit spectra along the m inor axis of bulges of edge-on spiral galaxies. Our results indicate that stellar populations in bulges are fairly old and encompass a range of m etallicities. The lum inosity-weighted ages of bulges range from those found for cluster ellipticals to slightly \younger" (by up to only a few Gyr, how ever). Their /Fe elem ent ratio is typically supersolar, consistent with those found in giant ellipticals. The radial line-strength gradients in bulges correlate with bulge lum inosity. Generally, these ndings are more compatible with predictions of the \dissipative collapse" m odel than with those of the \secular evolution" m odel.

K eyw ords: Bulges of Spiral G alaxies, Stellar Populations, R adial G radients

1. Introduction: The Form ation of Bulges of Spirals

Bulges of spiral galaxies are connerstones for constraining theories of galaxy form ation. Located at the centers of spiral galaxies, they hold the signature of the sequence of form ation | outside-in or inside-out| of the di erent sub-system s building a spiral galaxy: halo, disc, and bulge. The prom inence of bulges varies widely along the Hubble sequence. This contrasts with the situation for spiral disks whose m ass is nearly constant am ong all types of spirals (A rim oto & Jablonka, 1992). Hence, bulges constitute a main key to our understanding of spiral galaxy evolution. To set the context for our project, we describe below the two currently m ost popular scenarios on bulge form ation that have been proposed over the years.

Based on observations obtained at ESO, La Silla, Chile (O bserving P rogram m es 58 A-0192, 59 A-0774, and 61 A-0326), and at the Isaac N ew ton T elescope operated on the island of La P alm a by the Isaac N ew ton G roup in the Spanish O bservatorio delR oque de los M uchachos of the Instituto de A stro sica de C anarias



M onolithic D issipative C ollapse: Bulges form before disks do, on very short time scales. As the pregalactic gas collapses and form s stars, m etals are ejected by winds blown by m assive stars and supernovae. If the galaxy potential well is deep enough to retain these ejecta, the enriched gas is carried inward to the galaxy center. As new stars are form ed, their chem ical composition re ects that of the gas. The result is a radial m etallicity gradient, whose am plitude should increase with galaxy m ass and lum inosity (C arlberg, 1984; A rim oto & Yoshii, 1987).

Evidence in favor of this scenario has been presented by m eans of spectroscopic studies of ellipticals: Carollo, Danziger & Buson (1993) reported the presence of a correlation between radial M g_2 absorptionline-strength gradients and galaxy mass, velocity dispersion, and central line strength for low { to interm ediate-m ass (i.e., M $\leq 10^{11}$ M) ellipticals, as opposed to giant ellipticals (but see di erent results by Gonzalez & Gorgas 1996 and Kobayashi & Arim oto 1999). The gradients may thus be driven mostly by dissipation in smallish ellipticals, whereasm erging and violent relaxation play a more important role for giant ellipticals. W hat about the actual case of bulges of spirals? There is strong evidence that bulges are analogous to low { to interm ediatem assellipticals in several global principal properties: they populate the same location in the Fundam ental Plane (Bender, Burstein & Faber, 1992); they form a continuous sequence in the $V_{max} = 0$ vs. ellipticity diagram (being supported by rotation, Bender et al. 1992). As to the stellar populations, Jablonka, Martin & Arim oto (1996) performed a spectroscopic pilot study of the centers of bulges of face-on spirals. They revealed the existence of striking sim ilarities between bulges and ellipticals. E.g., their lum inosity-m etallicity relations, when derived from -elements (such as the $M g_2$ index), are consistent with one another.

Secular Evolution: Bulges form from disc material through redistribution of angular momentum. In this scenario, large amounts of gas are driven into the central region of the galaxy by a stellar bar and trigger intense star form ation (e.g., P fenniger & Norm an 1990). If enough mass is accreted, the bar itself will be dissolved and the resulting galaxy will reveal a bigger bulge than before bar form ation; galaxies would thus evolve from late to earlier types along the Hubble sequence.

Thism odel also certainly has its attractions. Bars appear in at least

2/3 of disk galaxies (Sellwood & W ilkinson, 1993), and several num erical simulations have indicated that bars can signi cantly in uence the dynam ical evolution of galaxies, through m echanism s such as disk thickening by box-peanut or bending instabilities (C om bes et al., 1990) or radial mass in ow towards the center (Friedli & Benz, 1995). Since bars appear to be able to a lect the global dynam ics of galaxies, it is natural to suspect that bars can be responsible for signi cant chem - ical evolution as well, due to mixing by the bar-induced kinematics. Recent N-body simulations (that include the e ect of star formation) have shown this to be a valid suspicion in the sense that any initial abundance gradient is indeed signi cantly washed out 1 Gyr after formation of a bar, both for gas and stars (Friedli, Benz & Kennicutt, 1994; Friedli, 1998). The slope of the abundance gradient is found to

atten beyond the co-rotation radius, by up to 50% through the e ect of one (strong) bar. O bservational evidence for this e ect exists, albeit only for the gas component so far. Several groups have shown that global radial gradients of the gas m etallicity (in terms of [D/H] in H ii regions) in barred spirals are shallower than gradients in \norm al" spirals of the same H ubble type (V ila-C ostas & Edm unds, 1992; Z aritsky et al., 1994; M artin & Roy, 1994). Furtherm ore, a clear relation seems s to exist between the relative length of the bar with respect to the size of the (optical) disk and the slope of the radial [D/H] gradient am ong barred spirals (M artin & Roy, 1994) in the sense that the larger the bar/disk ratio, the shallower the abundance gradient.

1.1. Line-Strength Gradients: Key to the Controversy? It thus seems that radial population gradients in bulges have great potential in discrim inating between the main bulge form ation scenarios. Previous work on stellar populations in bulges has mainly been limited to photometric studies: Balcells & Peletier (1994) in UBRI, and Terndrup et al. (1994) in J and K, show that for lum inous bulges, color gradients become increasingly negative with increasing lum inosity (sim ilar to the case am ong ellipticals). However, many faint bulges deviate from this trend and keep showing strong negative color gradients, and the two studies mentioned above do not converge in their conclusions. Besides, it is now well known that broad-band photom etry is incapable of accurately disentangling the e ects of age, m etallicity, and dust absorption, whereas stellar line strenghts are dust-independent and can separate the e ects of age and m etallicity (e.g., W orthey, 1994, hereafter W 94; Vazdekis et al., 1996). W ith this in m ind, we em barked on a spectroscopic survey of a signi cant sample of bulges to compare their possible star form ation histories with those in ellipticals.

2. Description of our Project

In order to avoid contam ination from disk light, we selected a sam ple of 28 edge-on, nearby spiral galaxies from the UGC and ESO-LV catalogs, and oriented the spectrograph slit along the m inor axis of the bulges. The sam ple encom passes a considerable range in lum inosities for each Hubble type (18:4 < M_V < 21:5). The observations (spectroscopy and two-color in aging) of the northerm spirals were obtained with the

2.5-m Isaac New ton Telescope at La Palma, while the southern spirals were observed with the ESO NTT and 3.6-m telescopes. As to the spectroscopy, the typical exposure time was 4 hours per galaxy, and the instrum ental resolution was typically of order 100 km s¹. The spectral range used (typically 3900 { 5500A) allowed the measurem ent of m ost Lick/ID S line-strength indices.

All spectral data reduction and line index measurements were performed using the $R_E D_m^{UCE}$ package¹ which propagates errors associated with all reduction steps along with the science data. The line indices were fully calibrated to the Lick/DS system (including broadening corrections), using measurements of 40 DS standard stars (G orgas et al., 1993). Full details of the sample selection, the observations, and the calibrations will be provided in a forthcoming paper.

3. Results

At the time this paper is written, data analysis of 16 bulges in our sample is completed. Here we describe some highlights of the results from this subsample.

3.1. Central Line Strengths

In order to e ectively compare \central" line strengths for bulges with those of ellipticals in the literature, we extracted spectra with spatial extent 4^{00} from the galaxy center but avoiding the innerm ost dust lane to elim inate disk light. In disentangling the e ects of age and m etallicity, m ost authors have used a combination of H and \m etallic" line-strengths such as M g₂, M gb, Fe5270 and Fe5335 (e.g., G onz alez 1993). However, the use of the H and M gb indices is very limited in case nebular emission is present (G oudfrooij & Em sellem, 1996), which is \unfortunately" often the case in our spectra. W e did not attempt to correct H for emission, but rather used the H A index as age-sensitive index (calibrated in age/m etallicity space by W orthey & O ttaviani 1997). Since the relative strength of B alm er line emission decreases rapidly with B alm er order (e.g., O sterbrock 1974), H A is signi cantly less diluted by nebular emission than H is.

In Fig.1 we present plots of the H $_{\rm A}$ index vs.two m etallicity indicators: (i) C₂4668 which Jones & W orthey (1995) identi ed as a particularly m etallicity-sensitive index, and (ii) < Fe> = (Fe5270+Fe5335)/2. For comparison, we also plot the results of K untschner & D avies (1998; hereafter KD 98) for the centers of E and S0 galaxies in the Formax cluster, as well as predictions from single-burst population synthesis m odels (W 94; W orthey & Ottaviani1997). It turns out that the bulges

4

¹ Cardiel & Gorgas, http://www.ucm.es/info/Astrof/reduceme/reduceme.html



Figure 1. The age-sensitive index H $_{\rm A}$ is plotted against metallicity indicators C₂4668 (left) and < Fe> (right) for the central regions of bulges in our sample. Age-metallicity grids from population synthesis models by W orthey (1994) and W orthey & Ottaviani (1997) are overplotted. Solid lines represent constant age, while dashed lines represent constant metallicity. Sym bol de nitions are shown in the inset; sym bol size is proportional to the bulge lum inosity. O pen circles and open squares represent centers of Formax ellipticals and S0s, respectively, from KD 98.

in our sam ple have ages similar to (or up to a few Gyr \younger"² than) those of cluster ellipticals. On the other hand, lum inosity-weighted ages of a sam ple biased towards eld ellipticals (G onz alez 1993) span the whole range found for bulges. The m etallicities of bulges cover a range similar to those of ellipticals. Interestingly, bulges of later-type (Sb { Sc) spirals are, in the mean, less metal rich than their counterparts in earlier-types. This seem s to be a bulge lum inosity e ect, judging from the symbol sizes in Fig. 1. There are no other obvious distinctions between bulges of di erent Hubble types in this context.

A nother interesting point is that, comparing the positions of the galaxies in Figs. 1a and 1b, it appears that C_2 (i.e., the C_24668 index) is overabundant in ellipticals and bulges with respect to solar abundance ratios. This is illustrated in Fig. 2a which compares C_24668 with < Fe>. C om paring the observations with the superim posed models of W 94, it is obvious that C_24668 is stronger than indicated by the models (which employed solar abundance ratios). Whether this is a real overabundance e ect (or, e.g., due to a problem in the tting functions) is an issue which deserves further analysis.

 $^{^2}$ Recall however that line-strength indices re ect lum inosity-weighted properties in a galaxy. A young population that is small in mass \mid but relatively large in lum inosity \mid , can dram atically change the index values.



Figure 2. (a) $C_24668 \text{ vs.} < \text{Fe} > \text{ indices for the central regions of bulges in our sample. M odels by W orthey (1994) are overplotted as in Fig. 1. Sym bols are shown in the inset. The sym bol size is proportional to the bulge lum inosity of the galaxies. Open pentagons represent Formax ellipticals from K untschner & D avies (1998). (b) M g₂ vs. < Fe> indices for bulges in our sample. Sym bols as in Fig. 2a. The led sym bols represent the \central" indices of the bulges, and the arrow s pointing dow nw ards from those sym bols denote the radial (vertical) gradients going outwards (length of arrow is proportional to the gradient slope). O verplotted are m odels by W orthey (1994) [only for ages 3 and 12 G yr; solid lines] and a correction for M g/Fe] = 0.45 for the 12 G yr isoage line (calculated from m odels by W eiss et al. 1995).$

3.2. Radial Line-strength Gradients

Fig. 2b shows a < Fe> vs. M g₂ plot for the bulges in our sample. The sym bols depict the \central" values, while the arrow spoint tow ands the outerm ost well-measured values; the length of the arrow s is a measure of the slope of the gradient. It is clear that, in most bulges, M g is overabundant with respect to solar abundance ratios. The [M g/Fe] abundance ratio in bulges is similar to those found in ellipticals, and staysm ore or less constant throughout the radial extent of bulges. From the (overplotted) -elem ent-overabundance models by W eiss, P eletier & M atteucci (1995) for [M g/Fe] = 0.45 and an age of 12 G yr (and mixing length parameter mult = 1.5), we estim ate that [M g/Fe] < 0.4.

Is there a correlation between the m etallicity gradient and lum inosity for bulges, as predicted by dissipative collapse m odels? This relation is depicted in Fig. 3. Bulge lum inosities³ were derived by performing ellipse ts to the isophotes of the galaxy in ages, after m asking out wedges encom passing the dusty disks (20 from the major axes). The gradient-lum inosity correlation indeed exists; any further distinction

³ using H₀ = 50 km s¹ Mpc¹



Figure 3. Logarithm ic radial gradient of M g_2 index vs. absolute V-band bulge lum inosity of the bulges analyzed to date. Symbols are shown in the inset (where T is the RC3 m orphological type). The dotted lines depict the extremes of the M $g_2/\log(r)$ vs.M $_V$ relation among low { and intermediate-lum inosity E and S0 galaxies from C arolb et al. (1993).

between bulges of di erent Hubble types within this relation is not obvious. Incidentally, the M g_2 gradients of the m ost lum inous bulges as well as the slope of the gradient-lum inosity relation are sim ilar to those am ong low -lum inosity ellipticals in the C arollo et al. (1993) sample.

4. Concluding Remarks

From the spectral data of bulges of spirals in our sample analyzed so far, we have established the following main results:

- 1. Bulges have lum inosity-weighted metallicities varying from roughly 0.50 to +0.20 in [Fe/H], as measured from H_A vs. < Fe> index diagram s. M any bulges are as old as cluster ellipticals, but som e (low -lum inosity) bulges have lum inosity-weighted ages up to a few G yr younger (note that this result m ay be in uenced by residual light from thick disks).
- 2. Bulges are typically overabundant in -elements, up to Mg/Fe] ' + 0.4 dex, throughout their radial extent. There is no obvious correlation between Mg/Fe] and bulge lum inosity. This is sim ilar to the situation among ellipticals, and indicates that the bulk of the stars in bulges typically form ed within a few Gyr (before the onset of SN Ia explosions; e.g., W orthey, Faber & Gonzalez 1992).
- 3. There is a correlation between the radialm etal-line index gradients and the bulge lum inosities for bulges in our sample.

These rst results seem to be generally more compatible with the predictions of the \dissipative collapse" models than with those of the \secular evolution" models (cf. Sect. 1). However, some individual bulges seem to be younger than the rest and show shallow radial M g_2 gradients, which in turn can be due to e ects induced by bar instabilities. Moreover, we postpone the announcement of $\$ nal" conclusions until the data of our full sample of 28 bulges has been analyzed.

A cknow ledgem ents. We thank the referee, R ichard de Grijs, for his timely and critical review of this paper. We are grateful to N icolas Cardiel for his help with the $R_E D \stackrel{uc}{m} E$ package, and to H arold K untschner for giving access to his data in electronic form.

References

Arim oto, N., Yoshii, Y., 1987, A&A, 173, 23 Arim oto, N., Jablonka, P., 1992, A & A, 249, 374 Balcells, M ., Peletier, R F ., 1994, A J, 107, 135 Bender, R., Burstein, D., Faber, S.M., 1992, ApJ, 399, 462 Carlberg, R.G., 1984, ApJ, 286, 403 Carollo, C.M., Danziger, I.J., Buson, L.M., 1993, MNRAS, 265, 553 Combes, F., Debbasch, F., Friedli, D., Pfenniger, D., 1990, A&A, 233, 82 Friedli, D., 1998, in: D. Friedliet al. (eds.). A SP Conf. Ser. 147, Abundance Pro les: D iagnostic Tools for Galaxy History. ASP, San Francisco, p. 287 Friedli, D., Benz, W., 1995, A&A, 310, 649 Friedli, D., Benz, W., Kennicutt, R., 1994, ApJ, 430, L105 Gonzalez, J.J., 1993, Ph.D. thesis, University of California, Santa Cruz Gonzalez, J. J., Gorgas, J., 1996, in: A. Buzzoni, A. Renzini, A. Serrano (eds.). A SP Conf. Ser. 86, Fresh V iews on Elliptical Galaxies. A SP, San Francisco, p. 225 Gorgas, J., Faber, S.M., Burstein, D., et al., 1993, ApJS, 86, 153 Goudfrooij, P., Em sellem, E., 1996, A & A, 306, L45 Jablonka, P., Martin, P., Arim oto, N., 1996, AJ, 112, 1415 Jones, L.A., W orthey, G., 1995, ApJ, 446, L31 Kobayashi, C., Arimoto, N., 1999, ApJ, in press (astro-ph/9907091) Kuntschner, H., Davies, R.L., 1998, MNRAS, 295, L39 (KD 98) Martin, P., Roy, J.-R., 1994, ApJ, 426, 601 Osterbrock, D.E., 1974, Astrophysics of Gaseous Nebulae. W.H.Freeman & Co., San Francisco Pfenniger, D., Norman, C., 1990, ApJ, 363, 391 Sellwood, J.A., Wilkinson, A., 1993, Rep. Prog. Phys., 56, 173 Temdrup, D.M., Davies, R.L., Frogel, J.A., et al., 1994, ApJ, 432, 518 Vila-Costas, M.B., Edm unds, M.G., 1992, MNRAS, 259, 121 Vazdekis, A., Casuso, E., Peletier, R.F., Beckman, J.E., 1996, ApJS, 106, 307 W eiss, A., Peletier, R.F., M atteucci, F., 1995, A&A, 296, 73 W orthey, G., 1994, ApJS, 95, 107 (W 94) W orthey, G., Faber, S.M., Gonzalez, J.J., 1992, ApJ, 398, 69 W orthey, G., Ottoviani, D.L., 1997, ApJS, 111, 377 Zaritsky, D., Kennicutt, R.C., Huchra, J.P., 1994, ApJ, 420, 87

8