

Real Arnold complexity versus real topological entropy for birational transformations

N. Abarenkova

Centre de Recherches sur les Très Basses Températures, B.P. 166, F-38042 Grenoble, France
Theoretical Physics Department, Sankt Petersburg State University, Ulyanovskaya 1, 198904 Sankt Petersburg, Russia

J.-Ch. Anglès d'Auriac*

Centre de Recherches sur les Très Basses Températures, B.P. 166, F-38042 Grenoble, France

S. Boukraa

LPTHE, Tour 16, 1er étage, 4 Place Jussieu, 75252 Paris Cedex, France
Institut d'Aéronautique, Université de Blida, BP 270, Blida, Algeria

S. Hassani†

CDTN, Boulevard F.Fanon, 16000 Alger, Algeria

J.-M. Maillard‡

LPTHE, Tour 16, 1er étage, 4 Place Jussieu, 75252 Paris Cedex, France

We consider a family of birational transformations of two variables, depending on one parameter, for which simple rational expressions with integer coefficients, for the exact expression of the dynamical zeta function, have been conjectured. together with an equality between the (asymptotic of the) Arnold complexity and the (exponential of the) topological entropy. This identification takes place for the birational mapping seen as a mapping bearing on two complex variables (acting in a complex projective space). We revisit this identification between these two quite “universal complexities” by considering now the mapping as a mapping bearing on two *real* variables. The definitions of the two previous “topological” complexities (Arnold complexity and topological entropy) are modified according to this real-variables point of view. Most of the “universality” is lost. However, the results presented here are, again, in agreement with an identification between the (asymptotic of some) “real Arnold complexity” and the (exponential of some) “real topological entropy”. A detailed analysis of this “real Arnold complexity” as a function of the parameter of this family of birational transformations of two variables is given. One can also slightly modify the definition of the dynamical zeta function, introducing a “real dynamical zeta function” associated with the counting of the real cycles only. Similarly one can also introduce some “real Arnold complexity” generating functions. We show that several of these two “real” generating functions seem to have the same singularities. Furthermore we actually conjecture several simple rational expressions for them, yielding again algebraic values for the (exponential of the) “real topological entropy”. In particular, when the parameter of our family of birational transformations becomes large, we obtain two interesting compatible non trivial rational expressions. These rational results for real mappings cannot be understood by any obvious Markov’s partition, or symbolic dynamics hyperbolic systems interpretation.

PACS numbers: 05.45.+b, 03.20, 46.10, 47.52.+j, 05.50.+q, 02.90.+p

Key words : Arnold complexity, topological entropy, discrete dynamical systems of real variables, birational mappings, Cremona transformations, rational dynamical zeta functions, complex mappings versus real mappings.

I. INTRODUCTION

The purpose of this paper is to sketch a classification of birational transformations based on various notions of “complexity”. In previous papers [1–3] an analysis, based on the examination of the successive (bi)rational expressions corresponding to the iteration of some given birational mappings, has been performed. When one considers the degree

*e-mail : dauriac@crtbt.polycnrs-gre.fr

†e-mail : hassani@ist.cerist.dz

‡e-mail : maillard@lpthe.jussieu.fr

$d(N)$ of the numerators (or denominators) of the corresponding successive rational expressions for the N -th iterate, the growth of this degree is (generically) exponential with N : $d(N) \simeq \lambda^N$. λ has been called the “growth complexity” [4] and it is closely related to the Arnold complexity [6]. A semi-numerical analysis, enabling to compute these growth complexities λ for these birational transformations, has been introduced in [1,2]. It has been seen, on particular sets of birational transformations [5], that these “growth complexities” correspond to a remarkable spectrum of *algebraic* values [4].

These “growth complexities”, summing up the (asymptotic) evolution of the *degree* of the successive iterates, amount to seeing these mappings as mappings of (two) *complex* variables. However, when one considers the phase portrait of these mappings, one also gets some “hint” of the “complexity” of these mappings seen as mappings of (two) *real* variables. In the following, we will consider a one-parameter dependent birational mapping of two variables. On this very example, it will be seen, considering phase portraits corresponding to various values of the parameter, that these “real complexities” vary for the different (positive) values of the parameter. Two universal (or “topological”) measures of the complexities were found to identify [1,2], namely the (asymptotic of the) *Arnold complexity* [6] (or growth complexity) and the (exponential of the) *topological entropy* [1]. The topological entropy is associated with the exponential growth h^N of the number of fixed points (real or complex) of the N -th iterate of the mapping : looking at various phase portraits, corresponding to different values of the parameter (see below), it is tempting to define, in an equivalent way, a “real topological entropy” associated with the exponential growth h_{real}^N of the number of *real fixed points only* of the N -th iterate of the mapping. This notion of “real topological entropy” would actually correspond to the “visual complexity” as seen on the phase portrait of the mapping. Such a concept, corresponding to the evaluation of the real complexity h_{real} of the mapping seen as a mapping bearing on *real* variables, would be less universal: it would have only “some” of the remarkable topological universal properties of the topological entropy. Similarly, it is also tempting to slightly modify the definition of the *Arnold complexity* [6]. The Arnold complexity [6], which corresponds (at least for the mappings of two variables) to the degree growth complexity [2,3], is defined as the number of (real or complex) intersections of a given (generic and complex) line with its N -th iterate : it is straightforward to similarly define a notion of “real Arnold complexity” describing the number of *real* intersections of a given (generic) *real* line with its N -th iterate. This real-analysis concept is, at first sight, also very well-suited to describe the “real complexity” of the mapping as it can be seen in the phase portrait (see Fig.2 below). Recalling the identification, seen on this one-parameter family of birational mappings, between the (asymptotic of the) Arnold complexity and the (exponential of the) topological entropy [1,2], it is natural to wonder if this identification also works for their “real” partners, or if, as the common wisdom could suggest, real analysis is “far less universal”, depending on a lot of details and, thus, requires a “whole bunch” of “complexities” (Lyapounov dimensions, ...) to be described properly.

In order to see the previous identification even more clearly, one can also slightly modify the definition of the dynamical zeta function, introducing a “real dynamical zeta function” associated with the counting of the real cycles only, and, similarly, one can also introduce some “real Arnold complexity” generating functions. We will show that several of these two “real” generating functions have the same singularities. Furthermore we will actually conjecture several *simple rational expressions* for them, yielding, again, *algebraic values* for the (exponential of the) “real topological entropy”. In particular, when the parameter of our family of birational transformations becomes large, we will get an interesting non trivial rational expression. These rational results for real mappings cannot be simply understood by any “obvious” Markov’s partition, or symbolic dynamics hyperbolic interpretation.

II. GROWTH (ARNOLD) COMPLEXITY FOR A BIRATIONAL MAPPING

A *one-parameter* family of birational mappings of *two* (complex) variables has been introduced in previous papers [2,8,9] (see definition (3) in [2]). This mapping actually originates from a lattice statistical mechanics framework that will not be detailed here [8,10,11]. In the following, we will use the extreme simplicity of this mapping of *two* (complex) variables to first compare two quite universal (topological) notions of “complexity” namely the growth complexity λ , which measures the exponential growth of the *degree* of the successive rational expressions encountered in an iteration (a notion which coincides with the (asymptotic of the) *Arnold complexity*¹ [6]), and the (exponential of the) *topological entropy* [12,13]. In section (IV C), we will go a step further and compare, more particularly, the

¹ More precisely the Arnold complexity $C_A(N)$ is proportional (for plane maps) to $d(N)$, the degree of the N -th iteration of the birational mapping which behaves like $d(N) \simeq \lambda^N$. This “degree notion” was also introduced by A. P. Veselov in exact correspondence with the general Arnold definition [6]. Note that the concept of Arnold complexity *is not* restricted to two-dimensional maps.

notion of “*real*” *Arnold complexity* versus the notion of “*real*” *topological entropy*. These two notions will be seen to be suitable to describe the properties of the mapping seen as a mapping of *real* variables.

A. A one-parameter family of birational transformation

Let us consider the following birational transformation (see (3) in [2]) of two (*complex*) variables k_ϵ , depending on one parameter ϵ :

$$k_\epsilon : \quad (y_{n+1}, z_{n+1}) = \left(z_n + 1 - \epsilon, \quad y_n \cdot \frac{z_n - \epsilon}{z_n + 1} \right) \quad (1)$$

In spite of its simplicity, this birational mapping can, however, have quite different behaviors according to the actual values of the parameter ϵ . For example, for $\epsilon = 0$, as well as $\epsilon = -1, 1/2, 1/3$ or 1 , the mapping becomes *integrable*, whereas it is not [9] for all other values of ϵ .

Let us now compare, in the following, two notions of “complexity” (*Arnold complexity versus topological entropy*) according to various values of ϵ .

B. Semi-numerical approach for the growth complexity λ

The (growth) complexity λ , which measures the exponential growth of the degrees of the successive rational expressions one encounters in the iteration of the birational transformation (1), can be obtained by evaluating the degrees of the numerators, or equivalently of the denominators, of the successive (bi)rational expressions obtained in the iteration process. One can actually build a *semi-numerical method* [1,2] to get the value of the complexity growth λ for any value of the parameter ϵ . The idea is to iterate, with the birational transformation (1), a generic *rational* initial point (y_0, z_0) and to follow the magnitude of the successive numerators, or denominators, of the iterates. During the first few steps some accidental simplifications may occur, but, after this transient regime, the integer denominators (for instance) grow like λ^n where n is the number of iterations. Typically, a best fit of the logarithm of the numerator as a linear function of n , between $n = 10$ and $n = 20$, gives the value of λ within an accuracy of 0.1%. Let us remark that an *integrable mapping* yields a *polynomial growth* of the calculations [11] : the value of the complexity λ has to be numerically very close to 1.




FIG. 1. Complexity λ , for k_ϵ , as a function of ϵ .

Fig.1 shows the values of the complexity growth λ as a function of the parameter ϵ . One remarks on Fig.1 that all the values of ϵ (except a zero measure set) give a growth complexity $\lambda \simeq 1.618$. The calculations have been performed

using an infinite-precision² C-library [14]. This semi-numerical analysis [2] clearly indicates, that, beyond the known integrable values [9] of ϵ , namely $-1, 0, 1/3, 1/2, 1$, two sets of values $\{1/4, 1/5, 1/6, \dots, 1/13\}$ and $\{3/5, 2/3, 5/7\}$ are singled out. This suggests that the growth complexity λ takes lower values than the generic one on two infinite sequences of values of ϵ , namely $\epsilon = 1/n$ and $\epsilon = (m-1)/(m+3)$ for n and m integers such that $n \geq 4$ and $m \geq 7$ and m odd.

C. Generating functions for the degree growth of the successive iterates

One can revisit all these results using the stability of the factorization schemes, performing exact formal (maple) calculations. For instance, one can consider, for various values of ϵ , the degrees of the successive rational expressions one encounters when performing the successive iterates, and build various generating functions³ corresponding to these successive degrees. In particular, having singled out a set of values of ϵ , one can revisit these various values, to see how the generic growth complexity $\lambda \simeq 1.618$ gets modified, and deduce the degree generating functions, and the associated complexity λ , in each case. Let us denote by $G_\epsilon(t)$ the degree generating function, corresponding, for some given value of the parameter ϵ , to the degree of (for instance) the numerator of the z component of the successive rational expressions obtained in the iteration process of transformation (1).

At this step, it is worth recalling, again, the notion of Arnold complexity [6] which corresponds to iterate a given (complex) line and count the number A_N of intersections of this N -th iterate with the initial line. It is straightforward to see that these Arnold complexity numbers A_N are closely linked to these successive degrees (see for instance [1,2]). Actually, if one considers the iteration of the $y = (1-\epsilon)/2$ line⁴, the generating function of the A_N 's “almost” identifies (often up to a simple $t/(1+t)$ factor) with the degree generating functions $G_\epsilon(t)$. The “Arnold” generating functions $A_\epsilon(t)$, and the degree generating functions $G_\epsilon(t)$, read (up to order fifteen for the “Arnold” generating functions and order ten, or eleven, for the degree generating functions) :

$$\begin{aligned} A_\epsilon(t) &= \frac{t}{1+t} \cdot G_\epsilon(t) = \frac{t}{1-t-t^2}, & A_{1/m}(t) &= \frac{t}{1+t} \cdot G_{1/m}(t) = \frac{t}{1-t-t^2+t^{m+2}}, \\ A_{(m-1)/(m+3)}(t) &= \frac{t}{1+t} \cdot G_{(m-1)/(m+3)}(t) = \frac{t}{1-t-t^2+t^{m+2}} & \text{for } m = 9, 13, 17, 21, \dots, \\ A_{(m-1)/(m+3)}(t) &= \frac{t \cdot (1-t^{(m+1)/2})}{1+t} \cdot G_{(m-1)/(m+3)}(t) = \frac{t \cdot (1-t^{(m+1)/2})}{1-t-t^2+t^{m+2}} & \text{for } m = 7, 11, 15, \dots, \end{aligned} \quad (2)$$

where the expression for $G_\epsilon(t)$ is valid for ϵ generic and the expressions for $G_{1/m}(t)$ are valid for $m \geq 4$, the $G_{(m-1)/(m+3)}(t)$ for $m \geq 7$ with m odd. One also has for various integrable value of ϵ :

$$\begin{aligned} A_{-1}(t) &= \frac{t}{1+t} \cdot G_{-1}(t) = \frac{t}{1-t^2}, & A_0(t) &= \frac{t}{1+t} \cdot G_0(t) = \frac{t}{(1-t)(1+t)}, \\ A_1(t) &= \frac{t \cdot (1-t)}{1+t} \cdot G_1(t) = \frac{t}{1-t^2}, & A_{1/3}(t) &= \frac{t \cdot (1+t) \cdot (1-t)^2}{1+t^4} \cdot G_{1/3}(t) = \frac{t \cdot (1+t)}{1-t^3}, \\ A_{1/2}(t) &= \frac{t}{1+t} \cdot G_{1/2}(t) = \frac{t \cdot (1-t^9)}{(1-t) \cdot (1-t^2) \cdot (1-t^3) \cdot (1-t^5)}. \end{aligned} \quad (3)$$

These various exact generating functions are in agreement with the previous semi-numerical calculations. In particular, the first expression in (2) yields an algebraic value for λ in agreement with the generic value of the complexity $\lambda \simeq 1.618$ of Fig.1 (and Fig.1 in [2]).

² The multi-precision library gmp (GNU MP) is part of the GNU project. It is a library for *arbitrary* precision arithmetic, operating on signed integers, rational numbers and floating points numbers. It is designed to be as fast as possible, both for small and huge operands. The current version is : 2.0.2. Targeted platforms and Software/Hardware requirements are any Unix machines, DOS and others, with an operating system with reasonable include files and a C compiler.

³ Similar calculations of generating functions have been performed [2] using other representations of the mapping related to 3×3 matrices [15]. These generating functions, denoted $G_\epsilon(x)$ in [2], are deduced from the existence of remarkable stable *factorization schemes* [2,4,5]. These results are in complete agreement with the one given here for the mapping of two variables (1) (see for instance equations (11), (12), (13) and (14) in [2]).

⁴ Which is known to be a singled-out line for this very mapping [2] (see also, below, in sections (IV B), (V)).

III. DYNAMICAL ZETA FUNCTION AND TOPOLOGICAL ENTROPY

It is well known that the periodic orbits (cycles) of a mapping k strongly “encode” dynamical systems [16]. The fixed points of the N -th power of the mapping being the cycles of the mapping itself, their proliferation with N provides a “measure” of chaos [17,18]. To keep track of this number of cycles, one can introduce the fixed points generating function

$$H(t) = \sum_N \# \text{fix}(k^N) \cdot t^N \quad (4)$$

where $\# \text{fix}(k^N)$ is the number of fixed points of k^N , *real or complex*. This quantity only depends on the number of fixed points, and *not* on their particular localization. In this respect, $H(t)$ is a *topologically invariant quantity*. The same information can also be coded in the so-called⁵ *dynamical zeta function* $\zeta(t)$ [13,20] related to the generating function $H(t)$ by $H(t) = t \frac{d}{dt} \log(\zeta(t))$. The dynamical zeta function is defined as follows [16,19,20] :

$$\zeta(t) = \exp \left(\sum_{N=1}^{\infty} \# \text{fix}(k^N) \cdot \frac{t^N}{N} \right) \quad (5)$$

The topological entropy $\log h$ is :

$$\log h = \lim_{N \rightarrow \infty} \frac{\log (\# \text{fix}(k^N))}{N} \quad (6)$$

If the dynamical zeta function is *rational*, h will be the inverse of the pole of smallest modulus of $H(t)$ or $\zeta(t)$. If the dynamical zeta function can be interpreted as the ratio of two characteristic polynomials of two linear operators⁶ A and B , namely $\zeta(t) = \det(1 - t \cdot B) / \det(1 - t \cdot A)$, then the number of fixed points $\# \text{fix}(k^N)$ can be expressed from $\text{Tr}(A^N) - \text{Tr}(B^N)$. In this case, the poles of a rational dynamical zeta function are related to the (inverse of the zeroes of the) characteristic polynomial of the linear operator A only. Since the number of fixed points remains unchanged under *topological conjugacy* (see Smale [24] for this notion), the dynamical zeta function is also a *topologically invariant function*, invariant under a large set of transformations, and does not depend on a specific choice of variables. Such invariances were also noticed for the growth complexity λ . It is thus tempting to make a connection between the *rationality* of the complexity generating function previously given, and a possible *rationality* of the dynamical zeta function. We will also compare the singularities of these two sets of generating functions, namely the growth complexity λ and h , the (exponential of the) topological entropy.

Some results for the dynamical zeta function : Let us now get the expansion of the dynamical zeta function of the mapping k_ϵ , for generic values of ϵ . We can first concentrate on the specific⁷, but arbitrary, value $\epsilon = 21/25$. Of course, there is nothing particular with this specific $\epsilon = 21/25$ value : the same calculations have been performed for many other *generic* values of ϵ yielding the same number of (complex) fixed points and, thus, the same dynamical zeta function. The total number of fixed points of k_ϵ^N , for N running from 1 to 14, yields, up to order fourteen, the following expansion for the generating function $H(t)$ of the number of fixed points :

$$H_\epsilon(t) = H_{21/25}(t) = t + t^2 + 4t^3 + 5t^4 + 11t^5 + 16t^6 + 29t^7 + 45t^8 + 76t^9 + 121t^{10} + 199t^{11} + 320t^{12} + 521t^{13} + 841t^{14} + \dots \quad (7)$$

This expansion coincides with the one of the *rational* function⁸ :

⁵The dynamical zeta function has been introduced by analogy with the Riemann ζ function, by Artin and Mazur [19].

⁶For more details on these Perron-Frobenius, or Ruelle-Araki transfer operators, and other shifts on Markov partition in a symbolic dynamics framework, see for instance [20–23]. In this linear operators framework, the *rationality* of the zeta function, and therefore the algebraicity of the (exponential of the) topological entropy, amounts to having a *finite dimensional representation* of the linear operators A and B .

⁷ Another generic value of ϵ , close to the $1/2$ value where the mapping is integrable [9], namely $\epsilon = 13/25 = 0.52$, has been analyzed in some detail in [2]. For this value $\epsilon = 0.52$, the enumeration of the number of fixed points, n -cycles and the actual status of these fixed points (elliptic, hyperbolic, points ...) are given in [2].

⁸Valid for generic values of ϵ , up to some algebraic values of ϵ corresponding to cycle-fusion mechanism see (27) and (28) below and see [3].

$$H_\epsilon(t) = \frac{t \cdot (1 + t^2)}{(1 - t^2) \cdot (1 - t - t^2)} \quad (8)$$

which corresponds to a very *simple*⁹ *rational* expression for the dynamical zeta function :

$$\zeta_\epsilon(t) = \frac{1 - t^2}{1 - t - t^2} \quad (9)$$

An alternative way of writing the dynamical zeta functions relies on the decomposition of the fixed points into *irreducible cycles* :

$$\zeta_\epsilon(t) = \frac{1}{(1 - t)^{N_1}} \cdot \frac{1}{(1 - t^2)^{N_2}} \cdot \frac{1}{(1 - t^3)^{N_3}} \cdots \frac{1}{(1 - t^r)^{N_r}} \cdots \quad (10)$$

For generic values of ϵ , one gets the following numbers of irreducible cycles : $N_1 = 1, N_2 = 0, N_3 = 1, N_4 = 1, N_5 = 2, N_6 = 2, N_7 = 4, N_8 = 5, N_9 = 8, N_{10} = 11, N_{11} = 18, \dots$ It has been conjectured in [2] that *the simple rational expression (9) is the actual expression of the dynamical zeta function for any generic value of ϵ* (up to some algebraic values of ϵ , see below and in section (IV B)). Similar calculations have been performed for the other values of ϵ that have been singled out in the semi-numerical analysis [3]. For the non generic values of ϵ , $\epsilon = 1/m$ with $m \geq 4$, we have obtained expansions compatible with the following *rational* expression :

$$\zeta_{1/m}(t) = \frac{1 - t^2}{1 - t - t^2 + t^{m+2}} \quad (11)$$

For the other non generic values, namely $\epsilon = (m - 1)/(m + 3)$ with $m \geq 7$ odd, the expansions are not large enough to conjecture a single formula valid for any m . For $m = 7$ (namely $\epsilon = 3/5$) one actually gets a dynamical zeta function given by (11) for $m = 7$ and this might also be the case for $m = 11, 15, \dots$. For $m = 9, 13, \dots$ the expansions are in agreement with a $1 - t - t^2 + t^{m+2}$ singularity. Comparing the various rational expressions in (2) corresponding to generic, and non-generic, values of ϵ , with (9), and (11), respectively, one sees that the singularities (poles) of the dynamical zeta function happen to *coincide* with the poles of the generating functions of the growth complexity λ , for all the values of ϵ . In particular, the growth complexity λ and h , the exponential of the topological entropy, are always *equal*.

Let us just mention, here, that the modification of the number of fixed points, from the “generic” values of ϵ to the particular values $(1/m, (m - 1)/(m + 3))$, corresponds to the *disappearance of some cycles* which become singular points (indetermination of the form 0/0). These mechanisms will be detailed in [25]. Actually the “non-generic” values of ϵ , like $\epsilon = 1/m$, correspond to such a “disappearance of cycles” mechanism which modifies the denominator of the rational generating functions, and, thus, the topological entropy and the growth complexity λ . In contrast, there actually exists for k_ϵ , *other singled-out values* of ϵ , like $\epsilon = 3$ for instance, which correspond to *fusion of cycles* (see section (IV B)) : in the $\epsilon \rightarrow 3$ limit, the order three cycle tends to coincide with the order one cycle, which amounts to multiplying the dynamical zeta function (9) by $1 - t^3$. Such “fusion-cycle” mechanism *does not modify* the denominator of the rational functions, and thus, the topological entropy, or the growth complexity λ , *remain unchanged*.

To sum up : Considering a (very simple) one parameter-dependent birational mapping of *only two (complex) variables*, we have deduced an exact identification between the (asymptotic of the) Arnold complexity, that is the growth complexity λ , and the (exponential of the) topological entropy for *all* the various ϵ cases (generic or not). This identification can be understood heuristically [2]. As a byproduct, one finds that these two complexities correspond, in this very example, to *simple algebraic numbers*.

A canonical degree generating function : This identification result is not completely surprising : the dynamical zeta function is a quite “universal” function, invariant under a large set of *topological conjugaisons* [24], and the concept of Arnold complexity (or the degree growth complexity λ) also has the same “large” set of (topological and projective) invariances [6].

Actually, as far as degree generating functions are concerned, it is natural to introduce, instead of some generating functions of the degrees of the numerator of the z component of the N -th iterate, a more “canonical” degree generating

⁹As far as *symbolic dynamics* is concerned, one can associate, to a dynamical zeta function like (9), a clipped Bernoulli shift with the “pruning rule” to forbid substring $-11-$ (that is 1 must be always followed by 0) in any sequence of 0 and 1. However, constructing the Markovian partitions (if any), yielding this simple pruning rule for the symbolic dynamics, remains to be done.

function $G_\epsilon^{Hom}(t)$ associated with the birational mapping (1) written in a *homogeneous way* (see the bi-polynomial mapping (4) in [2]). Iterating (1), written in a homogeneous way, and, factoring out at each iteration step the gcd's, one gets a new degree generating function $G_\epsilon^{Hom}(t)$ well-suited, at first sight, to describe such large (topological and projective) invariances. A simple calculation shows that this (projectively well-suited) degree generating function reads (for generic ϵ) :

$$G_\epsilon^{Hom}(t) = \frac{1}{(1-t) \cdot (1-t-t^2)} \quad (12)$$

For the $\epsilon = 1/m$ particular values, and for the two integrable values, $\epsilon = 1/2$ and $\epsilon = 1/3$, one gets respectively :

$$G_{1/m}^{Hom}(t) = \frac{1-t^{m+3}}{(1-t) \cdot (1-t-t^2+t^{m+2})} \quad (13)$$

$$G_{1/2}^{Hom}(t) = \frac{1-t^9}{(1-t)^2 \cdot (1-t^3) \cdot (1-t^5)} + \frac{t^2 \cdot (1-t^6)}{(1-t)^2 \cdot (1-t^2) \cdot (1-t^5)} \quad (14)$$

$$G_{1/3}^{Hom}(t) = \frac{1-t^6}{(1-t^3)(1-t^2)(1-t)^2} + \frac{t^4}{(1-t^3)(1-t)^2} \quad (15)$$

Since the expansions for the infinite set of values of the form $(m-1)/(m+3)$ for $m \geq 7$, can only be performed up to order eleven (or twelve), it is difficult to “guess” any expression valid for any m like (13). Recalling the results (2) given in section (II C) for the degree growth generating functions, one may suspect that, among these $(m-1)/(m+3)$ for $m \geq 7$ values, one should make a distinction between $m = 7, 11, 15, \dots$ on one side, and $m = 9, 13, 17, \dots$ on the other side. In fact, up to order eleven, all our calculations for various $(m-1)/(m+3)$ values for $m \geq 7$ ($\epsilon = 3/5, 2/3, 5/7, 3/4, 7/9, 4/5, \dots$) are in agreement with a general equality between $G_{(m-1)/(m+3)}^{Hom}(t)$ and $G_{1/m}^{Hom}(t)$. More details are available in Appendix A.

One gets simpler expressions for the integrable values $\epsilon = 0, 1$ and $\epsilon = -1$:

$$G_0^{Hom}(t) = G_1^{Hom}(t) = \frac{1+t^2}{(1-t)^2}, \quad \text{and :} \quad G_{-1}^{Hom}(t) = \frac{1}{1-t} \quad (16)$$

One remarks that $G_\epsilon^{Hom}(t)$ verifies the simple functional equation $G_\epsilon^{Hom}(t) + G_\epsilon^{Hom}(1/t) = 1$, for $\epsilon = -1, 1/2, 1/3$, and $G_\epsilon^{Hom}(t) = G_\epsilon^{Hom}(1/t)$ for $\epsilon = 0, +1$.

For $\epsilon = 1/2$ we have not written any dynamical zeta function $\zeta_{1/2}(t)$ because, for such an integrable birational mapping, there exist, at (almost) any order N of iteration, an *infinite number* of fixed points of order N (all the points of some elliptic curves [9]) and, therefore, our previous “simple” definition (5) for the dynamical zeta function is not valid anymore.

A possible universal relation. One can imagine many simple relations between the “canonical” degree generating function, $G_\epsilon^{Hom}(t)$, and the dynamical zeta function, $\zeta_\epsilon(t)$. For instance, for generic ϵ , one gets (among many ...) the relation $(1-t) \cdot (1-t^2) \cdot G_\epsilon^{Hom}(t) = \zeta_\epsilon(t)$, however this relation is not anymore valid for $\epsilon = 1/m$. One would like to find a “true universal” relation between $\zeta_\epsilon(t)$ and $G_\epsilon^{Hom}(t)$, that is a relation *independent* of ϵ (generic or non-generic). In order to achieve this goal one may imagine to barter $\zeta_\epsilon(t)$, and $G_\epsilon^{Hom}(t)$, for *projectively well-suited generating functions* taking into account the point at ∞ , namely a dynamical zeta function taking into account the fixed point at ∞ (see (52) in [3]), $\zeta^{(\infty)}(t)$, and $G_\infty^{Hom}(\epsilon, t)$ defined as follows :

$$\zeta_\epsilon^{(\infty)}(t) = \frac{\zeta_\epsilon(t)}{1-t} \quad \text{and :} \quad G_\infty^{Hom}(\epsilon, t) = G_\epsilon^{Hom}(t) + \frac{t}{1-t} \quad (17)$$

One verifies immediately that the relation :

$$G_\infty^{Hom}(\epsilon, t) = (1+t) \cdot \zeta_\epsilon^{(\infty)}(t) \quad \text{or equivalently :} \quad (1+t) \cdot \zeta_\epsilon(t) = (1-t) \cdot G_\epsilon^{Hom}(t) + t \quad (18)$$

is *actually verified* for generic values of ϵ , as well as, for the *non-generic values* of the form $\epsilon = 1/m$, and also some *non-generic values* of the form $\epsilon = (m-1)/(m+3)$ (see Appendix A). A similar relation for the two-parameters family of birational transformations depicted in [1–3] will be detailed elsewhere. Relation (18) should give some hint for a true mathematical proof of the relation between Arnold complexity and topological entropy.

Remark : Recalling the “Arnold” generating functions $A_\epsilon(t)$, (see (2)), which identifies “most of the time” (namely ϵ generic, $\epsilon = 1/m$, $\epsilon = (m+1)/(m+3)$ for $m = 9, 13, 17, \dots$) with the new well-suited generating function

$G_{\infty}^{Hom}(\epsilon, t)$, up to a simple multiplicative factor $t/(1+t)^2$, one can rewrite, for ϵ generic and $\epsilon = 1/m$, relation (18) as :

$$t \cdot \zeta_{\epsilon}(t) = (1 - t^2) \cdot A_{\epsilon}(t) \quad (19)$$

IV. REAL DYNAMICAL ZETA FUNCTION AND REAL TOPOLOGICAL ENTROPY

As far as the growth complexity λ is concerned, the generic values of ϵ (that is the values different from the previous $1/m$, $(m-1)/(m+3)$ singled-out values) are all on the same “complexity footing” (see the previous Fig.1). This is clearly confirmed by the exponential *growth of the computing time* during the iteration process, which seems to be similar for all these values (and clearly *smaller* for the $1/m$, $(m-1)/(m+3)$ particular values). It is however worth noticing that these generic values, which are all on the same λ -footing, clearly yield phase portraits which are quite different and, obviously, correspond to *drastically different* “visual complexities” of the phase portrait of the mapping. This “visual complexity” corresponds to the (exponential) growth of the number of (real) fixed points of the mapping seen as a mapping *bearing on two real variables*. The previous definitions of the dynamical zeta functions $\zeta_{\epsilon}(t)$ and of the generating function $H_{\epsilon}(t)$ counting the number of fixed points, can be straightforwardly modified to describe the counting of *real fixed points* :

$$H_{real}(t) = \sum_N H_N^R \cdot t^N = t \cdot \frac{d}{dt} \log(\zeta^{real}(t)), \quad \text{where :} \quad \zeta^{real}(t) = \sum_N z_N^R \cdot t^N \quad (20)$$

where the number of *real* fixed points H_N^R grow exponentially with the number N of iterates, like $\simeq h_{real}^N$. A quick examination of various phase portrait for various “generic values” of the parameter ϵ seems to indicate quite clearly that this “real topological entropy” $\log(h_{real})$ *varies with* ϵ , in contrast with the “usual” topological entropy $\log(h)$. An obvious inequality is : $h_{real} \leq h$.

A. “Phase portrait gallery”

Let us give here various phase portraits corresponding to different (generic except the first one) values of ϵ . Note the different scale for the frame of these various phase portraits. For most of the phase portraits below around 300 orbits of length 1000, starting from randomly chosen points inside the frame¹⁰, have been generated (only points inside the frame are shown) :

FIG. 2. Phase portrait of k_{ϵ} for $\epsilon = 1/100$ (left) and for $\epsilon = 9/50$ (right).

¹⁰With a special non-random treatment of the regular elliptic parts of the phase portraits.




FIG. 3. Phase portrait of k_ϵ for $\epsilon = 33/100$.




FIG. 6. Phase portrait of k_ϵ for $\epsilon = 34/100$.




FIG. 4. Phase portrait of k_ϵ for $\epsilon = 48/100$.




FIG. 7. Phase portrait of k_ϵ for $\epsilon = 51/100$.




FIG. 5. Phase portrait of k_ϵ for $\epsilon = 9/10$.




FIG. 8. Phase portrait of k_ϵ for $\epsilon = 11/10$.

FIG. 9. Phase portrait of k_ϵ for $\epsilon = 3$ (left) and for $\epsilon = 15$ (right).

On these various phase portraits one sees that one gets, near the integrable value $\epsilon \simeq 1/3$, quite different phase portraits which seem, however, to have roughly the same number of (real) fixed points (see Fig.3 and Fig.6). On these various phase portraits one also sees, quite clearly, that the number of fixed points seems to *decrease* when ϵ crosses the integrable value $\epsilon = 1/2$ and $\epsilon = 1$, going, for instance, from $\epsilon = .48$ to $\epsilon = .51$ (see Fig.4 and Fig.7) or, from $\epsilon = .9$ to $\epsilon = 1.1$ (see Fig.5 and Fig.8). These results will be revisited in a forthcoming section (see section (V)). Of course, exactly on the integrable value $\epsilon = 1$, the phase diagram corresponds to a (simple) foliation of the two-dimensional parameter space in (*rational*) curves (linear pencil of rational curves, see [9]) :

$$\Delta(y, z, 1) = \left(\frac{yz}{y-z-1} \right)^2 = \rho, \quad \text{or equivalently :} \quad \frac{yz}{y-z-1} = \pm \rho^{1/2} \quad (21)$$

where ρ denotes some constant. For the other integrable values one also has either a linear pencil of rational curves, namely $yz = \rho$ for $\epsilon = 0$, as well as :

$$\Delta(y, z, -1) = \frac{1}{(1+z-y)^2} = \rho \quad \text{or equivalently :} \quad (y-z) \cdot (y-z-2) = \frac{1}{\rho} - 1 \quad (22)$$

for $\epsilon = -1$, or a linear pencil of *elliptic* curves for $\epsilon = 1/2$, namely :

$$\Delta(y, z, 1/2) = \frac{(1+z+2yz) \cdot (1-y+2yz) \cdot (1+z-y-2yz)}{(1+z-y)^2} = \rho \quad (23)$$

and also :

$$\Delta(y, z, 1/3) = \frac{(5+3z-3y+9yz) \cdot (1-z-y+3yz) \cdot (1+z-y-3yz) \cdot (1+z+y+3yz)}{(1+z-y)^2} = \rho$$

for $\epsilon = 1/3$. One also remarks that $\epsilon = 3$, which corresponds to the generic $\lambda \simeq 1.618$ growth complexity, also yields a *remarkably regular* phase portrait, “visually” similar to a foliation of the two-dimensional parameter space in curves, suggesting a “real topological complexity” h_{real} very close, or even equal to 1. This fact will also be revisited in the next section (see section (V)). In order to describe, less qualitatively, the “real topological complexity” h_{real} as a function of the parameter ϵ , we have calculated in section (IV C), the first (ten, eleven or even twelve) coefficients of the expansions of $H_\epsilon^{real}(t)$, and of the “real dynamical function” $\zeta_\epsilon^{real}(t)$, for various values of ϵ .

B. Number of real fixed points as a function of ϵ .

Let us now try to understand why (and how) h_{real} varies as a function of ϵ , and why some other values of ϵ , like $\epsilon = 3$, different from the previous $1/m$ and $(m-1)/(m+3)$ non-generic values, seem to play a special role. The method to get the fixed points of the N -th iterate of k_ϵ has been detailed in previous papers [1,2]. Let us just mention here that, due to the symmetries of this mapping, there exist two singled out lines, namely $y = (1-\epsilon)/2$

and $y = -z$, playing a key role in classifying all these fixed points [7]. The fixed points of k_ϵ^N belong to N -cycles (if $\mathcal{M}$ is a fixed point of k_ϵ^N then $k_\epsilon^p(\mathcal{M})$ is also a fixed point of k_ϵ^N , for $p = 1, \dots, N-1$). It can be seen that *there always exist* a fixed point in these N -cycles which belongs, either to line $y = (1-\epsilon)/2$, or to the $y = -z$ line. We call the fixed points, corresponding to line $y = -z$, the “P-type” points and the ones, corresponding to line $y = (1-\epsilon)/2$, the “Q-type” points [25]. For $N \geq 9$, other N -cycles with no points lying on these two lines do occur: other remarkable sets occur like $y + \bar{z} = 0$ (see [2,3]). We call the fixed points, corresponding this “remaining” set of points, the points of the “R-type” [25].

One can use these localization properties to get, very quickly, a subset of all the fixed points, namely, for instance, the “Q-type” fixed points (one representant in the N -cycle belongs to line $y = (1-\epsilon)/2$). These calculations can be performed quite efficiently since one can eliminate the y variable ($y = (1-\epsilon)/2$) and, thus, reduce the calculations to looking for the roots (real or not) of an ϵ -dependent polynomial in this remaining z variable. One gets, for the first values of N , the following polynomial expressions relating z and ϵ :

$$\begin{aligned} Q_1(z, \epsilon) &= 2 \cdot z - (\epsilon - 1) = 0, & Q_3(z, \epsilon) &= z - (\epsilon - 2) = 0, \\ Q_5(z, \epsilon) &= (3\epsilon - 1) \cdot z^2 - 2 \cdot (\epsilon - 3) \cdot (2\epsilon - 1) \cdot z + (\epsilon^3 - 5\epsilon^2 + 10\epsilon - 4) = 0, \dots \end{aligned} \quad (24)$$

It is easy to see that the number of real roots z , of one of these $Q_N(z, \epsilon) = 0$ conditions, varies with ϵ by intervals. The changes of this number of real roots take place at *algebraic* values of ϵ (resultant of $Q_N(z, \epsilon)$ in z). The details of the calculations, and a description of these polynomials, will be given elsewhere [25]. The number of the fixed points of the “Q-type” (see [7]) is, thus, a function of ϵ constant by interval, the limits of the intervals corresponding to some algebraic values (resultants deduced from the Q_N ’s by eliminating z). For illustration, let us just plot here the real roots z , as a function of ϵ , for Q_{10} :




FIG. 10. The real roots z of Q_{10} , as a function of ϵ , in the interval $[-1.5, 2.5]$. We have included the plot of the total number of fixed points of k_ϵ^{10} (“P-type”, “Q-type” and “R-type”) as a function of ϵ in an interval around $[0, 2]$.

Let us give, for order 10, a few examples of these algebraic “threshold” real values of ϵ corresponding to the real roots of such “Q-type” polynomials (24) :

$$5\epsilon^2 - 10\epsilon + 1 = 0, \quad 5\epsilon^4 - 96\epsilon^3 + 114\epsilon^2 - 40\epsilon + 1 = 0 \quad (25)$$

The roots of the first polynomial are of the form (27). The “threshold” values of ϵ are thus given by two real roots $\epsilon \simeq .1055$ and 1.8944 . The second polynomial only gives two real roots $\epsilon \simeq .02703$ and $\simeq 17.9549$.

Similar calculations can be performed for the fixed points of the “P-type” (see [7]) corresponding to the line $y = -z$ (see [25]). One can also get the real roots z of P_{10} as a function of ϵ . The algebraic values of ϵ , occurring in this case,

are, again, the two roots of the first polynomial in (25) together with the only two real roots $\epsilon \simeq .1561$ and $.5013$ of polynomial :

$$\epsilon^8 - 26\epsilon^7 + 343\epsilon^6 - 2052\epsilon^5 + 6367\epsilon^4 - 7178\epsilon^3 + 3625\epsilon^2 - 824\epsilon + 64 = 0 \quad (26)$$

as well as two real roots $\epsilon \simeq .008999$ and $.1316$ of a polynomial of degree 24 in ϵ that will not be written here. The last set of points of the “*R*-type” (see [7]), which corresponds to fixed points that are neither of the “*P*-type”, nor of the “*Q*-type”, give the following real roots $\epsilon \simeq .2338$, $.51434$, and 33.2517 , corresponding to polynomial $\epsilon^3 - 34\epsilon^2 + 25\epsilon - 4 = 0$. On all these algebraic values of ϵ , one can see a variation of the total number of fixed points (“*P*-type”, “*Q*-type” and “*R*-type”) of order 10 (see insertion in Fig.10). These values are in fact particular examples of families of algebraic ϵ values. The simplest family of singled out *algebraic* values of ϵ , corresponds to the fusion on an N -cycle with the 1-cycle, and reads :

$$\epsilon = \frac{1 - \cos(2\pi M/N)}{1 + \cos(2\pi M/N)}, \quad \text{or equivalently :} \quad \cos(2\pi M/N) = \frac{1 - \epsilon}{1 + \epsilon} \quad (27)$$

for any integer N (with $1 < M < N/2$, M not a divisor of N). Other cycle-fusion mechanisms take place yielding new families of algebraic values for ϵ . For instance the coalescence of the $(3 \times N)$ -cycles in the 3-cycle, and the coalescence of the $(4 \times N)$ -cycles in the 4-cycle, yield respectively (with some constraints on the integer M that will not be detailed here) :

$$\cos(2\pi M/N) = 1 - \frac{3}{4} \frac{\epsilon(\epsilon - 3)^2}{(1 - \epsilon)(1 + \epsilon)}, \quad \cos(2\pi M/N) = 1 - 32 \frac{\epsilon(1 - \epsilon)^2}{(1 + \epsilon)^2(1 - 2\epsilon)} \quad (28)$$

Status of the fixed points : The fixed point of k_ϵ , which is elliptic for $\epsilon > 0$, becomes *hyperbolic* for $\epsilon < 0$. For three iterations ($N = 3$) one finds that, moving through the $\epsilon = 1/3$ value, also changes the status of these fixed points from elliptic to hyperbolic. In fact, the algebraic values, like the ones depicted in (27) or (28), *also occur* in such changes of status from elliptic to hyperbolic (see [25]). Therefore, the number of elliptic fixed points, or hyperbolic fixed points, is not as “universal” as the total number of (complex) fixed points, however it has “some universality” : for a given value of N , the number of elliptic (resp. hyperbolic) fixed points depends on ϵ also by *intervals* (staircase function). This has, again, to be compared with the dependence of the growth complexity λ , seen as a function of ϵ , depicted in Fig.1. The fact that the number of hyperbolic versus elliptic fixed points, as well as the number of *real versus non-real* fixed points, is modified when ϵ goes through the *same set* of values, like (27) or (28), seems to indicate that a modification of the number of *real fixed points is not independent of the actual status of these points* (hyperbolic versus elliptic). This phenomenon, in fact, corresponds to a quite involved, and interesting, *structure* [27] that will be sketched in [25].

C. Some expansions for the “real dynamical zeta function” and H^{real} .

Let us recall some results corresponding to $\epsilon = .52$, in particular the product decomposition (10) of the dynamical zeta function [2,3]. In [2,3], the number of irreducible cycles N_i (see (10)), as well as the number of irreducible cycles corresponding to hyperbolic points, elliptic points, real points is detailed (see Table I in [3]). These results (and further calculations) enable to write, for $\epsilon = .52$, the “real dynamical function” $\zeta_\epsilon^{real}(t)$ as the following product :

$$\zeta_{52/100}^{real}(t) = \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^5)^2(1-t^7)^2(1-t^8)(1-t^9)^4(1-t^{10})^2} \times \frac{1}{(1-t^{11})^6(1-t^{12})^{12}(1-t^{13})^{16}} \dots \quad (29)$$

yielding the following expansion for $\zeta_\epsilon^{real}(t)$ and $H_\epsilon^{real}(t)$:

$$\begin{aligned} \zeta_{52/100}^{real}(t) &= 1 + t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 20t^9 + 28t^{10} + 40t^{11} + 65t^{12} + 97t^{13} + \dots \\ H_{52/100}^{real}(t) &= t + t^2 + 4t^3 + 5t^4 + 11t^5 + 4t^6 + 15t^7 + 13t^8 + 40t^9 + 31t^{10} + 67t^{11} + 152t^{12} + 209t^{13} + \dots \end{aligned} \quad (30)$$

The number of *real* n -th cycles of the *P*-type, *Q*-type and *R*-type, denoted P_n , Q_n , and R_n respectively, are given in Table IV in Appendix A. For the R_n ’s one cannot reduce, in contrast with the *P*-type or *Q*-type analysis, the

calculations to a only one variable : one is obliged to perform a first resultant calculation where one eliminates one of the two variables and another resultant calculation where one eliminates the other one, and check back, in the cartesian product of these possible solutions, the solutions which are actually fixed points. In order to get integer values that can be trusted, one needs to perform these (maple) calculations with more than 2000 digits for order twelve, but then one faces severe memory limitations in the formal calculations. We have been able to find integer values for the R_n 's for orders larger than twelve, however it is clear that these integers are just lower bounds of the true integers (not enough precision does not enable to discriminate between fixed points that are very close). Therefore we prefer not to give these integers here, and put a “star” in Table IV, as well as in the forthcoming tables given in Appendix A, when we encounter these computer limitations.

The total number, T_n , of *real* cycles of the P -type, R -type and Q -type actually corresponds to the exponents in the product decomposition (29) for the “real dynamical zeta function”. Unfortunately, these series are not large enough to “guess” any possible (and simple, like (9)) rational expression for $\zeta^{real}(t)$, if any ... Series (30), however, give a first “rough estimate” for the “real topological complexity” h_{real} : $h_{real} \simeq (97)^{1/13} \simeq 1.4217$ or may be $h_{real} \simeq 209^{1/13} \simeq 1.508$, *clearly smaller* than the exact algebraic value for h corresponding to (9) : $h \simeq 1.61803$.

Let us consider other values of ϵ .

• For $\epsilon < 0$, one finds out that *all the fixed points seem to be real* and, thus, one can conjecture for $\epsilon < 0$ (but $\epsilon \neq -1$) :

$$\zeta_{\epsilon < 0}^{real}(t) = \frac{1 - t^2}{1 - t - t^2}, \quad \text{and :} \quad h_{real} = h \simeq 1.61803 \quad (31)$$

The number of cycles of the P -type, Q -type and R -type is given order by order in Table I in Appendix A. These successive integer values for the total number of irreducible real cycles, T_n , yield :

$$\begin{aligned} \zeta_{\epsilon < 0}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^5)^2(1-t^6)^2(1-t^7)^4(1-t^8)^5(1-t^9)^8(1-t^{10})^{11}} \\ &\times \frac{1}{(1-t^{11})^{18}(1-t^{12})^{25}(1-t^{13})^{40}(1-t^{14})^{58}(1-t^{15})^{90}(1-t^{16})^{135}(1-t^{17})^{210}(1-t^{18})^{316} \dots} \end{aligned} \quad (32)$$

• For $\epsilon = 3$, one has, at every order of iteration, up to order twelve, a *only one real* fixed point (the fixed point of order one but, of course, many complex fixed points) yielding :

$$H_3^{real}(t) = \frac{t}{1-t} \quad \text{and :} \quad \zeta_3^{real}(t) = \frac{1}{1-t} \quad (33)$$

and, for ϵ very close to 3, the expressions of $H^{real}(t)$ and $\zeta^{real}(t)$ cannot be distinguished (at the orders where we have been able to perform these fixed points calculations) from :

$$H_{\epsilon \simeq 3}^{real}(t) \simeq \frac{t \cdot (1 + t + 4t^2)}{1 - t^3} \quad \text{and :} \quad \zeta_{\epsilon \simeq 3}^{real}(t) \simeq \frac{1}{(1-t) \cdot (1-t^3)} \quad (34)$$

which just correspond to add an additional 3-cycle.

Expressions (33) are in agreement with the phase portrait of Fig.9 for $\epsilon = 3$. This indicates that, seen as a mapping of two *real variables*, the mapping “looks like” an integrable mapping: the “real topological complexity” h_{real} seems to be exactly equal to 1 for $\epsilon = 3$ (and $h_{real} \simeq 1$ for $\epsilon \simeq 3$). The “real topological entropy” $\log(h_{real})$ seems to be exactly zero for $\epsilon = 3$ and is, thus, drastically different from the generic “usual” topological entropy $\log(1.61803\dots)$. The possible foliation of the two-dimensional space in (transcendental) curves is discussed¹¹ elsewhere [26]. For other values of ϵ the series are not large enough to “guess” a rational expression (if any ...) for the “real dynamical zeta function” $\zeta^{real}(t)$.

• Miscellaneous examples are given in Appendix B. In particular the number of cycles of the P -type, Q -type and R -type is given in Table VI for $\epsilon = 11/10$, yielding the following expansion for the real dynamical zeta function :

$$\zeta_{11/10}^{real}(t) = 1 + t + t^2 + 2t^3 + 2t^4 + 2t^5 + 3t^6 + 5t^7 + 5t^8 + 6t^9 + 10t^{10} + 12t^{11} + 13t^{12} + \dots \quad (35)$$

¹¹ In particular it is shown that, at least, three of the (real) curves of the phase portrait correspond to *divergent series* satisfying an exact functional equation [26].

clearly yielding a value for h_{real} close to one (may be $h_{real} \simeq (13)^{1/12} \simeq 1.238$) significantly smaller than $h \simeq 1.618$. This result has to be compared with the equivalent one for $\epsilon = 9/10$ or for $\epsilon = 21/25$:

$$\begin{aligned}\zeta_{21/25}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^7)^2(1-t^8)(1-t^{10})^2(1-t^{11})^4(1-t^{12})^2} \cdots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 3t^5 + 4t^6 + 7t^7 + 9t^8 + 10t^9 + 15t^{10} + 23t^{11} + 28t^{12} + \dots\end{aligned}\quad (36)$$

yielding a larger value for h_{real} : $h_{real} \simeq (28)^{1/12} \simeq 1.32$. This expansion is actually compatible with the following simple rational expression and for its logarithmic derivative $H_{21/25}^{real}$:

$$\zeta_{21/25}^{real}(t) = \frac{1+t^2}{1-t+t^2-2t^3} \quad \text{and :} \quad H_{21/25}^{real} = \frac{t(5t^2+2t^4+1)}{(1+t^2) \cdot (1-t+t^2-2t^3)} \quad (37)$$

Note that all the coefficients of the expansion of the rational expression (37) and of its logarithmic derivative $H_{21/25}^{real}$ are positive (in contrast with Pade approximation (60) given in Appendix C for $\epsilon = 2/3$ which is ruled out because coefficient t^{54} of its expansion is negative). If this simple rational expression is actually the exact expression for the real dynamical zeta function $\zeta_{21/25}^{real}(t)$ this would yield the following algebraic value for h_{real} : $h_{real}(21/25) \simeq 1.353209964$.

For $\epsilon = 9/10$, one gets the same product decomposition, at least up to order ten. The number of n -th cycles of the P -type, Q -type and R -type for $\epsilon = 9/10$ are given in Appendix B. One thus sees that h_{real} decreases when ϵ crosses the $\epsilon = 1$ value.

- For $\epsilon = 1/4$ we have obtained (see Appendix B) :

$$\begin{aligned}\zeta_{1/4}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^5)^2(1-t^7)(1-t^8)(1-t^9)^3(1-t^{10})^2(1-t^{11})^4(1-t^{12})^4(1-t^{13})^8} \cdots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 8t^7 + 12t^8 + 18t^9 + 25t^{10} + 34t^{11} + 48t^{12} + 70t^{13} + \dots\end{aligned}\quad (38)$$

The “non-generic” values $\epsilon = 1/m$ and $(m-1)/(m+3)$ require a special and careful analysis¹². However, similarly to what was seen for $\zeta(t)$, one clearly verifies on all these “real dynamical zeta function” $\zeta^{real}(t)$ that the coefficients in these expansions are continuous in ϵ near these points except on these very values of ϵ where one gets smaller integers and thus smaller values for h_{real} (the limit on the left and on the right of h_{real} are equal and larger than h_{real} on these very “non-generic” values).

The numbers of irreducible *real* n -cycles of the P -type, Q -type and R -type are given in Appendix B for miscellaneous values of ϵ : $\epsilon = 11/100$, $\epsilon = 5$, $\epsilon = 10$, $\epsilon = 50$. We also give, in Appendix C, for various values of ϵ ($\epsilon = 9/50$, $\epsilon = 31/125$, $\epsilon = 12/25$, $\epsilon = 17/25$, $\epsilon = 66/125$, $\epsilon = 3/4$, $\epsilon = 3/2$), the product decomposition and expansions for $\zeta_{\epsilon}^{real}(t)$ up to order 11.

Similar calculations of the expansions of $H_{\epsilon}^{real}(t)$ and $\zeta_{\epsilon}^{real}(t)$, for many other values of the parameter ϵ , have been performed and will be detailed elsewhere. All these results confirm that h_{real} *varies with* ϵ when ϵ is positive, while h_{real} is constant (except $\epsilon = -1$) when $\epsilon < 0$. When ϵ is positive, the estimates of h_{real} are in agreement with the “visual complexity” as seen on the phase portraits (see the previous section). In particular one finds that h_{real} roughly decreases as a function of ϵ in the intervals $[0^+, \simeq 1/10]$ and $[\simeq 1/3, 1^-]$, and increases in the interval $[\simeq 1/10, \simeq 1/3]$, (with a sharp decrease near $\epsilon \simeq 1/2$ and $\epsilon \simeq 1$), that h_{real} is close or very close to one when ϵ belongs to an interval $[1^+, \simeq 16]$, that h_{real} grows monotonically with ϵ for $\epsilon > 16$ to reach some asymptotic value in the $\epsilon \rightarrow \infty$ limit. It will be seen, in the next subsection, that the “real topological complexity” h_{real} , in the $\epsilon \rightarrow \infty$ limit, tends to a value $h_{real} \simeq 1.429$ clearly different, again, from the generic “topological complexity” $h \simeq 1.618$.

D. Seeking for rationality for the “real dynamical zeta function”.

Recalling the large number of rational expressions, obtained for the dynamical zeta functions [2,3] and the degree generating functions [2,4,5], one may have a rationality “prejudice” for these “real dynamical zeta functions” $\zeta_{\epsilon}^{real}(t)$,

¹²Some fixed points near these “non-generic” values ($\epsilon \simeq 1/m$), disappear on these very values *stricto sensu* : they become singular. One has to verify carefully that all the points, obtained in such calculations, are fixed points and not singular points.

calculated for a *given* value of ϵ . However, the occurrence of any symbolic dynamic, and associated Markov partition, is *far from being natural in this real analysis framework* [28–30]. If one bets on the rationality of the real dynamical zeta function $\zeta_\epsilon^{real}(t)$ (see (20)), it must, however, be clear that $\zeta_\epsilon^{real}(t)$ depends on ϵ in a very complicated way (piecewise continuous functions, devil’s staircase ? ...). If, for some given value of the parameter ϵ , the partial dynamical zeta function $\zeta_\epsilon^{real}(t)$ actually corresponds to a rational expression, one should, in fact, have an *infinite set* of such rational expressions associated with the infinite number of steps (intervals¹³ in ϵ) in the “devil’s staircase”. The actual location of these “steps”, that is, the limits of these intervals in ϵ , corresponds to an infinite number of values of ϵ like (27) or (28) (and others [25]). For a given ϵ , the calculations of the first terms of the expansion of the “real dynamical zeta function” $\zeta_\epsilon^{real}(t)$ do not rule out, at the order for which we have been able to perform these calculations (ten, eleven), *rational expressions* (see for instance [26]).

The number of *real* n -th cycles of the P -type, Q -type and R -type, for $\epsilon = 50$, is given in Table IX in Appendix B. One remarks that, at order eleven, the number of irreducible *real* cycles, and therefore the expansion of the “real dynamical zeta function” are the same for $\epsilon = 50, 100, 1000, \dots$. For $\epsilon = 50, \epsilon = 100$ one has a product expansion for the dynamical zeta function identical, up to order eleven, to the product expansion corresponding to the ϵ large limit (see (39) below). These expansions are, however, different at order twelve, see Appendix B.

• **ϵ large.** For ϵ large enough one gets the following cycle product decomposition :

$$\zeta_{\epsilon=\infty}^{real}(t) = \frac{1}{(1-t)(1-t^3)(1-t^5)^2(1-t^7)^2(1-t^8)^2(1-t^9)^2(1-t^{10})^3(1-t^{11})^4(1-t^{12})^6} \dots \quad (39)$$

corresponding to the number of *real* P -type, Q -type and R -type n cycles for $\epsilon = 20000$ given in Table XI in Appendix B. Note that one gets the same table (up to order 16) for $\epsilon = 1000, 100000, 1000000$.

One finds out easily that these results, for the “real dynamical zeta function” $\zeta^{real}(t)$, are (up to order twelve) actually in *perfect agreement* with (the expansion of) the rational expression :

$$\zeta_{\epsilon=\infty}^{real}(t) = \frac{1+t}{1-t^2-t^3-t^5} = \frac{1-t^2}{(1-t-t^2)+t^4 \cdot (1-t+t^2)} \quad (40)$$

yielding an algebraic value for h_{real} : $h_{real} \simeq 1.4291$. If one “believes” in some symbolic dynamic coding interpretation, or in the existence of a linear transfer operator¹⁴, matrix A , such that the denominator of (40), $1-t^2-t^3-t^5$, can be written as $\det(Id - t \cdot A)$, one finds that a possible choice for this transition matrix is :

$$A = \begin{bmatrix} 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix} \quad (41)$$

In contrast with a Markov’s transition matrix the previous matrix is not such that the sums of the entries in each row, or column, are equal.

V. “REAL ARNOLD COMPLEXITY”

Let us recall, again, the identification between h , the (exponential of the) topological entropy, and λ , the (asymptotic of the) Arnold complexity [2]. Similarly to the topological entropy, the Arnold complexity can be “adapted” to define a “*real* Arnold complexity”. The Arnold complexity counts the number of intersections between a fixed (complex projective) line and its N -th iterate [6] : let us now count, here, the number of *real* points which are the intersections between a *real fixed line* and its N -th iterate. With this restriction to *real* points we have lost “most of


¹³In contrast with the situation for the “customary” dynamical zeta function which is equal to one generic universal expression (like (9)), up to a (zero measure) set of values of ϵ (see Fig.1).

¹⁴For more details on linear transfer operators in a symbolic dynamics framework, see for instance [20–23].

the universality properties” of the “usual” (complex) Arnold complexity. For various values of ϵ , we have calculated the number of intersections of various (real) lines with their N -th iterates. In contrast with the “usual” Arnold complexity [6], which does not depend on the (complex) line one iterates (topological invariance [24]), it is clear that the number of real intersections depends on the chosen line, but one can expect that the *asymptotic behavior* of these numbers for N large enough, will not depend too much of the actual choice of the (real) line one iterates. Actually, we have discovered on this very example, that this seems to be the case (except for some non generic lines). Furthermore, the real line $y = (1 - \epsilon)/2$, which is known to play a particular role for mapping (1) (see section (IV B)), is very well-suited to perform these numbers of intersections calculations : for this particular line the successive numbers of intersections are extremely regular, thus enabling to better estimate this asymptotic behavior λ_{real}^N of the “real Arnold complexity”, but of course similar calculations can be performed with an arbitrary (generic) line. λ_{real} can be seen as the equivalent, for *real mappings*, of the growth complexity λ (see section (II B) and [4]). Let us try to get λ_{real} for various values of ϵ .

Similarly to the semi-numerical method detailed in section (II B), we have developed a C-program using again the multiprecision library gmp [14], counting the number of (real) intersections of the $y = (1 - \epsilon)/2$ real line with its N -th iterate. This program does not calculate the precise location of the intersection points : it is based on the *Sturm’s theorem*¹⁵. All these calculations have been cross-checked by a (maple) program calculating these numbers of intersections using the sturm procedure in maple¹⁶. The results of these calculations are given in Fig.11.

Let us denote by $\mathcal{A}_N$ the number of (real) intersections for the N -th iterate. In order to estimate “real growth complexity” λ_{real} we have plotted $\mathcal{A}_N^{1/N}$, for various values of the number of iterations ($N = 13, 14, 15$), as a function of ϵ , in the range $[0, 1]$ where λ_{real} has a quite “rich” behavior.



¹⁵Assuming that a polynomial $P(x)$ has no multiple roots, one can build a finite series of polynomials corresponding to the successive Euclidean division of $P(x)$ by its first derivative $P'(x)$. See for instance [31] for more details on the Sturm sequences and Sturm’s theorem.

¹⁶The sturm procedure one can find in maple gives the number of real roots of a polynomial in any interval $[a, b]$, even the interval $]-\infty, +\infty[$. The procedure sturm uses *Sturm’s theorem* to return the number of real roots of polynomial P in the interval $[a, b]$. The first argument of this sturm procedure is a *Sturm sequence* for P , which can be obtained with another procedure, the procedure sturmseq which returns the Sturm sequence as a list of polynomials and replaces multiple roots by single roots.

FIG. 11. A representation of $\lambda_{real}(\epsilon)$ by $\mathcal{A}_N^{1/N}$, as a function of ϵ , in the $[0, 1]$ interval, for $N = 13$ (full line) 14 (dashed line), and 15 (dotted line). The singled out $\epsilon = 1/m$ values, for $m = 2, 3, 4, 5, \dots$, and $\epsilon = (m-1)/(m+3)$ for $m = 7, 9, \dots$ clearly play a special role for these various “staircase functions”, in the large N limit.

This behavior should be compared with the “universal” behavior of Fig. 1. On Fig.11, one remarks that the singled out values $\epsilon = 1/m$, as well as $\epsilon = (m-1)/(m+3)$ for $m = 7, 9, \dots$, seem, again, to play a special role in the large N limit. Of course, recalling the results of section (IV B), it is clear that $\mathcal{A}_N^{1/N}$ is a staircase function of ϵ , for N finite, the limits of each interval corresponding to *algebraic values* (like (27)) sketched in sections (IV C), (IV B). These algebraic values form an *infinite set* of values which accumulate everywhere in the $[0, 1]$ interval. What is the limit of these functions $\mathcal{A}_N^{1/N}(\epsilon)$ when N gets large : a devil’s staircase or a (piecewise) continuous function? The “shape” of $\mathcal{A}_N^{1/N}$, as a function of ϵ , is “monotonic enough” (see Fig.12 below) in different intervals, namely $\epsilon < 0$, and in the intervals of ϵ roughly given by : $[0^+, \simeq 1/10]$, $[\simeq 1/10, \simeq 1/3]$, $[\simeq 1/3, 1^-]$, $[1^+, \simeq 16.8]$ and $[\simeq 16.8, \infty]$, that one may expect that the infinite accumulation of these algebraic values (like (27)) could yield a perfectly continuous function $\lambda_{real}(\epsilon)$ (except on the non-generic values $\epsilon = 1/m$ and $\epsilon = (m-1)/(m+3)$) and not a devil’s staircase-like function. This question remains open at the present moment. When ϵ varies from $-\infty$ to ∞ the behavior of the “real growth complexity” λ_{real} , as a function of ϵ , is not as “rich” as in the interval $[0^+, 1^-]$ depicted in Fig.11. One finds that λ_{real} is close, or extremely close to 1, in a quite large interval $[1^+, \simeq 16.8]$ and that it increases monotonically with ϵ in the $[\simeq 16.8, \infty]$ interval to reach some asymptotic value in the $\epsilon \rightarrow \infty$ limit. In fact, a logarithmic scale in ϵ is better suited to describe λ_{real} as a function of ϵ . Fig.12 represents λ_{real} , more precisely $\mathcal{A}_{13}^{1/13}$, as a function of $\log(2 + \epsilon)$. For $\epsilon = -1, 0, 1/3, 1/2, 1$ we know that λ_{real} will be exactly equal to 1 (integrable cases [9]). On these points (represented by squares on Fig.12), as well as on the $\epsilon = 1/m$ and $\epsilon = (m-1)/(m+3)$ non-generic points, λ_{real} is *not continuous* as a function of ϵ . We have not represented these other non generic points. They have to be calculated separately.




FIG. 12. A representation of λ_{real} by $\mathcal{A}_{13}^{1/13}$, as a function of ϵ , in a $\log(2 + \epsilon)$ logarithmic scale. The integrable points $\epsilon = -1, 0, 1/3, 1/2, 1$ are represented by a square.

A first estimate of λ_{real} , for ϵ large, is $\lambda_{real} \simeq (214)^{1/15} \simeq 1.43008$. We are now ready to compare the “real topological entropy” and the “real Arnold complexity” for different values of ϵ , and see if the identification, between h , (the exponential of) the topological entropy, and λ , characterizing the (asymptotic behavior of the) usual Arnold complexity, also holds for their “real partners” namely h_{real} and λ_{real} . Actually, one finds that this identification (which is obviously true for $\epsilon < 0$) also holds for $\epsilon = 3$ and give numerical results for various values of ϵ (for which we have estimated the “real” topological entropy $\log(h_{real})$ (see section (IV C))), *quite compatible with this*

identification. In particular, for ϵ large, we do see that these two “real complexities” *give extremely close results*, namely $h_{real} \simeq 1.4291$ versus $\lambda_{real} \simeq 1.43$.

VI. “REAL ARNOLD COMPLEXITY” GENERATING FUNCTIONS : SEEKING FOR RATIONALITY.

Similarly to the introduction of the “real dynamical zeta functions” $\zeta^{real}(t)$, one can introduce the generating function of the previous “real Arnold complexities” $\mathcal{A}_N$:

$$\mathcal{A}_\epsilon(t) = \sum_N \mathcal{A}_N \cdot t^N \quad (42)$$

Recalling the large number of rational expressions, obtained for the dynamical zeta functions [2,3] and the degree generating functions [2,4,5], one may have, again, a rationality “prejudice” for these “real Arnold complexity generating functions”. Let us try to see if the expansions of these generating functions $\mathcal{A}_\epsilon(t)$ could coincide, for some given values of ϵ , with the expansion of some (hopefully simple) rational expressions.

In order to compare more carefully h_{real} and λ_{real} , and find some possible non trivial rational expressions for $\mathcal{A}_\epsilon(t)$, let us give, in the following, miscellaneous expansions of $\mathcal{A}_\epsilon(t)$ for various values of ϵ .

A. Expansions for the “real Arnold complexity” generating functions

In fact, we have not only calculated the real Arnold complexities $\mathcal{A}_{13}$, $\mathcal{A}_{14}$ and $\mathcal{A}_{15}$ required to plot Fig.11 and Fig.12, but actually obtained all the coefficients up to order 13 for 2000 values of ϵ , and up to order 15 for 200 values of ϵ . We thus have the expansion of $\mathcal{A}_\epsilon(t)$ up to order 13 (resp. 15) for several thousands of values of ϵ .

Let us first give the expansion of $\mathcal{A}_\epsilon(t)$ for $\epsilon = .52$ for which the expansion of the real dynamical zeta function has been given previously (see (29) and (30)). One gets the following expansion :

$$\mathcal{A}_{.52/100}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 11t^7 + 11t^8 + 16t^9 + 29t^{10} + 33t^{11} + 46t^{12} + 73t^{13} + \dots \quad (43)$$

This series yields a first rough approximation of λ_{real} corresponding to $\lambda_{real} \simeq (73)^{1/13} \simeq 1.391$, clearly smaller compared to the generic complexity $\lambda \simeq 1.61803$, and in good enough agreement with the estimation of h_{real} one can deduce from (30), namely $h_{real} \simeq (93)^{1/13} \simeq 1.417$. Of course these two series are too short to see if an identity like $h_{real} = \lambda_{real}$ really holds.

Considering h_{real} as a function of ϵ , it is clear that the general shape of this “curve” looks extremely similar to the curve corresponding to λ_{real} as a function of ϵ (see Fig.11 and Fig.12) : it is also constant for ϵ negative, gets to close to 1 around $\epsilon \simeq 3$, grows monotonically for $\epsilon > 10$ and tends to a non-trivial asymptotic value $h_{real} \simeq 1.4291$.

Therefore, in order to get some hint on the relevance of a possible $h_{real} = \lambda_{real}$ identity, it is necessary to see if this relation holds for various values of ϵ for which h_{real} , and λ_{real} , can be calculated exactly or for which very good approximations of can be obtained, namely $\epsilon < 0$, all the integrable values, or $\epsilon = 3$ and its neighborhood ...

For *any negative value* of ϵ (except $\epsilon = -1$ see below (49)) the expansion of the “real Arnold complexity” generating function $\mathcal{A}_\epsilon(t)$ is equal, up to order 15, to :

$$\mathcal{A}_{\epsilon < 0}(t) = \frac{t}{1 - t - t^2} \quad (44)$$

in agreement with (31).

For $\epsilon = 3$ the generating function for the “real Arnold complexity” generating function $\mathcal{A}_\epsilon(t)$ is equal, up to order 15, to the simple *rational* expression:

$$\mathcal{A}_3(t) = \frac{t}{1 - t} \quad (45)$$

which is in perfect agreement with the result for the “real dynamical zeta function” $\zeta_{\epsilon=3}^{real}(t)$ (see (33)). For ϵ very close to 3 one gets :

$$\mathcal{A}_{\epsilon \simeq 3}(t) \simeq \frac{t}{1 - t} + \frac{t^3}{1 - t^3} = \frac{t \cdot (1 + t + 2t^2)}{1 - t^3} \quad (46)$$

again in good agreement with (34).

Integrable values for ϵ and around. For $\epsilon = 1/2$ the generating function for the “real Arnold complexity”, $\mathcal{A}_\epsilon(t)$, is equal, up to order 38, to the expansion of the rational expression :

$$\mathcal{A}_{1/2}(t) = \frac{t(1-t^7)}{(1-t)^2(1-t^5)(1+t)} + \frac{t^4(1-t^9)}{(1-t)(1-t^5)(1-t^6)(1+t)} + \frac{t^5}{(1-t^5)(1-t^3)(1+t)} + 2\frac{t^{28}}{1-t^5} \quad (47)$$

to be compared with $A_{1/2}(t)$ given in (3).

For $\epsilon = 1/3$ the calculations corresponding to the generating function for the “real Arnold complexity” are, in contrast, quite trivial yielding :

$$\mathcal{A}_{1/3}(t) = \frac{t \cdot (1+t)}{1-t^3} = A_{1/3}(t) \quad (48)$$

For $\epsilon = -1$ the generating function for the “real Arnold complexity” is equal, up to order 15, to the rational expression :

$$\mathcal{A}_{-1}(t) = \frac{t}{1-t^2} = A_{-1}(t) \quad (49)$$

For $\epsilon = +1$ the generating function for the “real Arnold complexity” is equal, up to order 15, to the rational expression :

$$\mathcal{A}_1(t) = \frac{t}{(1-t^2) \cdot (1-t)} = A_1(t) \quad (50)$$

all these results have to be compared with the generating functions (3).

Non-generic values for ϵ and around. The non-generic values of ϵ require a special attention. For instance for $\epsilon = 1/4$ one obtains the following expansion¹⁷ :

$$\begin{aligned} \mathcal{A}_{1/4}(t) = & t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 8t^7 + 11t^8 + 17t^9 + 23t^{10} + 31t^{11} + 44t^{12} + 63t^{13} \\ & + 90t^{14} + 128t^{15} + 183t^{16} + \dots \end{aligned}$$

and for $\epsilon = 1/5$ one gets :

$$\begin{aligned} \mathcal{A}_{1/5}(t) = & t + t^2 + 2t^3 + 2t^4 + 4t^5 + 4t^6 + 6t^7 + 7t^8 + 12t^9 + 15t^{10} + 19t^{11} + 28t^{12} + 33t^{13} \\ & + 53t^{14} + 77t^{15} + \dots \end{aligned} \quad (51)$$

Since $\epsilon = 1/5$ is a non-generic value (it is of the form $1/m$), the previous expansion (51) can be compared with the ones corresponding to values very close to $1/5$ but not equal, for instance $\epsilon = 99/500$, and $\epsilon = 101/500$:

$$\mathcal{A}_{99/500}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 18t^9 + 27t^{10} + 37t^{11} + 62t^{12} + 89t^{13} + \dots$$

and :

$$\mathcal{A}_{101/500}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 18t^9 + 27t^{10} + 41t^{11} + 66t^{12} + 85t^{13} + \dots$$

Closer to $\epsilon = 1/5$ one obtains for $\epsilon = 999/5000$, and $\epsilon = 1001/5000$, the following expansions :

$$\mathcal{A}_{999/5000}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 18t^9 + 27t^{10} + 41t^{11} + 66t^{12} + 89t^{13} + \dots$$

and :

$$\mathcal{A}_{1001/5000}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 18t^9 + 27t^{10} + 41t^{11} + 66t^{12} + 89t^{13} + \dots$$

Similar expansions, corresponding to values close to the non-generic value $\epsilon = 1/4$, are given in Appendix D. All these results show that, similarly to the situation for the customary topological entropy, or the growth complexity λ

¹⁷These maple calculations have been performed with 6000 digits, but they are already stable with 2000 digits.

(see Fig.1), λ_{real} is continuous as a function of ϵ near the non-generic values of $\epsilon \simeq 1/m$, however exactly on these very non-generic values λ_{real} takes *smaller* values (continuous function up to a zero measure set).

Remark : It is natural to compare the expansion corresponding to $\epsilon = 3/5$ with the one corresponding to $\epsilon = 1/7$, since $\epsilon = 1/7$ and $\epsilon = 3/5$ have the *same topological entropy* (growth complexity λ) associated with $1 - t - t^2 + t^{m+2}$ for $m = 7$ (see relation (11)). One gets for $\epsilon = 3/5$:

$$\mathcal{A}_{3/5}(t) = t + t^2 + 2t^3 + 3t^4 + 2t^5 + 5t^6 + 9t^7 + 8t^8 + 11t^9 + 14t^{10} + 18t^{11} + 24t^{12} + 29t^{13} + 41t^{14} + 51t^{15} + \dots$$

and for $\epsilon = 1/7$:

$$\mathcal{A}_{1/7}(t) = t + t^2 + 2t^3 + t^4 + 4t^5 + 7t^6 + 7t^7 + 8t^8 + 13t^9 + 16t^{10} + 22t^{11} + 36t^{12} + 43t^{13} + 65t^{14} + 87t^{15} + \dots$$

These two expansions *do not seem to yield the same value for λ_{real}* ($\lambda_{real}(3/5) \simeq 51^{1/15} \simeq 1.2997$ and $\lambda_{real}(1/7) \simeq 87^{1/15} \simeq 1.3468$) *though they share the same growth complexity λ .*

The expansions of $\mathcal{A}_\epsilon(t)$ near $\epsilon = 3/5$ are given in Appendix D.

Miscellaneous values of ϵ . For most of the values of ϵ the expansions are not long enough to “guess” rational expressions (if any ...). One can however get some estimates of λ_{real} that can be compared with h_{real} .

- For $\epsilon = 21/25$ one gets the following results:

$$\mathcal{A}_{21/25}(t) = t + t^2 + 2t^3 + 3t^4 + 3t^5 + 6t^6 + 7t^7 + 11t^8 + 12t^9 + 21t^{10} + 25t^{11} + 36t^{12} + 45t^{13} + 69t^{14} + \dots$$

This expansions seem to yield the following estimated value for λ_{real} ($\lambda_{real}(21/25) \simeq 69^{1/14} \simeq 1.3531$ to be compared with (36)). In fact, one remarks that this expansion is *actually compatible* with the expansion of the rational expression:

$$\mathcal{A}_{21/25}(t) = \frac{t \cdot (1 + t + t^2 + t^3 - 2t^4)}{(1 - t)(1 + t)^2(1 - t + t^2 - 2t^3)} \quad (52)$$

One does remark that the rational expression (52) has actually the *same singularity* that the rational expression (37) suggested for the real dynamical zeta function $\zeta_{21/25}^{real}(t)$. All the coefficients of the expansion of (52) are positive (in contrast with (60) given in Appendix C which is ruled out because coefficient t^{54} of its expansion is negative). If this simple rational expression is actually the exact expression for $\mathcal{A}_{21/25}(t)$ this would yield the following algebraic value for λ_{real} : $\lambda_{real}(21/25) = h_{real}(21/25) \simeq 1.353209964$.

- For $\epsilon = 3/2$ one gets the following results:

$$\mathcal{A}_{3/2}(t) = t + t^2 + 2t^3 + t^4 + 3t^5 + 2t^6 + 3t^7 + 3t^8 + 2t^9 + 3t^{10} + 3t^{11} + 4t^{12} + 3t^{13} + \dots$$

- Near $\epsilon = 2$ (for instance for $\epsilon = 2001/1000$ or $\epsilon = 1999/1000$) one gets :

$$\mathcal{A}_{\epsilon \simeq 2}(t) = t + t^2 + 2t^3 + t^4 + t^5 + 2t^6 + t^7 + 3t^8 + 2t^9 + t^{10} + 3t^{11} + 2t^{12} + 3t^{13} + \dots$$

- Some results for ϵ larger than 3 (again obtained with 6000 digits) are given in Appendix D ($\epsilon = 4$, $\epsilon = 5$, $\epsilon = 6$, $\epsilon = 10$, $\epsilon = 20$, $\epsilon = 30$). These series indicate that an estimated value for λ_{real} could correspond to λ_{real} very close to 1 for $\epsilon = 10$, and quite close to 1 for $\epsilon = 20$.

B. “Real Arnold complexity” generating functions for ϵ large.

The examination of Fig.12 shows that λ_{real} goes to some non-trivial limit, $\lambda_{real} \simeq 1.429$, in the large ϵ limit. Let us examine the expansion of $\mathcal{A}_\epsilon(t)$ for various increasing values of ϵ , in order to study this $\epsilon \rightarrow \infty$ limit. The expansions of $\mathcal{A}_\epsilon(t)$ for $\epsilon = 40$, $\epsilon = 50$, $\epsilon = 100$ are given in Appendix D.

- For $\epsilon = 500$ the expansion of $\mathcal{A}_\epsilon(t)$ reads :

$$\mathcal{A}_{500}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 8t^6 + 11t^7 + 17t^8 + 24t^9 + 35t^{10} + 47t^{11} + 64t^{12} + 93t^{13} + \dots$$

- For $\epsilon = 1000$:

$$\mathcal{A}_{1000}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 8t^6 + 11t^7 + 17t^8 + 24t^9 + 35t^{10} + 51t^{11} + 72t^{12} + 101t^{13} + \dots$$

For ϵ large the expansion of $\mathcal{A}_\epsilon(t)$, the generating function for the “real Arnold complexity”, is equal, up to order 15, to (for instance¹⁸ for $\epsilon = 20000$) :

$$\begin{aligned} \mathcal{A}_{20000}(t) = & t + t^2 + 2t^3 + 3t^4 + 5t^5 + 8t^6 + 11t^7 + 17t^8 + 24t^9 + 35t^{10} + 51t^{11} + 72t^{12} + 105t^{13} \\ & + 149t^{14} + 214t^{15} + \dots \end{aligned} \quad (53)$$

which actually coincides with the expansion of the simple rational expression :

$$\mathcal{A}_\infty(t) = \frac{t \cdot (1 + t^4)}{(1 - t^2 - t^3 - t^5) \cdot (1 - t)} = \frac{t \cdot (1 + t^4)}{(1 - t - t^2) + t^4 \cdot (1 - t + t^2)} \quad (54)$$

This last result has to be compared with the equivalent one for the “real dynamical zeta function” $\zeta_{\epsilon=\infty}^{real}(t)$ (see (40) in section (IV D)). These two non-trivial rational results, for ϵ large, are in *perfect agreement*, yielding the same *algebraic value* for the two “real complexities” h_{real} and λ_{real} , namely $h_{real} = \lambda_{real} \simeq 1.42910832$.

All the results displayed in this section seem to show that the identification between h_{real} and λ_{real} actually holds.

Remark : Recalling the “universal” relation (18), or more precisely (19), which gives (for ϵ generic and for $\epsilon = 1/m$, $\epsilon = (m+1)/(m+3)$ for $m = 9, 13, 17, \dots$) a ratio $\zeta_\epsilon(t)/\mathcal{A}_\epsilon(t)$ equal to $(1-t^2)/t$, one can look at the “real ratio” $\zeta_\epsilon^{real}(t)/\mathcal{A}_\epsilon(t)$. Of course for $\epsilon < 0$ this “real ratio” is also equal to $(1-t^2)/t$, however in the $\epsilon \rightarrow \infty$ limit it tends to be equal to $(1-t^2)/t/(1+t^4)$. Therefore one should not expect any simple “universal” relation like (18) between $\zeta_\epsilon^{real}(t)$ and $\mathcal{A}_\epsilon(t)$.

These various Arnold complexity generating functions $\mathcal{A}_\epsilon(t)$ were associated with the iteration of the (real or complex) line $y = (1-\epsilon)/2$. One could introduce a generating function for each line (or fixed curve) one iterates. The corresponding series become slightly more difficult to extrapolate but give similar results in particular the asymptotic values for λ_{real} . The sensitivity of the previous analysis, according to the chosen curve one iterates, will be discussed elsewhere.

It would be interesting to see if the “real dynamical zeta functions” $\zeta_\epsilon^{real}(t)$, or the “real degree generating functions” $\mathcal{A}_\epsilon(t)$, could also be *rational* expressions, for *other* values of ϵ , or, even, if these “real generating functions” could be *rational* expressions for *any given* value of ϵ . In this last case there should be an infinite number of such *rational* expressions : it is clear that they could not all be “simple” like (40) or (54).

VII. CONCLUSION

The results presented here seem to be *in agreement with, again, an identification between λ_{real} , the (asymptotic) “real Arnold complexity”, and h_{real} , the (exponential of the) “real topological entropy”*. In contrast with the “universal” behavior of the “usual” Arnold complexity, or topological entropy, displayed in Fig. 1, λ_{real} and h_{real} are quite involved functions of the parameter ϵ the birational transformations depend on (see Fig. 11 and Fig. 12).

We have, however, obtained some remarkable *rational expressions for the real dynamical zeta function $\zeta_\epsilon^{real}(t)$ and for a “real Arnold complexity” generating function $\mathcal{A}_\epsilon(t)$* . In particular we have obtained in particular two non-trivial *rational* expressions (40) and (54), (yielding algebraic values for h_{real} and λ_{real}).

There is no simple down-to-earth Markov partition, symbolic dynamics, or hyperbolic systems interpretation of these rational results.

ACKNOWLEDGMENTS

One of us (JMM) would like to thank P. Lochak and J-P. Marco for illuminating discussions on symbolic dynamics and on non hyperbolic discrete dynamical systems. N. Abarenkova would like to thank the St Petersburg’s administration for a grant. S. Boukaraa would like to thank the CMEP for financial support.

¹⁸These calculations have to be performed with at least 6000 digits. With a number of digits lower than 2000 digits one gets *smaller* coefficients: the precision is not large enough to distinguish between very close intersection points.

VIII. APPENDIX A : DYNAMICAL ZETA FUNCTIONS VERSUS HOMOGENEOUS DEGREE GENERATING FUNCTION FOR NON-GENERIC VALUES

Let us consider here various non-generic values of the form $(m-1)/(m+3)$ (with $m \geq 7$, m odd).

- For $\epsilon = 3/5$ (corresponding to $m = 7$) the homogeneous generating function defined in section (III), reads :

$$G_{3/5}^{Hom}(t) = 1 + 2t + 4t^2 + 7t^3 + 12t^4 + 20t^5 + 33t^6 + 54t^7 + 88t^8 + 142t^9 + 228t^{10} + 366t^{11} + \dots$$

which is compatible with the expansion of the rational expression :

$$G_{3/5}^{Hom}(t) = \frac{1 - t^{10}}{(1-t) \cdot (1-t-t^2+t^9)}$$

Recalling a possible rational expression for the corresponding dynamical zeta function [3] :

$$\zeta_{3/5}(t) = \frac{1-t^2}{1-t-t^2+t^9}, \quad (55)$$

one immediately verifies that the “universal” relation (18) actually holds.

- For $\epsilon = 2/3$ (corresponding to $m = 9$) the homogeneous generating function defined in section (III), reads :

$$G_{2/3}^{Hom}(t) = 1 + 2t + 4t^2 + 7t^3 + 12t^4 + 20t^5 + 33t^6 + 54t^7 + 88t^8 + 143t^9 + 232t^{10} + 375t^{11} + 605t^{12} + \dots$$

This could be the expansion, up to order twelve, of the simple rational expression :

$$G_{2/3}^{Hom}(t) = \frac{1 - t^{12}}{(1-t) \cdot (1-t-t^2+t^{11})} \quad (56)$$

These results should be compared with the expansion of the dynamical zeta function. Unfortunately, here, the series for the dynamical zeta function are not sufficiently large to allow any “safe conjecture”. A possible exact expression does not seem to be equal to $(1-t^2)/(1-t-t^2+t^{11})$, but could be [3] :

$$\zeta_{2/3}(t) = \frac{1 - t^2 - t^{11} - t^{12} - t^{13}}{1 - t - t^2 + t^{11}} \quad \text{or :} \quad \frac{1 - t^2 - t^{11} - t^{12}}{1 - t - t^2 + t^{11}} \quad (57)$$

The “universal” relation (18) is verified with (56) together with $(1-t^2)/(1-t-t^2+t^{11})$, *but not with* (56) together with (57). One can however imagine that the “universal” relation (18) could be slightly modified on some of these $(m-1)/(m+3)$ values ($m = 9, 13, \dots$). For instance, (56) and (57) verify (up to order twelve) the simple relation :

$$t \cdot \zeta_{2/3}(t) - (1-t) \cdot G_{2/3}^{Hom}(t) + 1 - t^{m+2} - t^{m+3} = 0 \quad \text{where :} \quad m = 9 \quad (58)$$

These calculations need to be revisited.

- For $\epsilon = 5/7$ (corresponding to $m = 11$) the homogeneous generating function defined in section (III), reads :

$$G_{5/7}^{Hom}(t) = 1 + 2t + 4t^2 + 7t^3 + 12t^4 + 20t^5 + 33t^6 + 54t^7 + 88t^8 + 143t^9 + 232t^{10} + 376t^{11} + 609t^{12} + \dots$$

This could be the expansion of :

$$G_{5/7}^{Hom}(t) = \frac{1 - t^{14}}{(1-t) \cdot (1-t-t^2+t^{13})}$$

- For $\epsilon = 3/4$ (corresponding to $m = 13$) the homogeneous generating function reads :

$$G_{3/4}^{Hom}(t) = 1 + 2t + 4t^2 + 7t^3 + 12t^4 + 20t^5 + 33t^6 + 54t^7 + 88t^8 + 143t^9 + 232t^{10} + 376t^{11} + 609t^{12} + \dots$$

This series is not large enough. It could be the expansion of the simple expression :

$$G_{3/4}^{Hom}(t) = \frac{1 - t^{16}}{(1-t) \cdot (1-t-t^2+t^{15})}$$

IX. APPENDIX B : NUMBER OF REAL FIXED POINTS OF THE P -TYPE, Q -TYPE AND R -TYPE.

Let us just give the number of *real* n -th cycles of the P -type, Q -type and R -type for miscellaneous values of ϵ in increasing order.

For $\epsilon < 0$ (and $\epsilon \neq -1$) one gets :

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
P_n	1	0	1	1	2	2	4	4	6	8	12	12	20	24	30	38	54	65
Q_n	0	0	0	0	0	0	0	1	0	1	0	3	0	4	0	9	0	13
R_n	0	0	0	0	0	0	0	0	2	2	6	10	20	30	60	88	156	238
T_n	1	0	1	1	2	2	4	5	8	11	18	25	40	58	90	135	210	316

TABLE I. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon < 0$.

For $\epsilon = 11/100, 1/4, 52/100, 9/10, 11/10, 5, 10, 50, 100$, and 20000, one gets the following tables :

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
P_n	1	0	1	1	2	1	2	1	4	5	8	5	10	11	14	14	20	21
Q_n	0	0	0	0	0	0	0	0	0	1	0	3	0	3	0	5	0	5
R_n	0	0	0	0	0	0	0	0	0	0	*	*	*	*	*	*	*	*
T_n	1	0	1	1	2	1	2	1	4	6	*	*	*	*	*	*	*	*

TABLE II. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 11/100$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13
P_n	1	0	1	1	2	0	1	0	3	1	4	2	8
Q_n	0	0	0	0	0	0	0	1	0	1	0	2	0
R_n	0	0	0	0	0	0	0	0	0	0	0	0	0
T_n	1	0	1	1	2	0	1	1	3	2	4	4	8

TABLE III. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 1/4$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
P_n	1	0	1	1	2	0	2	0	4	1	4	2	6	2	8
Q_n	0	0	0	0	0	0	0	1	0	1	0	2	0	4	0
R_n	0	0	0	0	0	0	0	0	0	0	2	8	10	*	*
T_n	1	0	1	1	2	0	2	1	4	2	6	12	16	*	*

TABLE IV. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 52/100$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
P_n	1	0	1	1	0	0	2	0	0	1	4	0	2	1	6	1	6	3
Q_n	0	0	0	0	0	0	0	1	0	1	0	2	0	2	0	4	0	5
R_n	0	0	0	0	0	0	0	0	0	0	*	*	*	*	*	*	*	*
T_n	1	0	1	1	0	0	2	1	0	2	*	*	*	*	*	*	*	*

TABLE V. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 9/10$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
P_n	1	0	1	0	0	0	2	0	0	1	2	0	2	1	2	1	4
Q_n	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0
R_n	0	0	0	0	0	0	0	0	0	0	0	*	*	*	*	*	*
T_n	1	0	1	0	0	0	2	0	0	2	2	*	*	*	*	*	*

TABLE VI. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 11/10$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
P_n	1	0	1	0	0	0	0	0	0	0	2	0	*	1	0	0	2	0
Q_n	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0
R_n	0	0	0	0	0	0	0	0	0	0	0	*	*	*	*	*	*	*
T_n	1	0	1	0	0	0	0	0	0	0	2	*	*	*	*	*	*	*

TABLE VII. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 5$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
P_n	1	0	1	0	2	0	0	1	0	0	2	0	2	1	0	0	2	1
Q_n	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	1
R_n	0	0	0	0	0	0	0	0	0	0	0	*	*	*	*	*	*	*
T_n	1	0	1	0	2	0	0	2	0	0	2	*	*	*	*	*	*	*

TABLE VIII. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 10$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
P_n	1	0	1	0	2	0	2	1	2	0	4	1	4	1	2	2	6	3
Q_n	0	0	0	0	0	0	0	1	0	1	0	1	0	2	0	3	0	3
R_n	0	0	0	0	0	0	0	0	0	2	0	*	*	*	*	*	*	*
T_n	1	0	1	0	2	0	2	2	2	3	4	*	*	*	*	*	*	*

TABLE IX. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 50$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
P_n	1	0	1	0	2	0	2	1	2	0	4	1	6	1	6	2	8	3
Q_n	0	0	0	0	0	0	0	1	0	1	0	3	0	2	0	3	0	6
R_n	0	0	0	0	0	0	0	0	0	2	0	*	*	*	*	*	*	*
T_n	1	0	1	0	2	0	2	2	2	3	4	*	*	*	*	*	*	*

TABLE X. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 100$.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
P_n	1	0	1	0	2	0	2	1	2	0	4	1	6	1	6	2
Q_n	0	0	0	0	0	0	0	1	0	1	0	3	0	4	0	7
R_n	0	0	0	0	0	0	0	0	0	2	0	2	*	*	*	*
T_n	1	0	1	0	2	0	2	2	2	3	4	6	≥ 8	≥ 9	≥ 14	≥ 17

TABLE XI. Number of *real* n -th cycles of the P -type, Q -type and R -type for $\epsilon = 20000$.

X. APPENDIX C : EXPANSIONS OF SOME REAL DYNAMICAL ZETA FUNCTIONS

Let us just give some additional expansions for $\zeta_\epsilon^{real}(t)$ for increasing values of ϵ .
For $\epsilon = 9/50$, one obtains the following expansions for $\zeta_\epsilon^{real}(t)$:

$$\begin{aligned}\zeta_{9/50}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^5)^2(1-t^6)(1-t^7)^2(1-t^8)^3(1-t^9)^4(1-t^{10})^3(1-t^{11})^8} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 5t^5 + 7t^6 + 10t^7 + 16t^8 + 24t^9 + 34t^{10} + 52t^{11} + \dots\end{aligned}$$

yielding the following “rough” approximation for h_{real} : $h_{real} \simeq (52)^{1/11} \simeq 1.432$. For the “non-generic” value $\epsilon = 1/5$, $\zeta_{1/5}^{real}(t)$ reads :

$$\begin{aligned}\zeta_{1/5}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^5)^2(1-t^6)(1-t^7)(1-t^9)^4(1-t^{10})^2(1-t^{11})^5(1-t^{12})^4} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 5t^5 + 7t^6 + 9t^7 + 12t^8 + 20t^9 + 28t^{10} + 39t^{11} + 55t^{12} + \dots\end{aligned}$$

yielding the following “rough” approximation for h_{real} : $h_{real} \simeq (55)^{1/12} \simeq 1.3964$. For $\epsilon = 1/5$ the previous Q_n ’s and R_n ’s are equal to zero up to order twelve. The exponents in (59) are thus the P_n ’s.

For $\epsilon = 31/125$, $\epsilon = 12/25$, $\epsilon = 66/125$, $\epsilon = 2/3$, $\epsilon = 17/25$, $\epsilon = 3/4$, $\epsilon = 3/2$, one obtains, respectively, the following expansions for $\zeta_\epsilon^{real}(t)$:

$$\begin{aligned}\zeta_{31/125}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^5)^2(1-t^6)(1-t^7)^2(1-t^8)(1-t^9)^4(1-t^{10})^5(1-t^{11})^{12}} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 5t^5 + 7t^6 + 10t^7 + 14t^8 + 22t^9 + 34t^{10} + 54t^{11} + \dots\end{aligned}$$

yielding the approximation for h_{real} : $h_{real} \simeq (54)^{1/11} \simeq 1.437$,

$$\begin{aligned}\zeta_{12/25}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^5)^2(1-t^7)^2(1-t^8)(1-t^9)^6(1-t^{10})^5(1-t^{11})^{10}} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 22t^9 + 33t^{10} + 49t^{11} + \dots\end{aligned}$$

for $\epsilon = 12/25$, yielding the following rough approximation for h_{real} : $h_{real} \simeq (49)^{1/11} \simeq 1.424$,

$$\begin{aligned}\zeta_{66/125}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^7)^2(1-t^8)(1-t^9)^4(1-t^{10})^2(1-t^{11})^4} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 3t^5 + 4t^6 + 7t^7 + 9t^8 + 14t^9 + 19t^{10} + 27t^{11} + \dots\end{aligned}$$

for $\epsilon = 66/125$, yielding $h_{real} \simeq (27)^{1/11} \simeq 1.349$.

For $\epsilon = 2/3$ (that is $(m-1)/(m+3)$ for $m = 9$) the real dynamical zeta function reads :

$$\begin{aligned}\zeta_{2/3}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^7)^2(1-t^8)(1-t^9)^2(1-t^{10})^2(1-t^{11})^4(1-t^{12})^2} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 3t^5 + 4t^6 + 7t^7 + 9t^8 + 12t^9 + 17t^{10} + 25t^{11} + 32t^{12} + \dots\end{aligned} \tag{59}$$

yielding $h_{real} \simeq (32)^{1/12} \simeq 1.3348$. Let us note that one must be careful converting systematically a series to a rational function (Pade approximation). Up to order twelve, expansion (59) is in agreement with the expansion of the following simple rational expression :

$$\frac{1 + t + t^3 - t^6}{1 - t^2 - 2t^4 + t^5 - t^6} = \frac{(1 + t + t^3 - t^6) \cdot (1 - t)}{1 - t - t^2 + t^3 \cdot (1 - t + t^2)^2} \tag{60}$$

which is reminiscent of the exact expression (40). However, one easily finds that the coefficients of t^{54} in (60) becomes negative (the coefficients grow like $\simeq (-1.5252)^N$). Expression (60) *cannot* be the exact expression of a (real) dynamical zeta function.

For $\epsilon = 17/25$, the real dynamical zeta function reads :

$$\begin{aligned}\zeta_{17/25}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^7)^2(1-t^8)(1-t^9)^2(1-t^{10})^2(1-t^{11})^4} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 3t^5 + 4t^6 + 7t^7 + 9t^8 + 12t^9 + 17t^{10} + 25t^{11} + \dots\end{aligned}$$

yielding : $h_{real} \simeq (25)^{1/11} \simeq 1.3399$,

For the non-generic value $\epsilon = 3/4$ (that is $(m-1)/(m+3)$ for $m = 13$) the real dynamical zeta function reads :

$$\begin{aligned}\zeta_{3/4}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^4)(1-t^7)^2(1-t^8)(1-t^9)^3(1-t^{10})(1-t^{11})^2(1-t^{12})} \dots \\ &= 1 + t + t^2 + 2t^3 + 3t^4 + 3t^5 + 4t^6 + 7t^7 + 9t^8 + 13t^9 + 17t^{10} + 23t^{11} + 30t^{12} + \dots\end{aligned}$$

yielding $h_{real} \simeq (30)^{1/12} \simeq 1.3277$.

Finally, for $\epsilon = 3/2$, the real dynamical zeta function reads :

$$\begin{aligned}\zeta_{3/2}^{real}(t) &= \frac{1}{(1-t)(1-t^3)(1-t^7)^2(1-t^{10})^2} \dots \\ &= 1 + t + t^2 + 2t^3 + 2t^4 + 2t^5 + 3t^6 + 5t^7 + 5t^8 + 6t^9 + 10t^{10} + 10t^{11} + \dots\end{aligned}$$

yielding : $h_{real} \simeq (10)^{1/11} \simeq 1.233$.

XI. APPENDIX D : EXPANSIONS OF THE “REAL ARNOLD COMPLEXITY” GENERATING FUNCTIONS

Let us give here a few expansions for the “real Arnold complexity” generating functions $\mathcal{A}_\epsilon(t)$. Let us first give the expansion of $\mathcal{A}_\epsilon(t)$ corresponding to $\epsilon = 2/3$ in order to compare it with (59) and (60) :

$$\mathcal{A}_{2/3}(t) = t + t^2 + 2t^3 + 3t^4 + 3t^5 + 6t^6 + 7t^7 + 11t^8 + 14t^9 + 21t^{10} + 29t^{11} + 37t^{12} + 51t^{13} + \dots$$

yielding the following estimation for $\lambda_{real} \simeq (51)^{1/13} \simeq 1.3531$ to be compared with $h_{real} \simeq (32)^{1/12} \simeq 1.3348$ from (59). The coefficients of the expansions of $\zeta_{2/3}(t)$, and $\mathcal{A}_{2/3}(t)$, are very close. Up to order ten, the ratio $\zeta_{2/3}(t)/\mathcal{A}_{2/3}(t)$ coincides with the expansion of :

$$\frac{\zeta_{2/3}(t)}{\mathcal{A}_{2/3}(t)} \simeq \frac{1-t^2}{t} \cdot \frac{1-t^5}{1-2t^4+t^5} \quad (61)$$

The expansion of $\mathcal{A}_\epsilon(t)$ corresponding to the non-generic value $\epsilon = 3/4$ (that is $(m-1)/(m+3)$ for $m = 13$) reads:

$$\mathcal{A}_{3/4}(t) = 1 + t + t^2 + 2t^3 + 3t^4 + 3t^5 + 6t^6 + 7t^7 + 11t^8 + 12t^9 + 21t^{10} + 27 \cdot t^{11} + 36t^{12} + 47t^{13} + \dots \quad (62)$$

The expansions of $\zeta_{3/4}(t)$ and $\mathcal{A}_{3/4}(t)$ are again very close. Expansion (62) yields the following estimation for $\lambda_{real} \simeq (47)^{1/13} \simeq 1.3446$ to be compared with $h_{real} \simeq (30)^{1/12} \simeq 1.3276$ from (61).

Let us now give the expansion of $\mathcal{A}_\epsilon(t)$ corresponding to values very close to the non-generic value $1/4$, for instance $\epsilon = 99/400$ and $\epsilon = 101/400$:

$$\mathcal{A}_{99/400}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 22t^9 + 33t^{10} + 47t^{11} + 70t^{12} + 101t^{13} + \dots$$

and :

$$\mathcal{A}_{101/400}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 6t^6 + 9t^7 + 13t^8 + 22t^9 + 33t^{10} + 47t^{11} + 70t^{12} + 109t^{13} + \dots$$

Near the non-generic value $\epsilon = 3/5$, for instance for $\epsilon = 299/500$ and $\epsilon = 301/500$, one gets :

$$\mathcal{A}_{299/500}(t) = t + t^2 + 2t^3 + 3t^4 + 3t^5 + 6t^6 + 11t^7 + 11t^8 + 16t^9 + 21t^{10} + 29t^{11} + 42t^{12} + 57t^{13} + \dots$$

and :

$$\mathcal{A}_{301/500}(t) = t + t^2 + 2t^3 + 3t^4 + 3t^5 + 6t^6 + 11t^7 + 11t^8 + 16t^9 + 21t^{10} + 29t^{11} + 42t^{12} + 57t^{13} + \dots$$

For the non-generic value $\epsilon = 1/10$ one obtains the following results :

$$\mathcal{A}_{1/10}(t) = t + t^2 + 2t^3 + t^4 + 3t^5 + 8t^6 + 9t^7 + 11t^8 + 16t^9 + 21t^{10} + 31t^{11} + 48t^{12} + 58t^{13} + \dots$$

• $\epsilon > 3$. Let us finally give some results for ϵ larger than 3 (again obtained with 6000 digits).

For $\epsilon = 4$:

$$\mathcal{A}_4(t) = t + t^2 + 2t^3 + t^4 + t^5 + 2t^6 + t^7 + t^8 + 2t^9 + 3t^{10} + t^{11} + 2t^{12} + 3t^{13} + t^{14} + \dots$$

For $\epsilon = 5$:

$$\mathcal{A}_5(t) = t + t^2 + 2t^3 + t^4 + t^5 + 2t^6 + 3t^7 + t^8 + 2t^9 + 3t^{10} + t^{11} + 2t^{12} + 3t^{13} + 3t^{14} + \dots$$

For $\epsilon = 6$:

$$\mathcal{A}_6(t) = t + t^2 + 2t^3 + 3t^4 + t^5 + 2t^6 + 3t^7 + 3t^8 + 2t^9 + 3t^{10} + 3t^{11} + 4t^{12} + 3t^{13} + \dots$$

For $\epsilon = 10$:

$$\mathcal{A}_{10}(t) = t + t^2 + 2t^3 + 3t^4 + 3t^5 + 2t^6 + 3t^7 + 3t^8 + 4t^9 + 5t^{10} + 5t^{11} + 4t^{12} + 5t^{13} + \dots$$

For $\epsilon = 20$:

$$\mathcal{A}_{20}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 4t^6 + 5t^7 + 3t^8 + 8t^9 + 11t^{10} + 7t^{11} + 10t^{12} + 21t^{13} + \dots$$

For $\epsilon = 30$:

$$\mathcal{A}_{30}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 4t^6 + 5t^7 + 9t^8 + 8t^9 + 11t^{10} + 11t^{11} + 14t^{12} + 25t^{13} + \dots$$

For $\epsilon = 40$:

$$\mathcal{A}_{40}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 4t^6 + 5t^7 + 9t^8 + 10t^9 + 15t^{10} + 11t^{11} + 14t^{12} + 29t^{13} + \dots$$

For $\epsilon = 50$:

$$\mathcal{A}_{50}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 4t^6 + 7t^7 + 9t^8 + 10t^9 + 15t^{10} + 17t^{11} + 22t^{12} + 37t^{13} + \dots$$

For $\epsilon = 100$:

$$\mathcal{A}_{100}(t) = t + t^2 + 2t^3 + 3t^4 + 5t^5 + 8t^6 + 7t^7 + 9t^8 + 16t^9 + 19t^{10} + 29t^{11} + 36t^{12} + 51t^{13} + \dots$$

- [1] N. Abarenkova, J-C. Anglès d'Auriac, S. Boukraa, S. Hassani and J-M. Maillard, *Topological entropy and complexity for discrete dynamical systems*. *chao-dyn/9806026* .
- [2] N. Abarenkova, J-C. Anglès d'Auriac, S. Boukraa, S. Hassani and J-M. Maillard, *Rational dynamical zeta functions for birational transformations*. *chao-dyn/9807014* to be published in *Physica A*.
- [3] N. Abarenkova, J-C. Anglès d'Auriac, S. Boukraa, S. Hassani and J-M. Maillard, *From Yang-Baxter equations to dynamical zeta functions for birational transformations*, to be published by World Scientific
- [4] N. Abarenkova, J-C. Anglès d'Auriac, S. Boukraa and J-M. Maillard, *Growth complexity spectrum of some discrete dynamical systems*. *chao-dyn/9807031* to be published in *Physica D*.
- [5] S. Boukraa and J-M. Maillard, *Factorization properties of birational mappings*. *Physica A* **220**, 403 (1995).
- [6] V. Arnold, *Developments in Mathematics: The Moscow School*. Chapman & Hall, (1989). Problems on singularities and dynamical systems, Edited by V. Arnold and M. Monastyrsky, Chap. 7, pp. 261-274.
- [7] S. Boukraa, S. Hassani and J-M. Maillard, *Product of involutions and fixed points*. *Alg. Rev. Nucl. Sci.* **2** (1998) pp. 1–16.
- [8] S. Boukraa, J-M. Maillard and G. Rollet, *Almost integrable mappings*. *Int. J. Mod. Phys. B* **8** (1994), pp. 137–174.

- [9] S. Hassani, S. Boukraa and J-M. Maillard, *New integrable cases of a Cremona transformation : a finite order orbit analysis*. Physica **A 240** (1997), p. 586.
- [10] S. Boukraa, J-M. Maillard and G. Rollet, *Determinantal identities on integrable mappings*. Int. J. Mod. Phys. **B8** (1994), pp. 2157–2201.
- [11] S. Boukraa, J-M. Maillard and G. Rollet, *Integrable mappings and polynomial growth*. Physica A **208** (1994), pp. 115–175.
- [12] R.C. Adler, A.C. Konheim and M.H. McAndrew, *Topological entropy*. Trans. Am. Math. Soc. **114** (1965), p. 309.
- [13] V. Baladi, *Periodic orbits and dynamical spectra*. Ergod. Th. Dynam. Sys. **18**, 255-292 (1998).
- [14] Information on the multi-precision library gmp (GNU MP) can found in the following Home Site :
<http://www.nada.kth.se/~tege/gmp/>
 Other Links : [ftp://prep.ai.mit.edu/pub/gnu/\(gmp-*.tar.gz\)](ftp://prep.ai.mit.edu/pub/gnu/(gmp-*.tar.gz))
 Mailing Lists/USENET News Groups : bug-gmp@prep.ai.mit.edu
- [15] N. Abarenkova, J-C. Anglès d'Auriac, S. Boukraa and J-M. Maillard, *Elliptic curves from finite order recursions or non-involutive permutations for discrete dynamical systems and lattice statistical mechanics*. The European Physical Journal **B 5** (1998) pp. 647–661.
- [16] R. Bowen, *Periodic Orbits for hyperbolic flows*, Amer. J. Math. **94**, (1972), pp 1–30.
- [17] K.T. Alligood, T.D. Sauer and J.A. Yorke, *Chaos An introduction to dynamical systems*. Springer-Verlag New York (1997).
- [18] *Chaos Classical and Quantum* : a Web book to be found in Cvitanovic's web site <http://www.nbi.dk/ChaosBook>
- [19] M. Artin and B. Mazur, *On periodic points*. Ann. of Math. **81** (1965), p. 82.
- [20] D. Ruelle, *Thermodynamic Formalism*. Addison-Wesley, Reading, MA, (1978).
- [21] R. Bowen, *Symbolic dynamics for hyperbolic flows*. Amer. J. Math. **95** (1973), p. 429.
- [22] R. Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*. Springer Verlag, Berlin, (1975).
- [23] R. Bowen, *Markov partitions for Axiom A diffeomorphisms*, Amer. J. Math. **92**, (1970), pp 725–747.
- [24] C. Smale, *Differentiable dynamical systems*. Bull. Amer. Math. Soc. **73** (1967), p. 747.
- [25] N. Abarenkova, J-C. Anglès d'Auriac, S. Boukraa, S. Hassani and J-M. Maillard, *Cycle fusion for birational transformations*, in preparation.
- [26] J-C. Anglès d'Auriac, S. Boukraa, and J-M. Maillard, *Functional relations in lattice statistical mechanics, enumerative combinatorics and discrete dynamical systems*, to be published in Annals of Combinatorics.
- [27] R. W. Ghrist, P. J. Holmes and M. C. Sullivan, *Knots and Links in Three-Dimensional Flows*. Lecture Notes in Mathematics, **1654**, (1998), Springer.
- [28] R. W. Easton, *Geometrical Methods for Discrete Dynamical Systems*. Oxford Engineering Science Series, **50**, (1998).
- [29] C. Conley and R. Easton, *Isolated invariant sets and isolated blocks*. Trans. Amer. Math. Soc. **158** (1971), p. 35.
- [30] D. Fried, *Rationality for Isolated Expansive Sets*. Advances in Mathematics **65** (1987), p. 35.
- [31] M. Marden, *Geometry of polynomials*., Amer. Math. Soc. Surveys, second edition (1966).