## A New Class of Cellular Autom ata for Reaction-Diusion Systems

Jorg R.W eim ar and Jean-Pierre Boon Faculte des Sciences, C.P.231 Universite Libre de Bruxelles, B-1050 Bruxelles, Belgium E-m ail: jweimar@ulb.ac.be, jpboon@ulb.ac.be May 22, 1993)

We introduce a new class of cellular autom ata to model reaction-di usion systems in a quantitatively correct way. The construction of the CA from the reaction-di usion equation relies on a moving average procedure to im plem ent diffusion, and a probabilistic table-lookup for the reactive part. The applicability of the new CA is demonstrated using the G inzburg-Landau equation.

82.20 W t, 02.70 Rw, 05.50.+ q

Cellular autom ata m odels have been used in m any applications to model reactive and di usive system s [1,2]. Most uses of cellular autom ata (CAs) can be classi ed into one of four approaches: (i) Ising-type models of phase transitions; (ii) lattice gas models (the lattice gas m ethod was initially developed to m odel hydrodynam ic ows and has been extended in many directions [3,4]; (iii) system atic investigation of the behavior of CAs by investigating all rules of a certain class (e.g., all possible rules for one-dimensional automata with two states and nearest neighbor interaction) [1]; and (iv) qualitative discrete m odelling (including operational use of CAs as an alternative to partial di erential equations (PDEs) [5]). These models are generally based on qualitative rather than quantitative inform ation about the system to be modelled. A CA is constructed which preserves the qualitative features deem ed most relevant and it is then investigated (The lattice gas methods were also developed along these lines [6]). Existing CA models for reaction-diusion systems [7{11] fall into the category (iv), i.e., they show qualitatively \correct" behavior and are restricted to certain reaction-di usion (R-D) models and certain types of phenomena. This is the main criticism of experimentalists and researchers working with partialdi erential equation m odels, who search for quantitative predictions. In this letter we describe a class of CAs which is suitable for modelling many reactiondi usion systems in a quantitatively correct way. The new CAs are operationally more e cient than the reactive lattice gas m ethods, which also achieve quantitative correctness. We rst describe the construction of the new class of CAs; then we present the autom aton using the G inzburg-Landau equation as an example.

The main idea behind this class of CAs is careful discretization. Space and time are discretized as in normal

nite di erence methods for solving the PDE's. Finite di erence methods then proceed to solve the resulting coupled system of N s ordinary di erential equations (N points in space, s equations in the PDE system) by any of a number of num erical methods, operating on oating point num bers. The use of oating point num bers on computers in plies a discretization of the continuous variables. The errors introduced by this discretization and the ensuing roundo errors are often not considered explicitly, but assumed to be small because the precision is rather high (8 decim aldigits for usual oating point num bers). In contrast, in the CA approach, all variables are explicitly discretized into relatively small integers. This discretization allows the use of lookup tables to replace the evaluation of the nonlinear rate functions. It is this table lookup, combined with the fact that all calculations are perform ed using integers instead of oating point variables, that accounts for an im provement in speed of orders of magnitude on a conventional multi-purpose com puter. The undesirable e ects of discretization are overcom e by using probabilistic rules for the updating of the CA.

The state of the CA is given by a regular array of concentration vectors y residing on a d-dimensional lattice. Each y (r) is a s-vector of integers (s is the number of reactive species). For reasons of e ciency, and to fulll the niteness condition of the denition of cellular autom ata, each component  $y_i(r)$  can only take integer values between 0 and  $b_i$ , where the  $b_i$ 's can be di erent for each species i. The position index r is a d-dimensional vector in the CA lattice. For cubic lattices, r is a d-vector of integers.

The central operation of the autom aton consists of calculating the sum

$$y_{i}(r) = \sum_{r^{0} \geq N_{i}}^{X} y_{i}(r + r^{0})$$
 (1)

of the concentrations in some neighborhood N $_{\rm i}$ . The neighborhoods can be di errent for each species i. A neighborhood is specied as a set of displacement vectors, e.g. in two dimensions

$$N_{5 \text{ star}} = \begin{bmatrix} n \\ 0 \\ 0 \end{bmatrix}; \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}; \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}; \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}; \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
(2)

or

N<sub>1</sub> square =

$$N_{5 \text{ star}} \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$
(3)

For di usive system s, sym metry requires that

$$r^{0}2N$$
)  $r^{0}2N$  (4)

and if, as is usually the case in reaction-di usion system s, isotropy is required, then

$$(r_1; d)_r^T 2 N$$
  $(r_1; d)_r^T 2 N$  (5)

for all index permutations ( $_1$ ;  $_d$ ). For some neighborhoods, the sum m ing operation can be executed in an extrem ely e cient way by using m oving averages: in one dim ension, the sum of all 2k + 1 cells centered around cell r can be computed from the corresponding sum centered around cell r 1 with just one addition and subtraction:

$$\begin{split} y_{i}(\mathbf{r}) &= \sum_{j \in j \in \mathbf{k}}^{X} y_{i}(\mathbf{r} + \mathbf{r}^{0}) \\ &= y_{i}(\mathbf{r} - \mathbf{k}) + y_{i}(\mathbf{r} - \mathbf{k} + 1) + \frac{1}{2}(\mathbf{x} + \mathbf{k}) \\ &= y_{i}(\mathbf{r} - \mathbf{k} - 1) + y_{i}(\mathbf{r} - \mathbf{k} - 1) + y_{i}(\mathbf{r} - \mathbf{k}) + \frac{1}{2}(\mathbf{x} + \mathbf{k} - 1) + y_{i}(\mathbf{r} - \mathbf{k}) \\ &= y_{i}(\mathbf{r} - \mathbf{k} - 1) + y_{i}(\mathbf{r} - 1) + y_{i}(\mathbf{r} - \mathbf{k}): \end{split}$$

$$(6)$$

U sing this relationship recursively, and using the fact that the sum over a square neighborhood in d dimensions can be constructed as the convolution of such onedimensional sums applied in each dimension in turn, one obtains an algorithm to compute the sum of all cells in a square (cubic) neighborhood with only 2d additions per cell [8]. In a multispecies model several variables are necessary, but they can be packed into one computer word. In this case the averaging operation can be performed on several species at once if the di usion coe cients are equal.

In the following we use the normalized values  $x_i(r) = y_i(r) = b_i$  and  $\varkappa_i(r) = y_i(r) = (b_i N_i)$ , which are always between zero and one. The resulting elds  $\varkappa_i(r)$  are then the local averages of the  $x_i(r)$ . The averaging has the e ect of di usion. This can be seen from a Taylor expansion of  $x_i(r + r^0)$  around  $x_i(r)$ :

$$\mathbf{x}_{i}(\mathbf{r}) = \frac{1}{\mathcal{N}_{i} j} \frac{X}{r^{0} 2 N} = 0}{\mathcal{N}_{i} (\mathbf{r})} \frac{1}{\mathbf{r}} r^{0} \frac{\theta}{\theta r^{0}} = \mathbf{x}_{i} (\mathbf{r})$$
$$= \mathbf{x}_{i} (\mathbf{r}) + D_{i} r^{2} \mathbf{x}_{i} (\mathbf{r}) +$$

The factors D<sub>i</sub> can be computed as shown in [8] and are easily calculated from (7) for square neighborhoods with radius k: D<sub>i</sub> = k (k + 1)=6.

The second operation in the cellular autom atom is the implementation of the reactive processes described by a rate law.G iven the reaction-di usion equation

$$\frac{\partial z}{\partial t} = (z) + Dr^2 z$$
(8)

(where r  $^2$  is understood to be a spatial operator acting on each component of the vector z separately, and D is a diagonalmatrix), we discretize the time derivative to obtain

$$z^{t+t} = z^{t} + t (z^{t}) + tDr^{2}z^{t}$$
: (9)

C hanging the time and space scales by setting t = t = tand r = r = r, and using the variable x for the rescaled set gives

$$x^{t+1} = x^{t} + t (x^{t}) + \frac{t}{r^{2}} Dr^{2}x^{t}$$
: (10)

as the equation to be treated by the CA. Let us de ne

$$(x) = x + t (x)$$
: (11)

From eqns. (7) and (11)

$$(\mathbf{x}^{t}) = \mathbf{x}^{t} + D_{i}r^{2}\mathbf{x}^{t} + t + t^{t}(\mathbf{x} D_{i}r^{2}\mathbf{x}^{t} + t)$$
$$= \mathbf{x}^{t} + D_{i}r^{2}\mathbf{x}^{t} + t (\mathbf{x}^{t}) + O(t^{2}):$$
(12)

Then

$$\mathbf{x}^{t+1} = (\mathbf{x}^t) \tag{13}$$

is consistently rst order accurate in time and within this limit, eq. (13) can be validly idential with eq. (10) to describe the evolution of the system. The identiation yields

$$D_{i} = \frac{t}{r^{2}} D_{ii}$$
 (i= 1; ;s) (14)

which de nes the space scale. As y is the result of the di usion step, the average output of the CA reactiondi usion process should therefore be given by

$$g_{j} = b_{j} \qquad \frac{\overline{y}_{i}}{b_{i} N_{i} j} \sum_{i=1}^{s}$$
(15)

for species j.<sup>1</sup>

A simplistic discretization of the rate law may produce problem s: due to the discretization spurious steady states or oscillations can appear. It is at this stage that the probabilistic rules come into e ect. G iven an input con guration  $y^t$  (r), one assigns new values  $y^{t+1}$  (r) probabilistically in such a way that the average result corre-(7) sponds to the nite di erence approximation to the given reaction-di usion equation,  $g_i$ . The simplest CA nule for the reactive step is to treat each species separately and use  $y_j^{t+1} = bg_jc + 1$  with probability ( $g_j \mod 1$ ) and  $y_j^{t+1} = bg_jc$  otherw ise (bec is the largest integer smaller

<sup>&</sup>lt;sup>1</sup>Possibly the function needs to be truncated to conform to the condition  $2 [0;1]^{s}$ .

than or equal to e). In this way the average is exactly  $g_j$ . C learly this method introduces the minimal amount of noise for a given set of  $b_i$ 's. In case higher noise levels are desired (e.g., if one wants to evaluate the role of uctuations [12]), one can choose a di erent set of output values and associated probabilities with the same average  $g_j$ , but di erent variance. To do so, in general at least three di erent possible outcom es (e.g.bg<sub>j</sub>c 1;bg<sub>j</sub>c;bg<sub>j</sub>c+1) are necessary.

To demonstrate the applicability of the new CA we show how the G inzburg-Landau equation can be mapped onto the autom aton. We consider the PDE [13,14]

$$\frac{\partial z}{\partial t} = Dr^2 z + az \quad b \dot{z} \dot{f} z; \qquad (16)$$

which can be viewed (for D real) as a two-species reaction-diusion system by separating real and im aginary parts as z = x + iy, a = + i, and b = + i,

$$\frac{e^{2}x}{e^{2}t} = Dr^{2}x + x \qquad y + (x + y)(x^{2} + y^{2}) \quad (17a)$$

$$\frac{e^{2}y}{e^{2}t} = Dr^{2}y + y + x + (y + x)(x^{2} + y^{2}) \quad (17b)$$

where x, y are space- and time-dependent variables. W ith z =  $re^{\rm i}$ , the corresponding rate equation can be conveniently written as

$$\frac{d\mathbf{r}}{d\mathbf{t}} = \mathbf{r} \quad \mathbf{r}^3 = \mathbf{r} = \mathbf{r}^2 \tag{18a}$$

$$\frac{e}{et} = r^2 :$$
 (18b)

O ne can set = 1 by changing the time scale. The stable hom ogeneous solution is  $z = r_s e^{i t} w$  ith  $r_s = t^{-1} = t^{-1}$  and  $r_s = r_s^2$  (A steady but unstable solution is z = 0). Since the equation for r is independent of , one can transform the solution for = 0 to any given by multiplying z(t) with a factor  $e^{i!t}$ , yielding an oscillating solution. The full PDE system, eq. (17), also admits oscillating and rotating spirals, and other inhom ogeneous solutions.

For the CA simulations, the region of interest in concentration space needs to be included in the unit square  $[0;1]^2$  by shifting the origin to (0.5;0.5) and scaling the parameters in such a way that  $0 < R_s < 0.5$  (Here  $R_s = 0.25$  and = 1). In all CA simulations we use a square neighborhood (radius k = 1), i.e.,  $D_1 = D_2 = 1=3$ , and discretization levels  $b_1 = b_2 = 55$  (which gives a lookup-table size of 9 M bytes).

Study of the hom ogeneous solutions shows that the CA behaves like an explicit nite di erence method with added noise. The time step must be su ciently small to reproduce the correct solution. The noise, which is intrinsic in the autom atom and arises from the discretization, has little e ect on the amplitude of the solution, but it introduces random drifts in the phase.



FIG.1. Spiral wave for t = 2, system size 200<sup>2</sup> cells. Superim posed on the grayscale plot of x is the tted A rchimedian spiral.

W e now turn to more interesting cases by considering spiral wave solutions. W ithout loss of generality we use

= 0, which elim inates the hom ogeneous oscillations and thereby allows for larger time steps in the computation and better visualization.

W e initiate a spiral in a two-dimensional simulation by starting with the initial condition

$$R (r_x; r_y; t^0) = c_1 \frac{q}{(r_x r_x^0)^2 + (r_y r_y^0)^2}$$
(19a)

$$(\mathbf{r}_{x};\mathbf{r}_{y};\mathbf{t}^{0}) = c_{2} \arctan \frac{\mathbf{r}_{y}}{\mathbf{r}_{x}} \frac{\mathbf{r}_{y}^{0}}{\mathbf{r}_{x}};$$
 (19b)

which creates exactly one phase singularity at  $(r_x^0; r_y^0)$ . For nonzero smaller than a critical value, a spiral develops and rotates steadily after some time. In Figure 1 we show the spiral obtained for = 1, t = 2, and system size (200 x)<sup>2</sup>. Notice that in spite of the use of a square neighborhood for the CA, the spiral is perfectly round. W hile this feature is expected when the CA is viewed as a num ericalm ethod for solving the PDE, it is not universally accepted that CA dynam ics can produce such an isotropic behavior.

In order to determ ine the e ect of using large time steps, we measure the asymptotic wavelength by tting an Archimedian spiral to the contour x < 0; y = 0. The measured wavelengths for dierent time steps t are shown in Figure 2. The value expected from the theory described in [14] is around  $^0 = 81$ . We observe that the expected value is reached for small enough t. E ven for bigger t the deviations are relatively small and approximately linear in t. When the parameter becomes bigger than some critical value (estimated to be  $_c = 1:397:::in [14]$ ), the spiral wave solution becomes unstable.



FIG.2. Spiral wavelength as a function of the size of the time step t (logarithm ic scale). The data can be reasonably approximated as 80:6 + 1:7 t (solid curve). The size of the system is selected to be  $(6_{p_0})^2$  490<sup>2</sup> in non-dimensional units, corresponding to 490= 3 t cells and the spiral t is performed at t= 4500.

As an example of greater complexity, Figure 3 shows the simulation of a large system ( $500^2$  cells) initialized in the state z 0: Under such conditions m any interacting spirals develop. Indeed, the initial state is unstable, and because of the intrinsic (low level) noise, di erent regions depart from this unstable state with di erent phase values . This situation autom atically creates m any phase singularities (points with r = 0, surrounded by points with all values of ), which then develop into spirals. These phase singularities can merge and move as they are in uenced by each other [15].

Such large simulations are made possible by the speed advantage that the CA o ers over other numerical methods for solving PDEs. We found that on a NeXT workstation the CA runs about 5 times faster than an Euler integration using the same time step. This advantage is due to the use of integer arithmetic, table lookup, and the fact that the di usion calculation is faster in the CA than in nite di erence methods. The speed advantage is much more pronounced when the nonlinear rate functions arem ore complicated, as the table lookup operation does not slow down.

In conclusion, we have constructed a class of cellular autom ata that can be used to model many reactiondi usion system swith quantitatively correct results. The solutions are very isotropic despite the discreteness and the anisotropic nature of the CA. The adverse e ects of discretization are overcome by the use of probabilistic rules. We have demonstrated the applicability of the autom atom for the G inzburg-Landau equation. We have also successfully applied the method to more than ten di erent reaction-di usion models, som e of which will be reported in a forthcoming paper.

W e acknow ledge support from the European C om munity (SC 1-0212), Fonds N ational de la Recherche Scienti que (JPB), G ottlieb D aim ler-und K arlBenz-Stiffung (JRW) and Stiffung Stipendien-Fonds des Verbandes der C hem ischen Industrie (JRW).



FIG.3. Collection of many interacting spirals in a large domain  $(500^2 \text{ cells}, = 1, t = 2, t = 2000)$ 

- Theory and applications of cellular automata; including selected papers '83-'86, Vol. 1 of Advanced Series on Complex Systems, edited by S.W olfram (W orld Scientic, Singapore, 1986).
- [2] R.Kapral, J. of M athem atical Chem istry 6, 113 (1991).
- [3] Lattice gas autom ata: Theory, simulation, im plan entation, edited by J.P.Boon (J.Stat.Phys. 68 3/4, 1992).
- [4] Lattice gas methods for partial di erential equations, edited by G.Doolen (Addison-Wesley, Redwood City, CA, 1990).
- [5] T. To oli, Physica D 10, 117 (1984).
- [6] A. Lawniczak, D. Dab, R. Kapral, and J. P. Boon, Physica D 47, 132 (1991).
- [7] J.M.G meenberg and S.P.Hastings, SIAM J.Appl.M ath. 34, 515 (1978).
- [8] J.R.W eim ar, J.J.Tyson, and L.T.W atson, Physica D 55, 309 (1992).
- [9] J.R.W eim ar, J.J.Tyson, and L.T.W atson, Physica D 55, 328 (1992).
- [10] M . M arkus and B . Hess, N ature 347, 56 (1990).
- [11] H. E. Schepers and M. Markus, Physica A 188, 337 (1992).
- [12] J.R.W eim ar, D.Dab, J.P.Boon, and S.Succi, Europhysics Letters 20, 627 (1992).
- [13] Y. Kuram oto, Chem ical oscillations, waves, and turbulence (Springer-Verlag, Berlin, 1984).
- [14] P.S.Hagan, SIAM J.Appl.M ath. 42, 762 (1982).
- [15] X.-G. Wu, M.-N. Chee, and R. Kapral, Chaos 1, 421 (1991).