

CONNECTION BETWEEN THE SLAVE-PARTICLES AND X-OPERATORS PATH-INTEGRAL REPRESENTATIONS. A NEW PERTURBATIVE APPROACH.

A.Foussats*, A. Greco*, C.Repetto*, O.P.Zandron* and O.S.Zandron*

* Miembros del Consejo Nacional de Investigaciones Científicas y Técnicas - Argentina.

*Facultad de Ciencias Exactas Ingeniería y Agrimensura de la UNR,
Av.Pellegrini 250 - 2000 Rosario - Argentina.*

Abstract

Abstract

In the present work it is shown that the family of first-order Lagrangians for the t-J model and the corresponding correlation generating functional previously found can be exactly mapped into the slave-fermion decoupled representation. Next, by means of the Faddeev-Jackiw symplectic method, a different family of Lagrangians is constructed and it is shown how the corresponding correlation generating functional can be mapped into the slave-boson representation. Finally, in order to define the propagation of fermion modes we discuss two alternative ways to treat the fermionic sector in the path-integral formalism for the t-J model.

PACS: 75.10.Hk and 75.10.Jm

I. INTRODUCTION

Accounting the relevance in describing the behavior of strongly correlated electron systems, a renovate interest in the study of the supersymmetric generalisations of the Hubbard model appears in the last years. A complete review on strongly correlated electron systems together with its connection to high- T_c is given in Ref.[1].

The Hubbard models based on the superalgebras $\text{spl}(2,1)$, $\text{osp}(2,2)$ or $\text{su}(2,2)$ have been formulated by using several approaches²⁻⁶. For instance, as it was suggested⁷⁻⁹ the superalgebra $\text{spl}(2,1)$ could be useful to study the model in the limit of infinite on-site repulsion and with infinite-range hopping between all sites¹⁰.

Many problems about correlated electron systems were treated in the framework of the decoupled slave-particle representations. Two of them are the most important: the slave-boson and the slave-fermion representations. The first one privileges the fermion dynamics, and therefore the slave-boson representation seems to be more adequated to describe a Fermi liquid state^{11,12}. Instead, the slave-fermion representation seems to give a good response when the system is closed to the antiferromagnetic order^{13,14}.

An important question in order to understand the physics of the high- T_c superconductors, is to solve how to go from one representation to the other. This is because in high- T_c superconductors, both the Fermi liquid and the magnetic order states seem to be present.

On the other hand, one of the main problems appearing in these kind of models is to define the dynamics of fermions in the constrained Hilbert space, when the double occupancy of lattice sites is excluded. In this case also a convenient representation is given in terms of slave-particles¹⁵.

As it is known the slave-particle models exhibit a local gauge invariance which is destroyed in the mean field approximation. This local gauge invariance has associated a first-class constraint which is difficult to handle in the path-integral formalism.

Another possibility to attack the problem was given in Refs.[16,17] by using generalised coherent states in the framework of the functional integral formalism.

Recently, the t-J model was analysed in the context of the path-integral formalism^{18,19}.

Our starting point was the construction of a particular family of first-order constrained Lagrangians by using the Faddeev-Jackiw (FJ) symplectic method²⁰, in the supersymmetric version^{21,22}. In this approach any decoupling is used, but the field variables are directly the Hubbard X -operators which verify the superalgebra $\text{spl}(2,1)$. In this way, we always work with the real physical excitations.

Next, by using path-integral techniques, the correlation generating functional and the effective Lagrangian were constructed. Moreover, we have proved that our path-integral representation can be directly related with that found in Ref.[23].

As mentioned above, one of the interesting and non completely solved problem present in this constrained system, is to study the fermionic sector when the double occupancy of lattice sites is excluded. In particular the role of the fermionic constraints, and so the fermionic dynamics in the constrained Hilbert space is a crucial problem that must be investigated.

Therefore, one of the main purposes of the present paper is to go deep into the discussion of the different alternatives which allows us to define the fermionic propagator in the t-J model.

In Ref.[19], in the framework of the perturbative formalism the Feynman rules with appropriate propagators and vertices were found. In particular, a discussion on the fermionic propagator was also given.

In order to continue with the study of the fermionic propagation, different alternatives are exposed in the present paper. We show that there is another way that the one given in Ref.[19] to obtain fermionic propagation. This is done by working inside the path-integral and integrating out the two delta functions on the fermionic constraints.

Moreover, other interesting point is to check our formalism with those obtained by means of the slave-particle representations. More precisely, we will show how our path-integral expression for the partition function (see equation (4.1) of Ref.[18]), written in terms of the Hubbard operators, can be mapped in the partition function coming from the slave-fermion representation.

On the other hand, by following the FJ symplectic method, it is possible to show that a new family of first-order constrained Lagrangians written in terms of the Hubbard X -

operators exists. This family of classical Lagrangians is able to reproduce the Hubbard X -operators commutation rules verifying the graded algebra $\text{spl}(2,1)$. As it can be shown this family of Lagrangians, totally constrained in the boson-like Hubbard X -operators, can be mapped into the slave-boson representation.

The paper is organized as follow. In section II, the main results of Refs.[18,19] are collected. In section III, by analysing the change in the constraints structure of the t-J model, it is showed how starting from our path-integral expression for the partition function previously found, the partition function coming from the decoupled slave-fermion representation can be recovered. In section IV, by using the FJ symplectic method, a different family of first-order constrained Lagrangians is found. This family of Lagrangians corresponds to the situation in which the bosons are totally constrained. In such conditions it is possible to show how the correspondent partition function can be mapped in the partition function coming from the slave-boson representation. In section V, two alternative ways to define the fermion propagation are studied. In section VI, conclusions are given.

II. PRELIMINARY AND DEFINITIONS

In the t-J model the three possible states on a lattice site are $|\alpha\rangle = |0\rangle, |+\rangle, |-\rangle$. These states correspond respectively to an empty site, an occupied site with an electron of spin-up, or an occupied site with an electron of spin-down. Double occupancy is forbidden in the t-J model. In terms of these states the Hubbard \hat{X} -operators are defined as

$$\hat{X}_i^{\alpha\beta} = |i\alpha\rangle\langle i\beta|. \quad (2.1)$$

In Eq.(2.1), when one of the index is zero and the other different from zero, the corresponding \hat{X} -operator is fermion-like, otherwise boson-like.

The Hubbard \hat{X} -operators verify the following graded commutation relations

$$[\hat{X}_i^{\alpha\beta}, \hat{X}_j^{\gamma\delta}]_{\pm} = \delta_{ij}(\delta^{\beta\gamma}\hat{X}_i^{\alpha\delta} \pm \delta^{\alpha\delta}\hat{X}_i^{\gamma\beta}), \quad (2.2)$$

where the $+$ sign must be used when both operators are fermion-like, otherwise it corresponds the $-$ sign, and i, j denotes the site indices.

We assume that the family of classical constrained first-order Lagrangians in terms of the Hubbard \hat{X} -operators can be written as follows

$$L = a_{\alpha\beta}(X)\dot{X}^{\alpha\beta} - \mathbf{V}^{(0)} . \quad (2.3)$$

In the FJ language²⁰ the symplectic potential $\mathbf{V}^{(0)}$ is defined by

$$\mathbf{V}^{(0)} = H(X) + \lambda^a \Omega_a , \quad (2.4)$$

where λ^a are appropriate Lagrange multipliers, and so the constraints Ω_a are defined by

$$\Omega_a = \frac{\partial \mathbf{V}^{(0)}}{\partial \lambda^a} . \quad (2.5)$$

Therefore, the symplectic supermatrix associated to the Lagrangian (2.3) can be formally written²²

$$M_{AB} = \begin{pmatrix} \frac{\partial a_{\gamma\delta}}{\partial X^{\alpha\beta}} - (-1)^{|\alpha\beta||\gamma\delta|} \frac{\partial a_{\alpha\beta}}{\partial X^{\gamma\delta}} & \frac{\partial \Omega_b}{\partial X^{\alpha\beta}} \\ -(-1)^{|a||\gamma\delta|} \frac{\partial \Omega_a}{\partial X^{\gamma\delta}} & 0 \end{pmatrix} , \quad (2.6)$$

where the compound indices $A = \{(\alpha\beta), a\}$ and $B = \{(\gamma\delta), b\}$ run in the different ranges of the complete set of variables defining the extended configuration space, and $|A|$ indicates the Fermi grading.

Following Ref.[18], our starting point is to consider the following partition function for the t-J model written in terms of the four boson-like operators (X^{+-} , X^{-+} , X^{++} , X^{--}) and the four fermion-like operators (X^{0+} , X^{0-} , X^{+0} , X^{-0})

$$Z = \int \mathcal{D}X_i^{\alpha\beta} \delta(\Omega_{i1}) \delta(\Omega_{i2}) \delta(\Xi_{i3}) \delta(\Xi_{i4}) (sdet M_{AB})_i^{\frac{1}{2}} \exp i \int dt L(X, \dot{X}) , \quad (2.7)$$

where $L(X, \dot{X})$ is given by

$$\begin{aligned} L(X, \dot{X}) = & i \sum_i \frac{(1 + \rho_i)u_i - 1}{(2 - v_i)^2 - 4\rho_i - u_i^2} (X_i^{-+} \dot{X}_i^{+-} - X_i^{+-} \dot{X}_i^{-+}) \\ & + \frac{i}{2} \sum_{i,\sigma} (\dot{X}_i^{0\sigma} X_i^{\sigma 0} + \dot{X}_i^{\sigma 0} X_i^{0\sigma}) - \mu \sum_{i,\sigma} X_i^{0\sigma} X_i^{\sigma 0} - H_{t-J}(X) . \end{aligned} \quad (2.8)$$

where $u_i = X_i^{++} - X_i^{--}$ and $v_i = X_i^{++} + X_i^{--}$.

The Greek indices α, β takes the values $\{+, -, 0\}$, the index σ takes the values $\{+, -\}$, and $H_{t-J}(X)$ is the usual t-J Hamiltonian

$$H_{t-J} = \sum_{i,j,\sigma} t_{ij} X_i^{\sigma 0} X_j^{0\sigma} + \frac{1}{4} \sum_{i,j,\sigma,\bar{\sigma}} J_{ij} X_i^{\sigma\bar{\sigma}} X_j^{\bar{\sigma}\sigma} - \frac{1}{4} \sum_{i,j,\sigma,\bar{\sigma}} J_{ij} X_i^{\sigma\sigma} X_j^{\bar{\sigma}\bar{\sigma}} , \quad (2.9)$$

besides, in equation (2.8) a term depending on the chemical potential μ was added.

In Eq.(2.7) $s\det M_{AB}$ is the superdeterminant of the symplectic supermatrix M_{AB} defined in (2.6), and the bosonic and fermionic constraints at each site i are given respectively by

$$\Omega_{i1} = X_i^{++} + X_i^{--} + \rho_i - 1 = 0 , \quad (2.10a)$$

$$\Omega_{i2} = X_i^{+-} X_i^{-+} + \frac{1}{4}(X_i^{++} - X_i^{--})^2 - [1 - \frac{1}{2}(X_i^{++} + X_i^{--})]^2 + \rho_i = 0 , \quad (2.10b)$$

$$\Xi_{i3} = X_i^{0+} X_i^{+-} - X_i^{0-} X_i^{++} = 0 , \quad (2.10c)$$

$$\Xi_{i4} = X_i^{+0} X_i^{-+} - X_i^{-0} X_i^{++} = 0 . \quad (2.10d)$$

where was defined $\rho_i = X_i^{0+} X_i^{+0} + X_i^{0-} X_i^{-0}$.

In equation (2.8), the Lagrangian coefficients as well as the constraints (2.10) were determined by using the FJ symplectic method with the condition to reproduce at the classical level the generalised FJ brackets or graded Dirac brackets of the t-J model (see Ref.[18]).

In particular the constraint (2.10a) deduced by consistency, is the completeness condition which must be verified by the Hubbard \hat{X} -operators and plays an important physical role in this constrained model as it will be seen later on. At this stage it is important to remark that in the Dirac language²⁴ the two bosonic constraints (2.10a,b) are second-class one.

Now, it is useful to write the boson-like Hubbard X -operators in terms of the real components S_α ($\alpha = 1, 2, 3$) of a vector field \mathbf{S} and the fermion-like Hubbard X -operators in terms of suitable component spinors (Grassmann variables)^{18,23}

$$X_i^{++} = \frac{1}{2s}(1 - \rho_i)(s + S_{i3}) , \quad (2.11a)$$

$$X_i^{--} = \frac{1}{2s}(1 - \rho_i)(s - S_{i3}) , \quad (2.11b)$$

$$X_i^{+-} = \frac{1}{2s}(1 - \rho_i)(S_{i1} + iS_{i2}) , \quad (2.11c)$$

$$X_i^{-+} = \frac{1}{2s}(1 - \rho_i)(S_{i1} - iS_{i2}) , \quad (2.11d)$$

$$X_i^{-0} = \Psi_{i+} , \quad X_i^{0-} = \Psi_{i+}^* , \quad (2.11e)$$

$$X_i^{+0} = \Psi_{i-} , \quad X_i^{0+} = \Psi_{i-}^* , \quad (2.11f)$$

where s is a constant and the hole density in the new variables writes $\rho_i = \Psi_{i+}^* \Psi_{i+} + \Psi_{i-}^* \Psi_{i-}$. Accounting the fermionic constraints (2.10) it results $(1 - \rho_i)(1 + \rho_i) = 1$.

The real vector field \mathbf{S} can be identified with the spin only when $\rho = 0$, i.e., in the pure bosonic case.

By using the second-class constraints (2.10a) and after the change of variables is made, the partition function takes the form

$$Z = \int \mathcal{D}S_{i1} \mathcal{D}S_{i2} \mathcal{D}S_{i3} \mathcal{D}\Psi_{i\sigma} \mathcal{D}\Psi_{i\sigma}^* \mathcal{D}\lambda_i \mathcal{D}\xi_i \mathcal{D}\xi_i^* (s \det M_{AB})_i^{\frac{1}{2}} \left(\frac{\partial X}{\partial S} \right)_i \exp \left(i \int dt L_{eff} \right) . \quad (2.12)$$

where the quantity $\left(\frac{\partial X}{\partial S} \right)_i$ is the super Jacobian of the transformation (2.11).

The effective Lagrangian L_{eff} defined in Eq. (2.12), in terms of the new variables reads

$$\begin{aligned} L_{eff} = & \frac{1}{2s} \sum_i \frac{S_{i1} \dot{S}_{i2} - S_{i2} \dot{S}_{i1}}{s + S_{i3}} + i \sum_{i,\sigma} \dot{\Psi}_{i\sigma}^* \Psi_{i\sigma} + \mu \sum_{i,\sigma} \Psi_{i\sigma} \Psi_{i\sigma}^* - H_{t-J} \\ & + \sum_i \left[\lambda_i (S_{i1}^2 + S_{i2}^2 + S_{i3}^2 - s^2) + \xi_i^* (\Psi_{i-} (S_{i1} - iS_{i2}) - \Psi_{i+} (s + S_{i3})) \right. \\ & \left. + (\Psi_{i-}^* (S_{i1} + iS_{i2}) - \Psi_{i+}^* (s + S_{i3})) \xi_i \right] , \end{aligned} \quad (2.13)$$

where the Hamiltonian H_{t-J} of the t-J model writes

$$H_{t-J} = \sum_{i,j,\sigma} t_{ij} \Psi_{i\sigma} \Psi_{j\sigma}^* + \frac{1}{8s^2} \sum_{i,j} J_{ij} (1 - \rho_i)(1 - \rho_j) [S_{i1} S_{j1} + S_{i2} S_{j2} + S_{i3} S_{j3} - s^2] . \quad (2.14)$$

In equation (2.13) the parameters λ_i , ξ_i^* and ξ_i are respectively suitable bosonic and fermionic Lagrange multipliers.

At this stage it is important to remark that our Lagrangian formalism is independent of the underlying lattice dimension.

In the next section we are going to analyse the equation (2.7) in the framework of the decoupled slave-particle representations.

III. SLAVE-PARTICLE REPRESENTATIONS.

The standard way to construct the classical Hamiltonian formulation for slave-particle models and subsequently give the canonical quantisation is developed in Ref.[15] . The starting point is to consider the general classical first-order Lagrangian for n bosonic fields b_a and m fermionic fields f_b defined on a lattice

$$L(b_a^\dagger, b_a, f_b^\dagger, f_b) = \frac{i}{2} \sum_{i,a} (b_{ia}^\dagger \dot{b}_{ia} - \dot{b}_{ia}^\dagger b_{ia}) + \frac{i}{2} \sum_{i,b} (f_{ib}^\dagger \dot{f}_{ib} - \dot{f}_{ib}^\dagger f_{ib}) - H(b_a^\dagger, b_a, f_b^\dagger, f_b) . \quad (3.1)$$

Both, bosons and fermions fields, are submitted to the slave-particle first-class constraint at each lattice site i

$$\Omega_i = \sum_a b_{ia}^\dagger b_{ia} + \sum_b f_{ib}^\dagger f_{ib} - 1 = 0 . \quad (3.2)$$

Looking at the equations (3.1) and (3.2) it can be seen that when the index a takes the values \pm and the index b takes only one value, the six fields (four boson and two fermion fields) define the slave-fermion representation. On the contrary, when the index a takes only one value and the index b takes the values \pm , the six fields (two boson and four fermion fields) define the slave-boson representation.

With the aim to confront our results with others previously obtained, in this section we consider the slave-fermion representation. In particular it is possible to confront the correlation generating functional (2.7) with that obtained from the slave-fermion representation, and this relation is not trivial.

In our approach all the constraints are in the Dirac picture²⁴ second-class, while in the slave-particle representations the constraint (3.2) is first-class. Thus, when the Hubbard X -operators are decoupled a local gauge symmetry is made evident.

The starting point is the correlation generating functional (2.7) with the Lagrangian (2.8).

By computing the $(s\det M_{AB})_i^{\frac{1}{2}}$ appearing in equation (2.7) we find

$$(s\det M_{AB})_i^{\frac{1}{2}} = -i \frac{(1 + \rho_i)}{X_i^{++}} . \quad (3.3)$$

where ρ_i evaluated on the constraints is written $\rho_i = \frac{X_i^{0+} X_i^{+0}}{X_i^{++}}$.

Integrating out the fields components X_i^{--} , X_i^{0-} and X_i^{-0} by using the delta functions on the constraints written as follows

$$\delta(\Omega_{i2}) = \delta(X_i^{++}X_i^{--} - X_i^{+-}X_i^{-+}) = \frac{1}{X_i^{++}}\delta(X_i^{--} - \frac{X_i^{+-}X_i^{-+}}{X_i^{++}}), \quad (3.4a)$$

$$\delta(\Xi_{i3}) = \delta(X_i^{0+}X_i^{+-} - X_i^{0-}X_i^{++}) = X_i^{++}\delta(X_i^{0-} - \frac{X_i^{0+}X_i^{+-}}{X_i^{++}}), \quad (3.4b)$$

$$\delta(\Xi_{i4}) = \delta(X_i^{+0}X_i^{-+} - X_i^{-0}X_i^{++}) = X_i^{++}\delta(X_i^{-0} - \frac{X_i^{+0}X_i^{-+}}{X_i^{++}}), \quad (3.4c)$$

and taking into account the equality $(1+\rho_i)\delta(X_i^{-0} - \frac{X_i^{+0}X_i^{-+}}{X_i^{++}}) = \delta[(1-\rho_i)(X_i^{-0} - \frac{X_i^{+0}X_i^{-+}}{X_i^{++}})] = \delta(X_i^{-0} - \frac{X_i^{+0}X_i^{-+}}{X_i^{++}})$ (coming from the property of the Grassmann variables), the partition function (2.7) takes the form

$$Z = \int \mathcal{D}X_i^{++} \mathcal{D}X_i^{+-} \mathcal{D}X_i^{-+} \mathcal{D}X_i^{+0} \mathcal{D}X_i^{0+} \delta(X_i^{++} + \frac{X_i^{+-}X_i^{-+}}{X_i^{++}} + \frac{X_i^{0+}X_i^{+0}}{X_i^{++}} - 1) \\ \times \exp(i \int dt L^*(X, \dot{X})), \quad (3.5)$$

where $L^*(X, \dot{X})$ is given by

$$L^*(X, \dot{X}) = \frac{i}{2} \sum_i \frac{1}{X_i^{++}} (X_i^{-+} \dot{X}_i^{+-} - X_i^{+-} \dot{X}_i^{-+}) \\ + \frac{i}{2} \sum_{i,\sigma} \frac{1}{X_i^{++}} (X_i^{+0} \dot{X}_i^{0+} + X_i^{0+} \dot{X}_i^{+0}) - H(X). \quad (3.6)$$

As it is known the change of variables that allows to write the remaining five Hubbard X variables in terms of the fields variables in the decoupled slave-fermion representation is defined by

$$X_i^{++} = b_{i+}^\dagger b_{i+}, \quad (3.7a)$$

$$X_i^{+-} = b_{i+}^\dagger b_{i-}, \quad (3.7b)$$

$$X_i^{-+} = b_{i-}^\dagger b_{i+}, \quad (3.7c)$$

$$X_i^{0+} = b_{i+} f_i^\dagger, \quad (3.7d)$$

$$X_i^{+0} = b_{i+}^\dagger f_i . \quad (3.7e)$$

From equations (3.7) it can be seen that the five Hubbard X -fields are given in terms of the six fields of the slave-fermion representation, so it is necessary to introduce an additional condition among the six fields of the slave-fermion representation to make possible the transformation.

We assume the following general linear form for the conditions in each lattice site

$$\phi_i = \sum_a (G_{ia} b_{ia} + G_{ia}^\dagger b_{ia}^\dagger) + H_i f_i - H_i^\dagger f_i^\dagger + K_i = 0, \quad (3.8)$$

where $G_{ia}, G_{ia}^\dagger, K_i$ are bosonic parameters and H_i, H_i^\dagger are fermionic (Grassmannian) parameters.

As it was commented above, when the Hubbard X -operators are written in a decoupled representation a local gauge symmetry is made evident. Thus, from a constrained system with a set of second-class constraints, it changes into a constrained system with a first-class constraint, and therefore a gauge fixing condition must be imposed. Therefore, the equation (3.8) is not other than the gauge fixing condition which corresponds to the local gauge symmetry appearing in the decoupled representation¹⁵.

A convenient choice is to take in equation (3.8): $G_{i+} = i, G_{i+}^\dagger = -i$ and the remaining coefficients all zero, i.e equation (3.8) reads

$$\phi_i = i(b_{i+} - b_{i+}^\dagger) = 0 . \quad (3.9)$$

Later on, in equation (3.5) is introduced as usual the unity

$$\begin{aligned} 1 = & \int \mathcal{D}b_{i\sigma}^\dagger \mathcal{D}b_{i\sigma} \mathcal{D}f_i^\dagger \mathcal{D}f_i \delta(X_i^{++} - b_{i+}^\dagger b_{i+}) \delta(X_i^{+-} - b_{i+}^\dagger b_{i-}) \delta(X_i^{-+} - b_{i-}^\dagger b_{i+}) \\ & \times \delta(X_i^{0+} - f_i^\dagger b_{i+}) \delta(X_i^{+0} - b_{i+}^\dagger f_i) \delta(\phi_i) J_i , \end{aligned} \quad (3.10)$$

where J_i is the super Jacobian of the transformation (3.7) and (3.9), and its value is $J_i = (b_{i+} + b_{i+}^\dagger)$.

By integrating out the five variables X_i , the partition function can be written

$$Z = \int \mathcal{D}b_{i\sigma}^\dagger \mathcal{D}b_{i\sigma} \mathcal{D}f_i \mathcal{D}f_i^\dagger \delta(\Omega_i) \delta(\phi_i) J_i \exp(i \int dt L(b_\sigma, b_\sigma^\dagger, f, f^\dagger)). \quad (3.11)$$

It is easy to see that the super Jacobian J_i is equal to minus the determinant of the Dirac bracket constructed from the first-class constraint Ω_i and the gauge fixing condition (3.9), i.e $-det[\Omega_i, \phi_i]_D$, where the first-class constraint Ω_i in the slave-fermion representation is given by

$$\Omega_i = \sum_{i\sigma} b_{i\sigma}^\dagger b_{i\sigma} + f_i^\dagger f_i - 1 = 0. \quad (3.12)$$

In the equation (3.11) the Lagrangian $L(b_\sigma, b_\sigma^\dagger, f, f^\dagger)$ reads

$$L(b_\sigma, b_\sigma^\dagger, f, f^\dagger) = \frac{i}{2} \sum_{i,\sigma} (b_{i\sigma}^\dagger \dot{b}_{i\sigma} - \dot{b}_{i\sigma}^\dagger b_{i\sigma}) + \frac{i}{2} \sum_i (f_i^\dagger \dot{f}_i - \dot{f}_i^\dagger f_i) - H(b_{i\sigma}^\dagger, b_{i\sigma}, f_i^\dagger, f_i). \quad (3.13)$$

Therefore, the above considerations show that our correlation generating functional (2.7) can be mapped into the correlation generating functional (3.11) coming from the slave-fermion representation¹⁵. This mapping is a consequence of the fermionic constraints present in our expression for the correlation generating functional (2.7). We can also conclude that it is not possible to relate our correlation generating functional (2.7) with that corresponding to the slave-boson representation.

The next question is how to construct from the symplectic FJ formalism a new family of first-order Lagrangian by using the Hubbard X -operators of the graded algebra $\text{spl}(2,1)$ as fields variables, in such a way that the results can be mapped in the slave-boson representation. The problem is solved in the next section

IV. CLASSICAL LAGRANGIAN AND CONSTRAINTS. SLAVE-BOSON REPRESENTATION.

By following Refs.[18,25], we assume that the family of classical first-order Lagrangians in terms of the Hubbard \hat{X} -operators can be written as follows

$$L = a_{\alpha\beta}(X) \dot{X}^{\alpha\beta} - \mathbf{V}^{(0)}. \quad (4.1)$$

where the five Hubbard \hat{X} -operators $X^{\sigma\sigma'}$ and X^{00} are boson-like and the four Hubbard \hat{X} -operators $X^{\sigma 0}$ and $X^{0\sigma}$ are fermion-like. In the present case the symplectic potential is $\mathbf{V}^{(0)} = H(X)$.

The Lagrangian functional coefficients $a_{\alpha\beta}(X)$ that *a priori* are unknown must be determined by consistency in such a way that the graded algebra (2.2) for the Hubbard \hat{X} -operators is verified. By following the steps of Ref.[18] it is straightforward to construct the symplectic supermatrix associated to the Lagrangian (4.1). Thus the symplectic supermatrix M_{AB} is written in the form

$$M_{AB} = \begin{pmatrix} A_{bb} & B_{bf} \\ C_{fb} & D_{ff} \end{pmatrix}. \quad (4.2)$$

The Bose-Bose parts A_{bb} is a (10×10) -dimensional matrix and it takes the form

$$A_{bb} = \begin{pmatrix} \frac{\partial a_{\sigma\sigma'}}{\partial X^{\sigma''\sigma'''}} - \frac{\partial a_{\sigma''\sigma'''}}{\partial X^{\sigma\sigma'}} & \frac{\partial a_{00}}{\partial X^{\sigma''\sigma'''}} - \frac{\partial a_{\sigma''\sigma'''}}{\partial X^{00}} & \frac{\partial \Omega_{\sigma\sigma'}}{\partial X^{\sigma''\sigma'''}} & \frac{\partial \Omega_{00}}{\partial X^{\sigma''\sigma'''}} \\ -\frac{\partial a_{00}}{\partial X^{\sigma\sigma'}} + \frac{\partial a_{\sigma\sigma'}}{\partial X^{00}} & 0 & \frac{\partial \Omega_{\sigma\sigma'}}{\partial X^{00}} & \frac{\partial \Omega_{00}}{\partial X^{00}} \\ -\frac{\partial \Omega_{\sigma''\sigma'''}}{\partial X^{\sigma\sigma'}} & -\frac{\partial \Omega_{\sigma''\sigma'''}}{\partial X^{00}} & 0 & 0 \\ -\frac{\partial \Omega_{00}}{\partial X^{\sigma\sigma'}} & -\frac{\partial \Omega_{00}}{\partial X^{00}} & 0 & 0 \end{pmatrix}, \quad (4.3)$$

The Bose-Fermi parts B_{bf} (the Fermi-Bose parts $C_{fb} = -B_{bf}^T$) is a (4×10) -dimensional rectangular supermatrix given by

$$B_{bf} = \begin{pmatrix} \frac{\partial a_{0\sigma}}{\partial X^{\sigma''\sigma'''}} - \frac{\partial a_{\sigma''\sigma'''}}{\partial X^{0\sigma}} & \frac{\partial a_{\sigma 0}}{\partial X^{\sigma''\sigma'''}} - \frac{\partial a_{\sigma''\sigma'''}}{\partial X^{\sigma 0}} \\ \frac{\partial a_{0\sigma}}{\partial X^{00}} - \frac{\partial a_{00}}{\partial X^{0\sigma}} & \frac{\partial a_{\sigma 0}}{\partial X^{00}} - \frac{\partial a_{00}}{\partial X^{\sigma 0}} \\ -\frac{\partial \Omega_{\sigma''\sigma'''}}{\partial X^{0\sigma}} & -\frac{\partial \Omega_{\sigma''\sigma'''}}{\partial X^{\sigma 0}} \\ -\frac{\partial \Omega_{00}}{\partial X^{0\sigma}} & -\frac{\partial \Omega_{00}}{\partial X^{\sigma 0}} \end{pmatrix}. \quad (4.4)$$

The Fermi-Fermi parts D_{ff} is the (4×4) -dimensional matrix given by

$$D_{ff} = \begin{pmatrix} \frac{\partial a_{0\sigma}}{\partial X^{0\sigma'}} + \frac{\partial a_{0\sigma'}}{\partial X^{0\sigma}} & \frac{\partial a_{\sigma 0}}{\partial X^{0\sigma'}} + \frac{\partial a_{0\sigma'}}{\partial X^{\sigma 0}} \\ \frac{\partial a_{0\sigma}}{\partial X^{\sigma 0}} + \frac{\partial a_{\sigma' 0}}{\partial X^{0\sigma}} & \frac{\partial a_{\sigma 0}}{\partial X^{\sigma 0}} + \frac{\partial a_{\sigma' 0}}{\partial X^{\sigma 0}} \end{pmatrix}, \quad (4.5)$$

where $\Omega_{\sigma\sigma'}$ and Ω_{00} are the appropriate bosonic second-class constraints defining the structure of the constrained model.

Once the symplectic algorithm is applied and the correspondent differential equations are solved the solution we found is

$$a_{i0\sigma} = \frac{i}{2X_i^{00}} X_i^{\sigma 0}, \quad a_{i\sigma 0} = \frac{i}{2X_i^{00}} X_i^{0\sigma} \quad (4.6)$$

and the boson-like Lagrangian coefficients are all zero.

The set of bosonic second class constraints is given by

$$\Omega_i^{00} = X_i^{00} + X_i^{++} + X_i^{--} - 1 = 0 , \quad (4.7a)$$

$$\Omega_i^{\sigma\sigma'} = X_i^{\sigma\sigma'} - \frac{X_i^{\sigma 0} X_i^{0\sigma'}}{X_i^{00}} = 0 . \quad (4.7b)$$

In particular the constraint (4.7a) is the completeness condition necessary to avoid the double occupancy at each site.

In these conditions the symplectic supermatrix is invertible and the matrix elements of its inverse gives the correct Hubbard graded brackets (2.2), i.e

$$(M^{AB})^{-1} = -i(-1)^{|\varepsilon_A|} [\hat{A} , \hat{B}]_{\pm} , \quad (4.8)$$

where $|\varepsilon_A|$ is the Fermi grading of the field variable A .

Consequently the dynamics in this condition is given by the Lagrangian

$$L(X, \dot{X}) = -\frac{i}{2X_i^{00}} \sum_{i,\sigma} (\dot{X}_i^{0\sigma} X_i^{\sigma 0} + \dot{X}_i^{\sigma 0} X_i^{0\sigma}) - H(X) . \quad (4.9)$$

The Lagrangian (4.9) together with the bosonic constraints (4.7) correspond to a situation in which the bosons are totally constrained and the dynamics is carried out only by the fermions.

The partition function corresponding to this solution reads

$$\begin{aligned} Z = & \int \mathcal{D}X_i^{\alpha\beta} \delta[X_i^{00} + X_i^{++} + X_i^{--} - 1] \delta[X_i^{\sigma\sigma'} - \frac{X_i^{\sigma 0} X_i^{0\sigma'}}{X_i^{00}}] (sdet M_{AB})_i^{\frac{1}{2}} \\ & \times \exp(i \int dt L(X, \dot{X})) . \end{aligned} \quad (4.10)$$

By computing the superdeterminant of the symplectic matrix appearing in (4.10) it results

$$(sdet M_{AB})_i^{\frac{1}{2}} = \left(\det A \left[\det(D - CA^{-1}B) \right]^{-1} \right)^{\frac{1}{2}} = (X_i^{00})^2 . \quad (4.11)$$

Now, in order to confront the correlation generating functional (4.10) with those coming from the slave-boson representation some algebraic manipulations are needed.

The first step is to make the following change of variables

$$\varphi_{i1} = X_i^{00} - b_i^\dagger b_i = 0 , \quad (4.12a)$$

$$\varphi_{i\sigma 0} = X_i^{\sigma 0} - f_{i\sigma}^\dagger b_i = 0 , \quad (4.12b)$$

$$\varphi_{i0\sigma} = X_i^{0\sigma} - f_{i\sigma} b_i^\dagger = 0 . \quad (4.12c)$$

Analogously to what happens in the slave-fermion representation, in the decoupled slave-boson one also an additional condition among the fields is needed. From the general linear equation (3.8) we choose by simplicity the following reality condition

$$\varphi_{i2} = b_i^\dagger - b_i = 0 , \quad (4.12d)$$

The super Jacobian J_i of the transformation (4.12) is given by

$$J_i = -\frac{(b_i^\dagger + b_i)}{(b_i^\dagger b_i)^2} . \quad (4.13)$$

Now, by introducing in the equation (4.10) the unity

$$1 = \int \mathcal{D}f_{i\sigma}^\dagger \mathcal{D}f_{i\sigma} \mathcal{D}b_i^\dagger \mathcal{D}b_i J_i \delta(X_i^{\sigma 0} - f_{i\sigma}^\dagger b_i) \delta(X_i^{0\sigma} - f_{i\sigma} b_i^\dagger) \delta(b_i^\dagger b_i - X_i^{00}) \delta(b_i^\dagger - b_i) , \quad (4.14)$$

and integrating out all the fields variables $X_i^{\alpha\beta}$, after same algebraic manipulations it is possible to show that the partition function (4.10) takes the form

$$Z = \int \mathcal{D}b_i^\dagger \mathcal{D}b_i \mathcal{D}f_{i\sigma}^\dagger \mathcal{D}f_{i\sigma} \delta(\Omega_i) \delta(\phi_i) (b_i + b_i^\dagger) \exp \left(i \int dt L(b^\dagger, b, f_\sigma^\dagger, f_\sigma) \right) , \quad (4.15)$$

where Ω_i and ϕ_i are respectively the first-class constraint and the gauge fixing condition in the radial gauge¹², appearing in the partition function of the slave-boson representation¹⁷. They respectively read

$$\Omega_i = b_i^\dagger b_i + \sum_{i,\sigma} f_{i\sigma}^\dagger f_{i\sigma} - 1 = 0 . \quad (4.16)$$

$$\phi_i = i(b_i - b_i^\dagger) = 0 . \quad (4.17)$$

Again, we note that the factor $(b + b^\dagger)$ in the equation (4.15) is precisely the value of the $\det[\Omega_i, \phi_i]_D$ appearing in the gauge theories containing first-class constraints.

The Lagrangian $L(b^\dagger, b, f_\sigma^\dagger, f_\sigma)$ defined in the equation (4.15) is given by

$$L(b^\dagger, b, f_\sigma^\dagger, f_\sigma) = \frac{i}{2} \sum_i (b_i^\dagger \dot{b}_i - \dot{b}_i^\dagger b_i) + \frac{i}{2} \sum_{i,\sigma} (f_{i\sigma}^\dagger \dot{f}_{i\sigma} - \dot{f}_{i\sigma}^\dagger f_{i\sigma}) - H. \quad (4.18)$$

In summary, from our approach and working with the Hubbard X -operators without using any decoupling representation, the new family of Lagrangians (4.9) is obtained. The respective correlation generating functional (4.10) is mapped into the solution provided by the slave-boson representation. It is important to note that the new path-integral (4.10) in terms of the Hubbard X -operators up to our knowledge was developed in the present paper for the first time.

We can see once more how a second-class constrained model written in terms of the Hubbard X -operators, when written in terms of the decoupled slave-particle representations, is transformed in a constrained system where a local gauge symmetry is made evident.

We think that the results we can obtain by using the partition function (4.10) with the Lagrangian (4.9) can be useful for regimes where the system is close to a Fermi liquid state.

In a forthcoming paper the partition function (4.10) is studied in detail within the context of the perturbative formalism. Having in mind the difficulty to treat the first-class constraint inside the path-integral (4.15), our purpose is to construct the Feynman rules and the diagrammatics starting from the path-integral (4.10). Once an appropriate fermion propagator can be found, our first objective will be to analyse the properties of the fermion spectral function.

V. TWO ALTERNATIVE WAYS TO DEFINE THE PROPAGATOR OF FERMION MODES.

In this section we discuss two alternative ways to treat the fermionic sector in order to define the propagation of the fermion modes. As commented above, a crucial problem in the t-J model is to define the fermionic propagation in the constrained Hilbert space, when the double occupancy is forbidden.

With the purpose to study this problem, in Ref.[19] the correlation generating functional (2.12) was considered at finite temperature by means of the "Euclideanization procedure". Moreover it was assumed that we are close to an undoped regime where the system is an

antiferromagnetic insulator. Under this condition there is a small number of holes and it can be assumed that the hole density $\rho_i = \langle \rho_i \rangle = \text{constant}$. The constant value ρ of the hole density must be determined later on by consistency, for a given value of the chemical potential μ .

In these conditions, it is possible to treat the non-polynomic Lagrangian (2.13) in the framework of the perturbative formalism, and so it can be partitioned as follows

$$L_{eff} = L^B(\mathbf{S}, \lambda) + L^F(\boldsymbol{\eta}) + L^I(\mathbf{S}, \boldsymbol{\eta}) , \quad (5.1)$$

where

$$\begin{aligned} L^B(\mathbf{S}, \lambda) = & -\frac{i}{2s} \sum_i \frac{\tilde{S}_{i1}\dot{\tilde{S}}_{i2} - \tilde{S}_{i2}\dot{\tilde{S}}_{i1}}{s + s'} + 2s' \sum_i \lambda_i \tilde{S}_{i3} \\ & + \frac{1}{8s^2} \sum_{i,I} J' \left[\tilde{S}_{i1}\tilde{S}_{(i+I)1} - \tilde{S}_{i2}\tilde{S}_{(i+I)2} - \tilde{S}_{i3}\tilde{S}_{(i+I)3} + \tilde{S}_i^2 \right] , \end{aligned} \quad (5.2a)$$

and

$$\begin{aligned} L^F(\boldsymbol{\eta}) + L^I(\mathbf{S}, \boldsymbol{\eta}) = & \sum_{i,\sigma} \dot{\Psi}_{i\sigma}^* \Psi_{i\sigma} + \mu \sum_{i,\sigma} \Psi_{i\sigma} \Psi_{i\sigma}^* + \sum_{i,j,\sigma} t_{ij} \Psi_{i\sigma} \Psi_{j\bar{\sigma}}^* \\ & + \sum_i \bar{\boldsymbol{\eta}}_i \mathcal{M}_i \boldsymbol{\eta}_i . \end{aligned} \quad (5.2b)$$

Considering the bilinear bosonic part of the equation (2.13), for a constant value of the hole density and taking $J_{ij} = \text{constant}$, we arrive at the equation (5.2a). In the equation (5.2a) was defined $J' = J(1 - \rho)^2$. Moreover, the symbol \sum_I indicates sum over nearest-neighbor sites.

Besides, in equation. (5.2a) it was assumed that the vector \mathbf{S} is written

$$\mathbf{S} = (0, 0, s') + (\widetilde{S}_1, \widetilde{S}_2, \widetilde{S}_3) \quad (5.3)$$

where $\widetilde{S}_1, \widetilde{S}_2, \widetilde{S}_3$ are the fluctuations. So, the equation (5.2a) corresponds to the lowest order of the system fluctuating around on antiferromagnetic state. Moreover, we must consider $s' \neq s$ because as known the local magnetisation in an antiferromagnetic state reduced from its classical value, even for the pure Heisenberg model. The value of s' must be determined also by consistency.

In equation (5.2b) the four-component spinor $\boldsymbol{\eta} = \begin{pmatrix} \boldsymbol{\Psi} \\ \boldsymbol{\xi} \end{pmatrix}$, is constructed from the two spinors $\boldsymbol{\Psi}$ and $\boldsymbol{\xi}$. The physical two-component spinor $\boldsymbol{\Psi} = \begin{pmatrix} \Psi_+ \\ \Psi_- \end{pmatrix}$ is restricted by the fermionic constraint equations (2.10c,d), and the two-component spinor $\boldsymbol{\xi} = \begin{pmatrix} \xi_+ \\ \xi_- \end{pmatrix}$ is a Majorana spinor.

In the same equation the 4×4 dimensional matrix \mathcal{M} is defined by

$$\mathcal{M} = \begin{pmatrix} 0 & \mathbf{I}s + \mathbf{S} \cdot \boldsymbol{\sigma} \\ \mathbf{I}s + \mathbf{S} \cdot \boldsymbol{\sigma} & 0 \end{pmatrix}, \quad (5.4)$$

where $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ are the Pauli matrices.

In this regime the diagrammatics and the Feynman rules can be found. In particular the bilinear part of the bosonic sector written in equation (5.2a) gives rise to the usual antiferromagnetic magnon propagator (see Ref.[19]).

The bilinear fermionic part $L^F(\boldsymbol{\eta})$ of the equation (5.2b) can be written in terms of the four-component spinor $\boldsymbol{\eta}$ and it is given by

$$L^F(\boldsymbol{\eta}) = \sum_{i,j} \bar{\eta}_{i\alpha} (G_{(0)ij}^{-1})^{\alpha\beta} \eta_{j\beta}, \quad (5.5)$$

where in the Fourier space the symmetric non-singular 4×4 dimensional matrix $G_{(0)}^{-1}$ is defined by

$$(G_{(0)}^{\alpha\beta})^{-1}(k, \nu_n, \nu'_n) = \begin{pmatrix} -(i\nu_n + \mu) & \varepsilon_k & \frac{1}{2}(s + s') & 0 \\ \varepsilon_k & -(i\nu_n + \mu) & 0 & \frac{1}{2}(s - s') \\ \frac{1}{2}(s + s') & 0 & f & g \\ 0 & \frac{1}{2}(s - s') & g & -f \end{pmatrix} \delta(\nu_n, \nu'_n). \quad (5.6)$$

In Eq.(5.6) the quantities k and ν_n are respectively the momentum and the Matsubara frequency of the fermionic field, and was defined $\varepsilon_k = -t \sum_I \exp(-i\mathbf{I} \cdot \mathbf{k})$.

The functions f and g appearing in Eq. (5.6) are totally arbitrary. As easily can be seen, these functions do not appear in the Lagrangian due to the Majorana condition on the two-component spinor $\boldsymbol{\xi}$.

In this scenario, the symmetric matrix defining the fermionic free propagator $G_{(0)\alpha\beta}$ is given by the inverse of the matrix (5.6). The physical components of the free propagator are given by the matrix elements $G_{(0)11}, G_{(0)12}, G_{(0)21}$ and $G_{(0)22}$ and they were explicitly written and analysed in section IV. of Ref.[19]. So, the Feynman rules propagators and vertices are given straightforward and therefore the boson and fermion self-energy can be computed.

However, some particular features of the fermionic free propagator must be emphasised.

The trick of introducing an auxiliary two-component Majorana spinor is a way to obtain a free functional $G_{(0)}$ that really propagates physical fermionic modes.

The electron spectral function is defined from the fermionic propagator $G_{(0)\alpha\beta}$ by considering the components $G_{(0)11}$ and $G_{(0)22}$. The matrix elements directly connected with the electronic properties, as for example the Fermi surface (FS), are precisely $G_{(0)11}$ and $G_{(0)22}$. The electronic spectral function measured in photoemission experiments²⁶ must be related with minus the imaginary part of these matrix elements.

Contrary to the fermion propagator obtained by means of the standard Green function method²⁷, our fermion propagator contains two poles. It is important to say that for a given filled factor the chemical potential we obtain is exactly the same obtained by using the standard Green function method.

By plotting the electron spectral function (see Fig. 1 Ref.[19]) we can see that the peak at negative energy must be interpreted as the extraction of an electron, while the peak at positive energy represents the addition of an electron to the system. Therefore, the two peaks account for the photoemission and the inverse of the photoemission respectively. The presence of these two peaks implies that for a given value of k , the state is not completely filled or empty. Photoemission experiments are only sensitive to the first peak. Then the first peak of our propagator must be related with the excitation measures in photoemission.

An alternative way to treat the fermionic sector is to start from the equation (2.7) and to work inside the path-integral. So, by integrating out the two delta functions on the fermionic constraints Ξ_{i3} and Ξ_{i4} , the partition function can be written as follows

$$Z = \int \mathcal{D}X_i^{\sigma\sigma'} \mathcal{D}X_i^{0+} \mathcal{D}X_i^{+0} \delta(\Omega_{i1}) \delta(\Omega_{i2}) (sdet M_{AB})_i^{\frac{1}{2}} \exp i \int dt L^*(X, \dot{X}) , \quad (5.7)$$

where $L^*(X, \dot{X})$ is given by

$$\begin{aligned} L^*(X, \dot{X}) = & \frac{i}{2} \sum_i \frac{1}{X_i^{++}} \left(X_i^{-+} \dot{X}_i^{+-} - X_i^{+-} \dot{X}_i^{-+} \right) \\ & + \frac{i}{2} \sum_{i,\sigma} \frac{1}{X_i^{++}} \left(\dot{X}_i^{0+} X_i^{+0} + \dot{X}_i^{+0} X_i^{0+} \right) - H(X) . \end{aligned} \quad (5.8)$$

The total Hamiltonian H is defined by

$$H = H_{t-J} + \mu \sum_{i,\sigma} X_i^{0\sigma} X_i^{\sigma 0} , \quad (5.9)$$

where the Hamiltonian H_{t-J} defined in equation (2.9) must be evaluated on the fermionic constraints Ξ_{i3} and Ξ_{i4} .

Due to the non-linearity of the constraints (2.10c,d), when the path-integration on the two fermionic fields X_i^{0-} and X_i^{-0} is carried out, the non-polynomial structure of the kinetic fermionic part of the Lagrangian is made evident, as it can be seen from the equation (5.8).

After the four boson-like X -Hubbard operators are related to the real components S_α ($\alpha = 1, 2, 3$) of a vector field \mathbf{S} and the remaining two fermion-like X -Hubbard operators written in terms of suitable component spinor fields (see equations (2.11)), the correlation generating functional (5.7) takes the form

$$Z = \int \mathcal{D}S_{i1} \mathcal{D}S_{i2} \mathcal{D}S_{i3} \mathcal{D}\Psi_{i-} \mathcal{D}\Psi_{i-}^* \mathcal{D}\lambda_i (s \det M_{AB})_i^{\frac{1}{2}} \left(\frac{\partial X}{\partial S} \right)_i \exp \left(i \int dt L_{eff} \right) . \quad (5.10)$$

Now, the Lagrangian L_{eff} defined in equation (5.10) is given by

$$\begin{aligned} L_{eff} = & \frac{1}{2s} \sum_i (1 - \rho_i) \left(\frac{S_{i1} \dot{S}_{i2} - S_{i2} \dot{S}_{i1}}{s + S_{i3}} \right) + is \sum_i \frac{1}{s + S_{i3}} \left(\dot{\Psi}_{i-}^* \Psi_{i-} + \dot{\Psi}_{i-} \Psi_{i-}^* \right) \\ & - H + \sum_i \lambda_i (S_{i1}^2 + S_{i2}^2 + S_{i3}^2 - s^2) . \end{aligned} \quad (5.11)$$

where in equation (5.11) and hereafter the tilde over the fluctuations is omitted.

The Hamiltonian H written in terms of the real vector variable \mathbf{S} and the two spinor component fields Ψ_{-i} and Ψ_{-i}^* , reads

$$\begin{aligned} H = & \sum_{i,j} \frac{t_{ij}}{(s + S_{i3})(s + S_{j3})} \left[S_{i1} S_{j1} + S_{i2} S_{j2} + S_{i3} S_{j3} + s^2 + s(S_{i3} + S_{j3}) \right. \\ & + i(S_{i1} S_{j2} - S_{i2} S_{j1}) \Psi_{-i} \Psi_{-j}^* + \frac{1}{8s^2} \sum_{i,j} J_{ij} (1 - \rho_i)(1 - \rho_j) [S_{i1} S_{j1} + S_{i2} S_{j2} \\ & \left. + S_{i3} S_{j3} - s^2] + 2s\mu \sum_i \left(\frac{1}{s + S_{i3}} \right) \Psi_{-i} \Psi_{-i}^* . \right] \end{aligned} \quad (5.12)$$

Again, the path-integral (5.8) is considered in the framework of the perturbative formalism at finite temperature, and we assume that we are close to an undoped regime (antiferromagnetic insulator).

After a rotation of spins on the second sublattice by 180° about the S_1 axis is performed the Euclidean and rotated Lagrangian L_{eff}^{ER} is obtained, and so the lowest order of the effective Lagrangian (5.11) can be partitioned as follows

$$L_{eff}^{ER} = L^B(\mathbf{S}, \lambda) + L^F(\Psi_{-i}, \Psi_{-i}^*) + L^I(\mathbf{S}, \Psi_{-i}, \Psi_{-i}^*) , \quad (5.13)$$

where

$$\begin{aligned} L^B(\mathbf{S}, \lambda) = & -\frac{i}{2s}(1-\rho) \sum_i \frac{S_{i1}\dot{S}_{i2} - S_{i2}\dot{S}_{i1}}{s+s'} + 2s' \sum_i \lambda_i S_{i3} \\ & + \frac{1}{8s^2} \sum_{i,I} J' [S_{i1}S_{(i+I)1} - S_{i2}S_{(i+I)2} - S_{i3}S_{(i+I)3} + S_i^2] , \end{aligned} \quad (5.14a)$$

and

$$L^F = \frac{s}{s+s'} \sum_i (\dot{\Psi}_{i-}^* \Psi_{i-} + \dot{\Psi}_{i-} \Psi_{i-}^*) + \frac{2\mu s}{s+s'} \sum_i \Psi_{i-} \Psi_{i-}^* \quad (5.14b)$$

$$L^I = \sum_{i,j} \frac{t_{ij}}{s+s'} (S_{i1} - iS_{i2} + S_{j1} + iS_{j2}) \Psi_{i-} \Psi_{j-}^* \quad (5.14c)$$

At this stage it is important to note that our effective theory does not contain fermion dispersion. This feature is also present in the slave-fermion theories when the fermion dynamics is generated *via* de interaction with virtual magnons.

By making a Fourier transformation it is possible to see that the bilinear bosonic part of the Lagrangian (5.14a) allows to recover the structure of the bosonic propagator (antiferromagnetic magnons) given by

$$\mathcal{D}_{(0)}^{ab}(q, \omega_n, \omega'_n) = \begin{pmatrix} \frac{J'z}{4s^2d_{(0)}}(1-\gamma_q) & -\frac{2\omega_n}{(s+s')d_{(0)}}(1-\rho) & 0 & 0 \\ \frac{2\omega_n}{(s+s')d_{(0)}}(1-\rho) & \frac{J'z}{4s^2d_{(0)}}(1+\gamma_q) & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2s'} \\ 0 & 0 & \frac{1}{2s'} & -\frac{J'z(1-\gamma_q)}{16s^2s'^2} \end{pmatrix} \delta(\omega_n, \omega'_n) . \quad (5.15)$$

In Eq.(5.15) the quantity $d_{(0)}$ is given by

$$d_{(0)} = \left(\frac{2(1-\rho)}{s+s'} \right)^2 (\omega_q^2 + \omega_n^2) , \quad (5.16)$$

where the frequency ω_q^2 is defined by

$$\omega_q^2 = \left[\frac{zJ'}{4s^2} \left(\frac{s+s'}{2(1-\rho)} \right) \right]^2 (1 - \gamma_q^2) . \quad (5.17)$$

Moreover, in Eq.(5.15) z is the number of nearest neighbors and was defined the quantity $\gamma_q = \frac{1}{z} \sum_I \exp(i\mathbf{I} \cdot \mathbf{q})$.

Analogously, the bilinear fermionic part (5.14b) reads

$$L^F = \sum_{k, \nu_n} \Psi_-^*(k, \nu_n) G_0^{-1} \Psi_-(k, \nu_n) , \quad (5.18)$$

where we have named

$$G_0^{-1} = \frac{2s}{s+s'} (i\nu_n - \mu) . \quad (5.19)$$

The inverse of this scalar function given by

$$G_0 = \frac{s+s'}{2s} \frac{1}{i\nu_n - \mu} , \quad (5.20)$$

is a (non-propagating) functional which only depends on the Matsubara frequency ν_n .

Finally, in the approximation that we consider the unique three-leg vertex is defined by

$$U_a = \frac{1}{s+s'} \begin{pmatrix} \varepsilon(k') + \varepsilon(k) \\ i(\varepsilon(k') - \varepsilon(k)) \\ -\frac{s}{s+s'} [i(\nu + \nu') - 2\mu] \\ 0 \end{pmatrix} \delta(q + k - k') \delta(\omega + \nu - \nu') . \quad (5.21)$$

At this point, the problem is to analyse the bilinear fermionic sector, in order to give the prescriptions for the propagation of the fermionic modes. The usual way to solve the propagation of fermions is by means of the Dyson equation. As known the Dyson theorem allows to compute the inverse of the corrected fermion propagator in terms of the free fermion propagator and the self-energy. Therefore the propagator $G(k, \nu_n) = [G_0^{-1}(\nu_n) - \Sigma(k, \nu_n)]^{-1}$ can be calculated in a straightforward way within the self-consistent Born approximation^{28,29}.

By using standard techniques the following expression for the self-energy at zero temperature is found

$$\begin{aligned} \Sigma(k, i\nu_n) = & \frac{(1+\rho)}{2N} t^2 z^2 \sum_q \frac{[\gamma_k[1 - (1 - \gamma_q^2)^{1/2}]^{1/2} - \gamma_{k+q}[1 + (1 - \gamma_q^2)^{1/2}]^{1/2}]^2}{(1 - \gamma_q^2)^{1/2}} \\ & \times \frac{1}{i\nu_n - \omega_q - \mu - \Sigma(k+q, i\nu_n - \omega_q)} \end{aligned} \quad (5.22)$$

The expression (5.22) is useful in the strong coupling case ($t > J$). On the other hand the self-consistent solution of this equation is necessary in order to obtain fermionic propagation, and it must be performed numerically. Once an appropriate self-energy function $\Sigma(k, i\nu_n)$ is found the propagator $G(k, \nu)$ remains well defined and it is possible to compute numerically the spectral function defined by $A(k, \nu) = -\frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} G(k, \nu + i\varepsilon)$. Finally, as well known, the correction to the bosonic propagator is given by

$$\mathcal{D}_{ab} = [\mathcal{D}_{(0)ab}^{-1} - \Pi_{ab}]^{-1}, \quad (5.23)$$

where $\mathcal{D}_{(0)ab}^{-1}$ is the inverse of the free bosonic propagator (5.15) and the bosonic self-energy Π_{ab} reads

$$\Pi_{ab}(q, \omega_n) = \left(\frac{s + s'}{2s} \right)^2 \sum_{k, \nu_n} \frac{U_a U_b}{[i\nu_n + \mu - \Sigma(k, \nu_n)][i(\nu_n + \omega_n) + \mu - \Sigma(q + k, \nu_n + \omega_n)]}. \quad (5.24)$$

As it can be seen from the above equation the dimension of the underlying lattice and the physics depend on the parameters z , γ_q and ε_k , though our general formalism is dimensional independent.

It is important to confront our results with other previously given in the literature related with the spin-polaron theories²⁸. Like in these theories our starting point was to assume an antiferromagnetic order state. This physical assumption is directly connected with the fact that at lowest order the fermion is not propagating (see equation (5.20)). Then, in order to describe a metallic phase where the holes move coherently on the lattice, it is necessary to solve the self-consistent equation (5.22).

The solution of our equations (5.22) and (5.24) together with a quantitative comparison with the spin-polaron theories is an important matter that deserves further study.

Another point to take into account in a future work is to study the relationship between our matricial propagator $G_{(0)\alpha\beta}$ (see equation (4.13) of Ref[19]) and those obtained by solving self-consistently the equation (5.22).

As it is known, any model or approach will be considered as a good candidate to describe high T_c superconductors when it is able to answer the question of why the antiferromagnetic long range order disappears for small values of doping (for instance $\rho = 0.04 - 0.05$). In the last years this problem was attacked from different approaches^{13,14,30}. In a future work and from our formalism, we will also have a response to give about this important point related with the disappearance of the antiferromagnetism.

In the present section, the magnetic excitations that we have considered are antiferromagnetic magnons and in addition we have assumed a strong long range antiferromagnetic order. Therefore, our next step must be to study the instability of the antiferromagnetic order and to analyse against which phase this is unstable. In order to have some idea about this fact, in Ref.[19] we have studied the magnon self energy effects on the magnetic spectral function. Besides the softening of the antiferromagnetic magnon we have also found a reduction of the magnetic spectral signal. These results were got by using our two pole bare fermionic propagator. In the near future and in order to improve our calculation we will solve a self energy coupled problem for both magnetic and electronic dynamics.

VI. CONCLUSIONS

As showed first in Ref.[25] for the pure bosonic case ($\text{su}(2)$ algebra), in a classical Lagrangian formalism it is not possible to introduce the full Hubbard algebra by means of constraints. Consequently, in a path-integral formulation it cannot be introduced the complete information of the Hubbard algebra namely, the commutation rules, the completeness condition and the multiplication rules for the Hubbard X -operators. So, the Heisenberg model treated in the Lagrangian picture only admits two second-class constraints, and these are the completeness condition $X^{++} + X^{--} + X^{00} = 1$ and the non-linear constraint $X^{+-} X^{-+} + \frac{1}{4}(X^{++} - X^{--})^2 = s^2$.

The above last constraint is not really the group Casimir operator. It can be shown that the presence of such constraint is consistent with the quantisation of a spin system in the limit of large spin s , or equivalently for magnetic order state.

A similar situation actually occurs in the case in which the Hubbard X -operators verify the graded algebra $\text{spl}(2,1)$, but in this case at least two solutions are possible. When the Hubbard X -operators closes the graded algebra $\text{spl}(2,1)$ the t-J model described in terms of a first-order Lagrangian has the following possible solutions:

i) One is the family of first-order Lagrangians (2.8) together with the set of second-class constraints (2.10). Two of them are bosonics and the other two are fermionics. In particular the constraint (2.10a) is the completeness condition. As it was shown the corresponding path-integral formalism is mapped into the decoupled slave-fermion representation. So, in this case our correlation generating functional (2.7) privilege the magnon dynamics of the system with a strong feature of magnetic order state (consistent with the large s nature of the constraint (2.10b)).

ii) The different family of first-order Lagrangians (4.9) together with the new set of second-class constraints (4.7) is also a possible solution. In this case all the constraints are bosonic, and (4.7a) is again the completeness condition. As it can be seen the remaining four constraints (4.7b) are related to the multiplication rules. In this situation the bosons are totally constrained and the dynamics is carried out only by the fermions. Moreover, we note that not any non-linear constraint of the type (2.10b) appears. As it was shown the path-integral formalism corresponding to this dynamical situation is mapped in the slave-boson representation. Therefore, in our correlation generating functional (4.10) the fermion dynamics with a strong feature of Fermi liquid is privileged.

It is possible to conclude that once the set of second class-constraints is chosen, different families of Lagrangians are obtained, and so we can insure that each family contains different physics.

Worthwhile to remark that both Lagrangian formalism are independent of the dimension of the underlying lattice.

Moreover, as it was seen in all the cases the completeness condition appears as necessary.

As it is well known the completeness condition involve an important physical meaning. Such condition avoids at the quantum level the double occupancy at each lattice site.

Finally, in section V two alternative way to define the fermion propagator were developed. By means of the trick of introducing auxiliary Majorana spinors, a free fermion matricial propagator having two poles was found. Later on, by integrating out two of the fermions using the delta functions, it was possible to obtain the non-propagating scalar function (5.20) in the fermionic sector. From this free "propagator" and by means of the Dyson equation the fermion self-energy can be evaluated straightforward within the self-consistent Born approximation.

Since our path-integral formalism is mapped in the slave-fermion formalism and taking into account that the equation (5.22) for the self-energy at zero temperature has a similar structure to that obtained from the spin-polaron theories²⁸, both results allows us to insure that our approach is consistent.

REFERENCES

- ¹ A. Izyumov, Physics - Uspekhi **40** (1997), 445.
- ² M. W. Kirson, Phys. Rev. Lett. **78** (1997), 4241.
- ³ A. Angelucci, Physical Review **B51** (1995), 11580.
- ⁴ A. J. Bracken, M.D. Gould, J.R. Links and Y.Z. Zhang, Phys. Rev. Lett. **74** (1995), 2768.
- ⁵ Z. Maassarani, J. Phys. **A28** (1995), 1305.
- ⁶ M. J. Martins and P. B. Ramos, Physical Review **B56** (1997), 6376.
- ⁷ S. Sarkar, Journal of Physics **A24** (1991), 1137.
- ⁸ A. Angelucci and R. Link, Phys. Rev. **B46** (1992), 3089.
- ⁹ A. Förster and M. Karowski, Nucl. Phys **B 396** (1993), 611.
- ¹⁰ "Supersymmetry in models with strong on-site Coulomb repulsion - application to t-J model", T.K. Ng and C. H. Cheng, HKUST, Kowloon, Hong Kong, preprint 1998.
- ¹¹ G. Baskaran, Z. Zou and P.W. Anderson, Solid State Commun. **63** (1987) 973.
- ¹² G. Kotliar and J. Liu, Phys. Review **B 38** (1988) 5142.
- ¹³ C. Jayaprakash, H. R. Krishnamurthy and S. Sarker, Phys. Rev. **B 40** (1989), 2610.
- ¹⁴ C. L. Kane, P. A. Lee, T.K. Ng, B. Chakraborty and N. Read, Phys. Rev. **B 41** (1990), 2653.
- ¹⁵ J. C. Le Guillou and E. Ragoucy, Phys. Rev. **B52** (1995), 2403.
- ¹⁶ J. R. Klauder, Ann. Phys. (NY) **254** (1997), 419.
- ¹⁷ E. . Tüngler and T. Kopp, "Functional integrals for Hubbard operators and projection methods for strong interaction". Preprint Institut für Theorie der Kondensierten Materie, Universität Karlsruhe (1999), and bibliography cuoted therein.
- ¹⁸ A. Foussats, A. Greco and O. S. Zandron, Annals of Physics (NY) **275**(1999), 238.

- ¹⁹ A. Foussats, A. Greco and O. S. Zandron, Annals of Physics(NY) **279**(2000), 263.
- ²⁰ L. D. Faddeev and R. Jackiw, Phys. Rev. Lett.**60** (1988), 1692.
- ²¹ A. Foussats and O. S. Zandron, Journal of Physics **A 30** (1997), L513
- ²² A. Foussats, C. Repetto, O. P Zandron and O. S. Zandron, Annals of Physics(NY) **268** (1998), 225.
- ²³ P. B. Wiegmann, Phys. Rev. Lett. **60** (1988), 821; Nucl. Phys. **B323** (1989), 311.
- ²⁴ P. A. M. Dirac, Can. J. Math. **2** (1950), 129 ; Lectures on Quantum Mechanics (1964),(New York: Yeshiva University Press); K. Sundermeyer, Constrained Dynamics (1982), (Berlin: Springer).
- ²⁵ A. Foussats, A. Greco and O. S. Zandron, Int. J. of Theor.Phys. **38** (1999), 1439.
- ²⁶ Z. X. Shen and D. S. Dessau, Physics Reports **253** (1995) 1.
- ²⁷ G. Mahan "Many - Particle Physics" (1981) Plenum Press, New York.
- ²⁸ G. Martinez and P. Horsch, Physical Review **B44** (1991) 317.
- ²⁹ S. Schmitt-Rink, C. M. Varma and A. E. Ruckenstein, Phys. Rev. Lett **60** (1988) 2793.
- ³⁰ B. Shraiman and E. Siggia, Phys. Rev. Lett **61** (1988) 467; Phys. Rev. Lett **62** (1989) 1565; A. Auerbach and B. Larson, Phys. Rev. B **43** (1991) 7800; J. Igarashi and P. Fulde, Phys. Rev. B **45** (1992) 10419; G. Khaliullin and P. Horsch, Phys. Rev. B **47** (1993) 463; S. Caprara, M. Avignon and V. Tugueshev, Phys. Lett A **255** (1999) 98.