

Long-range interacting classical systems: universality in mixing weakening

Alessandro Campa^{1,‡}, Andrea Giansanti^{2,§}, Daniele Moroni^{2,†} and
Constantino Tsallis^{3,4,5,*}

¹Laboratorio di Fisica, Istituto Superiore di Sanità
and INFN Sezione di Roma1, Gruppo Collegato Sanità
Viale Regina Elena 299, 00161 Roma, Italy

²Dipartimento di Fisica, Università di Roma “La Sapienza”
and INFN Unità di Roma1,
Piazzale Aldo Moro 2, 00185 Roma, Italy

³Centro Brasileiro de Pesquisas Físicas
Rua Xavier Sigaud 150, 22290 -180 Rio de Janeiro - RJ, Brazil

⁴The Santa Fe Institute,
1399 Hyde Park Road, Santa Fe, New Mexico 87501, USA

⁵Department of Mechanical Engineering
Massachusetts Institute of Technology
77 Massachusetts Avenue, Cambridge, Massachusetts 02139, USA

Abstract

Through molecular dynamics, we study the $d = 2, 3$ classical model of N coupled rotators (inertial XY model) assuming a coupling constant which decays with distance as $r_{ij}^{-\alpha}$ ($\alpha \geq 0$). The total energy $\langle H \rangle$ is asymptotically $\propto N\tilde{N}$ with $\tilde{N} \equiv [N^{1-\alpha/d} - (\alpha/d)]/[1 - \alpha/d]$, hence the model is thermodynamically extensive if $\alpha/d > 1$ and nonextensive otherwise. We numerically show that, for energies above some threshold, the (appropriately scaled) maximum Lyapunov exponent is $\propto N^{-\kappa}$ where κ is an *universal* (one and the same for $d = 1, 2$ and 3, and all energies) function of α/d , which monotonically decreases from 1/3 to zero when α/d increases from 0 to 1, and identically vanishes above 1. These features are consistent with the nonextensive statistical mechanics scenario, where thermodynamic extensivity is associated with *exponential* mixing in phase space, whereas *weaker* (possibly *power-law* in the present case) mixing emerges at the $N \rightarrow \infty$ limit whenever nonextensivity is observed.

PACS: 05.20.-y, 05.70.Ce, 05.10.-a

[§]Corresponding author; Andrea.Giansanti@roma1.infn.it

[‡]Campa@iss.infn.it

[†]smoroni@lucifero.phys.uniroma1.it

*tsallis@cbpf.br

In the last few years there has been a noticeable interest in the study of the thermodynamics and statistical mechanics of anomalous systems exhibiting nonextensivity [1, 2]. A system is extensive if its energy and entropy, as functions of intensive internal parameters (e.g. temperature), grow linearly with the size of the system (i.e. with N , the number of its microscopic components). Nonextensivity can be brought into scene by long-range interactions. From the static point of view, nonextensivity means that thermodynamic quantities like the internal or the free energies *per particle* are not constant with N , but rather diverge in the thermodynamic limit $N \rightarrow \infty$. From the dynamical point of view, one might think that long-range interactions induce persistent spatial and temporal correlations leading to the breaking of standard mixing and ergodicity properties, hence to a possible violation of the usual Boltzmann-Gibbs (BG) statistics. Classical Hamiltonian systems with many degrees of freedom are paradigmatic in the discussion of the foundations of equilibrium statistical mechanics. The maximal Lyapunov exponent (MLE) is a well established indicator of chaos. If it is positive, the system will generically be strongly chaotic and will satisfy the standard ergodic hypothesis attained through *exponentially* quick mixing. If it is instead zero non ergodicity could emerge, giving origin to less-than-exponential (typically *power-law*) mixing, hence to anomalous thermostistical behavior. It is then interesting to study maximal Lyapunov exponents in systems with long-range interactions, where nonextensive thermodynamical and dynamical anomalies are expected. In a previous work [3] the maximal Lyapunov exponent of a one-dimensional system of N planar rotators coupled with interactions decaying as the inverse power α of their distances (see model (1) below) has been studied as a function of N . For total energy above some threshold, the maximal Lyapunov exponent is proportional to $1/N^\kappa$, where $\kappa(\alpha)$ is a function which goes from $\kappa(0) > 0$ to zero while α increases from zero to 1 (long range forces) and remains zero for $\alpha > 1$ (short range forces). In this letter we want to investigate if the exponent κ , describing the weakening of the mixing properties of the dynamics in the long-range-interacting regime, is universal. In particular, we check the hypothesis that $\kappa(\alpha, d)$ is a universal function of α/d , where d is the dimensionality of space.

Let us consider a simple model of N planar rotators (XY spins) placed at the sites $i = 1 \dots N$ of a square and a cubic lattice (i.e., dimension $d = 2$ and 3). The rotators have unit moment of inertia, angular momentum L_i and position specified by the angle $\theta_i \in [0, 2\pi]$. Rotations occur inside an arbitrary reference plane. The model is described by the following classical hamiltonian in the conjugate canonical pairs $\{\theta_i, L_i\}$:

$$H = K + V = \frac{1}{2} \sum_i L_i^2 + \frac{1}{2} \sum_{ij, i \neq j} \frac{1 - \cos(\theta_i - \theta_j)}{r_{ij}^\alpha}; \quad (1)$$

where r_{ij} is the distance between lattice site i and j with periodic boundary conditions in the form of nearest image convention; $\alpha \geq 0$ is a variable parameter specifying the range of the interaction, which is defined long-range if $\alpha \leq d$ and short-range if $\alpha > d$.

This system is nonextensive in the case of long-range interactions [4]. To verify that, let us consider as an example a quantity related to the internal energy per rotator at $T = 0$ (i. e., when $K = 0$ and all rotators are equally oriented), namely the integral $\int_1^\infty dr r^{d-1} r^{-\alpha}$, which converges for $\alpha > d$ and

diverges for $0 \leq \alpha \leq d$. In the latter case, the thermodynamics of the model is not well defined in the usual sense. However, it is possible to define energy quantities that are bounded in the thermodynamic limit if they are properly scaled (see [2, 5, 6]). It is interesting to remark that, in system (1) for $0 \leq \alpha \leq d$, not only equilibrium (static) quantities, but also a quantity related to the local instability of the dynamics such as the maximal Lyapunov exponent might diverge with N [7], and needs to be properly rescaled by a function of N if one wants to evaluate the degree of chaoticity as a function of the size of the system.

An alternative procedure is to force the system to be extensive via a largely used mean-field-like rescaling [8] of the potential energy in (1) which then becomes:

$$H' = K' + V' = \frac{1}{2} \sum_i L_i^2 + \frac{1}{2\tilde{N}} \sum_{ij, i \neq j} \frac{1 - \cos(\theta_i - \theta_j)}{r_{ij}^\alpha} \quad (2)$$

The rescaling factor \tilde{N} in front of the potential energy V of hamiltonian (2), is a function of α, d, N and of the geometry of the lattice and has to be chosen to guarantee the existence of a bounded energy density $U = \langle H'/N \rangle$ in the thermodynamic limit.

The model (1) has been introduced in [3] on a one-dimensional lattice and its thermodynamics has been studied by means of simulations in [9]. When $\alpha = 0$ and $\tilde{N} = N$ the model is also called the Hamiltonian Mean-Field XY Model (HMF) [10], whose dynamics and thermodynamics have been extensively studied [11, 12, 13]. In this case the underlying lattice has no meaning and any dependence from its dimension d disappears. It is found that the HMF model has a phase transition from a ferromagnetic state to a paramagnetic one at the critical temperature $T_c = 0.5$ corresponding to a critical energy density $U_c = 0.75$ ($k_B = 1$). In this letter we use [14]

$$\tilde{N} \equiv 1 + d \int_1^{N^{1/d}} dr r^{d-1-\alpha} = \begin{cases} \frac{N^{1-\alpha/d} - \alpha/d}{1-\alpha/d} & \text{if } \alpha \neq d \\ 1 + \ln N & \text{if } \alpha = d \end{cases} \quad (3)$$

whose properties are shown in figure 1. (Unity has been added to the integral in the definition of \tilde{N} in order to have the convenient limit $\tilde{N} \rightarrow 1$ when $\alpha/d \rightarrow \infty$.) Using a similar rescaling factor it was shown in [15] that model (2) and HMF model have the same thermodynamics, and in particular the same critical energy density.

Chaoticity properties and maximal Lyapunov exponent (MLE) of HMF model have been studied by Latora, Rapisarda and Ruffo [13, 16, 17, 18]. They calculated the curve MLE versus U numerically for various N and analytically for $N \rightarrow \infty$; MLE goes to 0 both when U tends to 0 and when U tends to ∞ . In these two limits the system becomes integrable, being represented by a set of harmonic oscillators around the ferromagnetic fundamental state of aligned rotators in the former case and by free rotators in the latter. Finite-size effects are present for $U > U_c$ where MLE vanishes in the thermodynamic limit as $\sim N^{-1/3}$. For fixed $U \leq U_c$ the curve is smooth and presents a positive maximum at the critical energy U_c , where therefore in the thermodynamic limit exists a finite discontinuity.[19]

The behavior of MLE in the one-dimensional model (2) has been studied, as said above, in [3], as a function of N and α at a fixed high energy (above U_c). It

has been shown that it goes to 0 in the thermodynamic limit when $\alpha \leq d$, like in the HMF model, but it remains constant when $\alpha > d$. The way the exponent vanishes is still of the form $\sim N^{-\kappa}$ where now κ is a function of α . The curve $\kappa(\alpha)$ starts from the value $1/3$ at $\alpha = 0$ and goes to 0 when $\alpha \rightarrow 1^-$.

In the present work we extend this study to cases $d > 1$. We still expect that when $\alpha < d$ and $U > U_c$ MLE vanishes in the thermodynamic limit with a power law of the type $N^{-\kappa}$ with $\kappa = \kappa(\alpha, d)$ satisfying $\kappa(\alpha = 0, d) = 1/3$ [19] and vanishing when $\alpha \rightarrow d^-$. Two questions seem quite relevant. Is this curve $\kappa(\alpha, d)$ universal in the sense that it *only* depends on α/d , or it also depends on the dimensionality d of the system? Moreover, is this curve independent from the energy density U at which it is computed? In what follows we will positively answer to both questions.

We have simulated the constant energy dynamics of the nonextensive hamiltonian (1) and computed the maximal Lyapunov exponents.

Hamiltonians H (eq. (1)) and H' (eq. (2)) are strictly connected. If we multiply H' by \tilde{N} , the potential energy V' becomes V . Then, if we rescale time as $t' = t\sqrt{\tilde{N}}$, and using the fact that L_i 's involve first order time derivatives, we obtain that also the kinetic energy K' becomes K . Finally we get $H = \tilde{N}H'$.

Integrating the equations of motion of H is identical to integrating those of $\tilde{N}H'$ but for times longer by a factor $\sqrt{\tilde{N}}$. All thermodynamic quantities that can be obtained as temporal averages of time dependent observables are not influenced by the rescaling of time. In particular, once one has T, V, U from H simulations, one can get the corresponding values for H' simply dividing by \tilde{N} . The evaluation of maximal Lyapunov exponents λ'_{max} of hamiltonian H' can also be obtained through rescaling of the one measured for H , noted λ_{max} hereafter.

In order to calculate maximal Lyapunov exponents one must consider the limit, [20]

$$\lambda_{max} = \lim_{t \rightarrow \infty} \frac{1}{t} \ln \frac{d(t)}{d(0)} = \lim_{t \rightarrow \infty} \lambda(t) \quad (4)$$

with $d(t) = \sqrt{\sum_i (\delta\theta_i)^2 + (\delta L_i)^2}$ being the metric distance calculated from infinitesimal displacements at time t obtained in turn through the double integration of both the normal equations of motion

$$\dot{\theta}_i = L_i \quad (5)$$

$$\dot{L}_i = \sum_{j, j \neq i} \frac{\sin(\theta_j - \theta_i)}{r_{ij}^\alpha} \quad (6)$$

and the linearized ones

$$\delta\dot{\theta}_i = \delta L_i \quad (7)$$

$$\delta\dot{L}_i = \sum_{j \neq i} \frac{\cos(\theta_j - \theta_i)}{r_{ij}^\alpha} (\delta\theta_j - \delta\theta_i) \quad (8)$$

(we report the ones derived from H , the ones for H' are exactly the same with the substitution $r_{ij}^\alpha \rightarrow r_{ij}^\alpha \tilde{N}$). Because of formula (4) which involves $1/t$, time rescaling yields $\lambda'_{max} = \lambda_{max}/\sqrt{\tilde{N}}$.

We computed λ_{max} for typical values of $N \simeq 30$ to 4000 and a fixed energy density $U = 5.0$ above the critical one $U_c = 0.75$. We made simulations on a

square and simple cubic lattices (i.e., $d = 2, 3$) with unitary lattice step, for various values of the α/d ratio.

Simulations started from an high temperature initial state: θ_i 's are randomly extracted from a uniform distribution in $[0, 2\pi]$, momenta L_i 's are extracted from a uniform distribution in $[-0.5, 0.5]$ and then translated and rescaled to have zero total momentum and the desired total energy. We used the velocity-Verlet algorithm [21] with a time-step chosen to have a relative energy conservation of 10^{-4} or better. Length of simulations were chosen looking at the asymptotic behavior of quantity $\lambda(t)$, eq. (4), where $d(0)$ is randomly chosen (see [20]). We report in figure 2, the curves $\lambda'_{max}(N)$ for various α in dimensions $d = 2$ and $d = 3$.

These curves extend those of fig. 3 in [3]. In both cases we observe that, for growing N , if $\alpha > d$ then λ'_{max} is positive and constant, whereas if $\alpha < d$ the maximal Lyapunov exponent tends to zero.

We fitted the data with the following functional form: $\lambda'_{max} \propto N^{-\kappa}$; κ being the slope in the log-log plots of fig. 2.

In figure 3 we collect the slopes κ as a function of the ratio α/d , for $d = 1$ (data from [3]), $d = 2$ and $d = 3$. Remarkably, through the simple scaling α/d we show that the N dependence of the rescaled maximal Lyapunov exponent of model (1) is universal. All curves start at $\alpha = 0$ from a value close to $1/3$, analytically and numerically found for the HMF model [19, 16]; when α/d increases from zero to unity all our data collapse into a single curve, and then remain zero for α/d greater than unity. All of our data are roughly fitted by the heuristic expression $\kappa = [1 - (\alpha/d)^2]/[3 + (\alpha/d)^2/2]$.

Extrapolating for N going to infinity the results in fig. 2, it is interesting to note that the constant κ behavior for $\alpha > d$ is consistent with the results found for the $\alpha = \infty$ (only first-neighboring coupling) model [22].

The universal function κ shown in fig. 3 does not appear to depend on the energy density, provided this is greater than the critical one. For the case $\alpha = 0$ this has been proved analytically in [19] and numerically in [16]. We show in fig. 4 that this is also true for $\alpha = 0.8$ and $d = 2$, the asymptotic slope of λ'_{max} being sensibly the same for the two energy densities $U = 5$, $U = 8$.

Summarizing our results, we can say that the classical hamiltonian H is non extensive for $0 \leq \alpha \leq d$ and extensive for $\alpha/d > 1$. Through a \tilde{N} rescaling of its potential energy it can artificially be made extensive, thus merging with a long standing tradition in mean-field-like approaches. Above some critical value for the (rescaled) total energy per particle, the weakening of the mixing properties is universal; more precisely, the maximal Lyapunov exponents scale as $N^{-\kappa}$, κ being a function *only* of the ratio α/d (and not d nor U). In fact, it seems reasonable to expect that this $\kappa(\alpha/d)$ function is model-independent. More precisely, for total energies high enough, it is expected to have one and the same functional form $\kappa(\alpha/d)$ for *all* classical models whose potential is nowhere singular and whose attractive tail decays as $1/r^\alpha$. Consistently with this speculation, some specific gravitational and fluid models [23] do exhibit $\kappa(0) = 1/3$, in agreement with the present results for the XY models under analysis. An universal $\kappa(\alpha/d)$ fits in fact very well the generic expectation for such models within nonextensive statistical mechanics. Indeed, the available studies of hamiltonian systems including long range interactions provide a variety of indications that a crossover occurs, as a function of time, in the thermostistical properties of the system. For times below a crossover time $\tau(N)$, several properties are anoma-

lous (diffusion, velocity distribution and others), whereas they become normal for times much larger than τ . The fact that τ seems to diverge with N makes the anomalous regime to become the only observable one in the thermodynamic limit. It is precisely this regime which is focused on within that generalized formalism. For the specific case of the mixing properties, the present results suggest that a less-than-exponential mixing takes place until a crossover time after which the mixing becomes exponential, i.e., the usual one. This crossover time might well scale like $1/\lambda'_{max}$ (or a positive power of it). Therefore, since λ'_{max} vanishes in the limit $N \rightarrow \infty$, we expect the crossover time to diverge, thus emerging the anomalous regime we were just discussing.

References

- [1] C. Tsallis, J. Stat. Phys. **52**, 479 (1988); E.M.F. Curado and C. Tsallis, J. Phys. A **24**, L69 (1991) [E: **24**, 3187 (1991) and **25**, 1019 (1992)]; C. Tsallis, R.S. Mendes and A.R. Plastino, Physica A **261**, 534 (1998).
- [2] A set of mini-reviews is contained in "Nonextensive Statistical Mechanics and Thermodynamics", eds. S.R.A. Salinas and C. Tsallis, Braz. J. Phys. **29** (1999) [<http://sbf.if.usp.br/WWWpages/Journals/BJP/Vol29/Num1/index.htm>]. See <http://tsallis.cat.cbpf.br/biblio.htm> for further bibliography on the subject.
- [3] C. Anteneodo and C. Tsallis, Phys. Rev. Lett. **80**, 5313 (1998).
- [4] M.E. Fisher, Arch. Rat. Mech. Anal. **17**, 377 (1964); J. Chem. Phys. **42**, 3852 (1965); J. Math. Phys. **6**, 1643 (1965); M.E. Fisher and D. Ruelle, J. Math. Phys. **7**, 260 (1966); M.E. Fisher and J.L. Lebowitz, Commun. Math. Phys. **19**, 251 (1970).
- [5] C. Tsallis, Fractals **3**, 541 (1995).
- [6] R. Salazar and R. Toral, Phys. Rev. Lett. **83**, 4233 (1999).
- [7] Definition (4) assumes an exponential separation. It is our opinion that a diverging Lyapunov exponent indicates that the rate of separation in these cases is more than exponential.
- [8] This rescaling implies that the microscopic coupling between rotators is a function of the size N of the system which is a macroscopic quantity.
- [9] F. Tamarit and C. Anteneodo, Phys. Rev. Lett. **84**, 208 (2000).
- [10] M. Antoni and S. Ruffo, Phys. Rev. E **52**, 2361 (1995).
- [11] M. Antoni and A. Torcini, Phys. Rev. E **57**, R6233 (1998).
- [12] V. Latora, A. Rapisarda and S. Ruffo, Phys. Rev. Lett. **83**, 2104 (1999).
- [13] V. Latora, A. Rapisarda, and S. Ruffo, Prog. Theor. Phys. Supplement. (2000), in press [cond-mat/0001010]
- [14] C. Tsallis, Braz. J. Phys **29**, 1 (1999).

- [15] A. Campa, A. Giansanti and D. Moroni, Phys. Rev. E (2000), in press [cond-mat/0002168].
- [16] V. Latora, A. Rapisarda, and S. Ruffo, Phys. Rev. Lett. **80**, 692 (1998).
- [17] V. Latora, A. Rapisarda, and S. Ruffo, Physica D **131**, 38 (1999).
- [18] V. Latora, A. Rapisarda, and S. Ruffo, Physica A **81**, 280 (2000).
- [19] M. C. Firpo, Phys. Rev. E **57**, 6599 (1998).
- [20] G. Benettin, L. Galgani and J. M. Strelcyn, Phys. Rev. A **14**, 2338 (1976).
- [21] W. C. Swope, H. C. Andersen, P. H. Berens and K. R. Wilson, J. Chem. Phys. **76**, 637 (1982).
- [22] P. Butera and G. Caravati, Phys. Rev. A **36**, 962 (1987).
- [23] V.G. Gurzadyan and G.K. Savvidy, Astron. Astrophys. **160**, 203 (1986); Ch. Dellago and H.A. Posch, Physica A **240**, 68 (1997); M. Antoni, S. Ruffo and A. Torcini, preprint [cond-mat/9908336].




Figure 1: Scaling factor $\tilde{N}(N, \alpha/d)$: (a) versus α/d , for typical values of N (dotted line: the $N \rightarrow \infty$ asymptote); (b) versus N , for typical values of α/d (dotted line: the $\alpha/d = 2$ asymptote). Note the divergence of \tilde{N} in the thermodynamic limit ($N \rightarrow \infty$), in the long-range case.




Figure 2: Curves λ'_{max} vs N (log-log scale) for $d = 2$ (upper part) and $d = 3$ (lower part), $U = 5$. Solid lines are fits with the functional form $(a - \frac{b}{N}) / (\tilde{N})^c$. Consequently, $\lambda'_{max} \propto N^{-\kappa(\alpha, d)}$ where $\kappa(\alpha, d) = (1 - \alpha/d)c(\alpha, d)$ for $0 \leq \alpha < d$ and $\kappa(\alpha, d) = 0$ for $\alpha > d$; for $\alpha = d$, λ'_{max} is expected to vanish as a power of $1/\ln N$. The ordinate indistinctively represents λ'_{max} corresponding to Hamiltonian (2) with $U = H'/N = 5$, or $\lambda_{max}/\sqrt{\tilde{N}}$ corresponding to Hamiltonian (1) with $U = H/(N\tilde{N}) = 5$.




Figure 3: The mixing weakening exponent κ vs α/d for $d = 1, 2, 3$ ($d = 1$: from [3]; $d = 2, 3$: present work). It describes the asymptotic N behaviour of the maximal Lyapunov exponent λ'_{max} at fixed energy above the critical one, i.e., $\lambda'_{max} \propto 1/N^\kappa$. The solid line is a guide to the eye consistent with universality. For $\alpha = 0$ we have [19] $\kappa(0) = 1/3$ ($\forall d$).




Figure 4: $\lambda'_{max}(N)$ for fixed $\alpha = 0.8$, $d = 2$ and two different energies U . The asymptotic N behavior, for all values of U above the critical one $U_c(\alpha, d)$ and $0 \leq \alpha/d < 1$, appears to be $\lambda'_{max} \propto 1/N^{\kappa(\alpha/d)}$, where the proportionality coefficient decreases from a finite (α, d) -dependent positive value to zero when U increases from $U_c(\alpha, d)$ to infinity.