

A protein model exhibiting three folding transitions

Audun Bakk

Department of Physics, Norwegian University of Science and Technology, NTNU,
N-7491 Trondheim, Norway

Alex Hansen¹ and Kim Sneppen

NORDITA and Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen,
Denmark

(April 14, 2024)

Abstract

We explain the physical basis of a model for small globular proteins with water interactions. The water is supposed to access the protein interior in an "all-or-none" manner during the unfolding of the protein chain. As a consequence of this mechanism (somewhat speculative), the model exhibits fundamental aspects of protein thermodynamics, as cold and warm unfolding of the polypeptide chain, and hence decreasing the temperature below the cold unfolding the protein folds again, accordingly the heat capacity has three characteristic peaks. The cold and warm unfolding has a sharpness close to a two-state system, while the cold folding is a transition where the intermediate states in the folding is energetical close to the folded/unfolded states, yielding a less sharp transition. The entropy of the protein chain causes both the cold folding and the warm unfolding.

PACS: 05.70.Jk, 87.14.Ee, 87.15.Cc, 87.10.+e

1 Introduction

In order to have a precise function in the biological "machinery", it is important for proteins to have a unique conformation at physiological temperatures. This is termed the native state. Anfinsen [1] proved in his famous experiment with ribonuclease the important fact that the folding of the polypeptide chain is thermodynamically determined.

One simplified view of protein folding is that the protein is supposed to follow a specific folding pathway of conformations in a descending landscape of Gibbs free energy [2,9]. This is a picture of a folding protein that is forced to follow a specific "path" of successive conformational steps of increasing structural order. We will use this pathway assumption in this paper.

A protein in physiological environments (pH, ionic strength etc.), and temperatures is packed in a very compact way. It is then termed folded. An increase of the temperature will eventually denature the protein, i.e. it unfolds. Other ways to unfold the protein are for instance to change the pressure, denaturant concentration or the pH. The fact that proteins also unfold at low temperatures, termed as cold unfolding [10,11], makes the system very unusual. A major difficulty in experiments of cold unfolding is that the temperature is around and below the freezing point of water. In a frozen aqueous solution, one cannot observe any conformational transitions [12].

¹Permanent Address: Department of Physics, Norwegian University of Science and Technology, NTNU, N-7491 Trondheim, Norway

A general feature of small globular proteins is that they thermodynamically seem to unfold in an "all-or-none" manner. This means that they unfold cooperatively without noticeable intermediates [13,18], with a deviation from a two-state system not exceeding 5%. The deviation from a single macroscopic system can be explained by presence of unstable intermediates [8,14,20]. It is worth noting that all these experiments have been done only for the warm unfolding. The occurrence of intermediate states in larger proteins [2,3,19] is not a contradiction to the two-state behavior in the experiments in Refs. [13,18], because the latter only considers small globular proteins.

The van't Hoff enthalpy relation (for heat of reaction) [14,21]

$$H = k_B T_c^2 \frac{C}{Q} ; \quad (1)$$

is a powerful way to quantify the sharpness of a smoothed out first order phase transition. As shown in Fig.1, T_c is the transition temperature (at the middle of the peak), Q , which is the same as H (no pressure), is the released energy (latent heat), and C is the peak height of the transition. k_B is a dimensionless proportionality factor. For a given H and Q , then the value of T_c is inversely proportional to C . In this respect a smaller T_c corresponds to a sharper transition.

In this article we will explain the physical basis of a protein model, that reformulates the water interactions proposed in earlier models by Hansen et al. [5,6] and Bakk et al. [8,9]. We will compare thermodynamical quantities, as the heat capacity, to experiments. The protein is also investigated in a temperature region below accessible experimental data.

2 Modeling the protein

2.1 The polypeptide chain

The polypeptide chain is modeled as in earlier articles by Hansen et al. [5,7] and Bakk et al. [8,9], where the protein is supposed to follow a pathway as described in Section 1. The protein is equipped with N contact points, which we here call nodes. A node in contact means that the protein has a "correct" conformation on the folding pathway, and we say that the node is folded. Due to the fact that a protein is a complex 3-dimensional system, a folded node likely has non-local contacts with respect to the amino acid sequence in the polypeptide chain.

Each node is assigned only two energies $\epsilon_i = 0$ or 1 [22,23]. By using the binary variables $\epsilon_i \in \{0,1\}$, the energy associated to each individual node is written $E_i = \epsilon_i \epsilon_{i-1} \epsilon_{i-2} \dots \epsilon_1$. The value $\epsilon_i = 0$ means an unfolded node, while $\epsilon_i = 1$ is equivalent to a folded node. The product term meets the assumption about a folding pathway, because $E_i = \epsilon_i$ only if all nodes before i is folded, in addition to a folded node i itself. For a system of N nodes the Hamiltonian of the polypeptide chain (vacuum energy of the protein) is

$$H_c = \sum_{i=1}^N \epsilon_i \epsilon_{i-1} \epsilon_{i-2} \dots \epsilon_1 \quad (2)$$

Folding of node i is only proper in the case of an unique conformation before this folding step. According to Eq.2, every attempt to fold i while j (describing one or several nodes $< i$) is unfolded will not gain energy in the system, because then the protein is supposed to be in an energetical unfavorable conformation.

The unfolded protein has more degrees of freedom relative to the folded, because the unfolded polypeptide backbone will have rotational freedom. We incorporate

the water entering upon unfolding of node $N-1$, will not interact with the protein surface, because it is regarded as a second layer of "free" water. Choh and Edsall [27] states that roughly a monolayer of water is bounded to the protein, implying that the protein is only interacting with the first monolayer, thus the second, and third etc. water layers, successively entering the protein during unfolding, are regarded as "free water". Hence, according to the latter possible (but somewhat speculative) explanation of how the water access the apolar interior of the protein, unfolding of nodes $i < N$ does not contribute energetical to the water Hamiltonian (H_w) and thus not to the thermodynamics.

2.3 The statistical framework

The system Hamiltonian (H) describing both chain specific energy (H_c) and water interactions (H_w) is

$$H = H_c + H_w = \sum_{i=1}^N (\epsilon_i + \sum_{j=1}^{i-1} \epsilon_{ij}) + \sum_{i=1}^N (1 - \epsilon_i) \sum_{j=1}^M \epsilon_{ij} \quad (5)$$

Let now Z_i be term number i in the partition function which corresponds to folding of all nodes i (pathway assumption), thus

$$Z_i = f^{N-i} e^{i\epsilon_0} e^{\beta \sum_{j=1}^M \epsilon_{ij}} \sum_{g=0}^{\infty} \frac{e^{-g}}{1 - e^{-g}}^M \quad (i < N) \quad (6)$$

$\beta = 1/T$ is a rescaled inverse absolute temperature in which the Boltzmann constant is absorbed. Z_0 is the term where all nodes are zero, i.e. a complete unfolded protein, while Z_N corresponds to a folded protein. The factor f^{N-i} in Eq.6 is the degrees of freedom in the polypeptide chain that is available in the $N-i$ unfolded nodes. $e^{i\epsilon_0}$ is the Boltzmann factor from i contact energies ϵ_0 in the polypeptide chain. The last term in brackets is simply the sum over all distinct levels in one water molecule raised to the power of the number of water molecules M bounded to the unfolded parts of the protein. We assume that $\epsilon_0 \gg 1$ (i.e. $g \gg 1$), which is equal to an infinite small level spacing in Eq.3. A first order Taylor expansion of the denominator in Eq.6 yields

$$Z_i = f^{N-i} \frac{e^{\beta \sum_{j=1}^M \epsilon_{ij}}}{1 - e^{-g}}^M e^{i(\epsilon_0 - \ln f)} \quad (i < N) \quad ; \quad (7)$$

assuming $1 - e^{-g} \approx 1$ when $g \gg 1$. The last term in the partition function (Z_N) corresponds to a complete folded protein, where there are g^M degrees of freedom from M unbounded water molecules and N contact energies ϵ_0 , hence

$$Z_N = g^M e^{N\epsilon_0} \quad (8)$$

By summing up the Z_i terms in Eqs.7 and 8, we obtain the partition function

$$Z = \sum_{i=0}^N Z_i = f^N g^M \frac{ae^{\beta \sum_{j=1}^M \epsilon_{ij}}}{1 - r} + r^N \quad ; \quad (9)$$

where $r = e^{-\ln f}$, $a = e^{\beta \sum_{j=1}^M \epsilon_{ij}}$ and $\beta \epsilon_0 = g$. The inverse temperature is here rescaled by ϵ_0 . The parameter m measures the strength of the water interactions relative the chain contact energy, thus m is interpreted as an effective chemical potential. Changing m means adding denaturants, changing pH or salt concentration etc.

The order parameter n in this system measures the degree of folding, i.e. the mean number of folded nodes divided by N

$$n = \frac{\sum_{i=0}^N i Z_i}{Z} = \frac{1}{N} \frac{\frac{ae}{1-r} \frac{(N-1)r^{N+1} + Nr^N + r}{(1-r)^2} + Nr^N}{\frac{ae}{1-r} + r^N} \quad (10)$$

$n = 0$ corresponds to an unfolded protein, while $n = 1$ is interpreted as a completely folded protein.

3 Calculations and discussion

The heat capacity is $C = -k_B T^2 \partial^2(\ln Z) / \partial T^2$. Fig.2 shows a typical plot of the heat capacity $C(T)$ with three peaks (numbered 1, 2 and 3 from left). These characteristic peaks corresponds to three critical transition temperatures: T_1, T_2 and T_3 , measuring the temperatures at the respective peak maxima. The corresponding order parameter $n(T)$ in Fig.3, calculated from Eq.10, shows that the protein is essential folded for $T < T_1$ and $T_2 < T < T_3$, while the protein is nearly unfolded in the temperature intervals $T_1 < T < T_2$ and $T > T_3$. From this picture it is reasonable to state that the physiological temperature interval is between peak 2 and 3. Accordingly, with reference to this temperature region, we call peak 1 for cold folding and peak 2 and 3 respectively for cold and warm unfolding. Peak 2 and 3 are both observed in experiments [10,11] and are also seen in the model of Hansen et al. [5,6] and Bakk et al. [8,9]. The model considered in this paper has, in addition to the cold and warm unfolding, the peculiarity of cold folding.

Experiments on cold unfolding are very difficult because most proteins unfolds below the freezing point of water. Chen and Chellman [11] and Pivalov et al. [28] have all done experiments where the cold unfolding temperature is elevated by denaturants, but denaturants make the interpretation of the data very difficult. However, Pivalov [10] did experiments in super cooled water, which is easier to interpret. Unfortunately he was not able to detect the sharpness of the cold unfolding, and not at all the heat capacity below the cold unfolding. This means that our model may predict a cold folding transition at a temperature below the cold unfolding transition.

For temperatures below the cold folding ($T \rightarrow 0$) analysis of the model gives $n \rightarrow 0.99$, i.e. only the last node is unfolded, and corresponds to the global energy minimum. From Fig.2 it is seen that $T_1 = 1.45$. In Eq.9 the critical $r = e^{1/T} \ln f = 1$, caused by the contact energies of the polypeptide chain, implies $T_1 = 1/\ln 2 = 1.44$. Hence, this is nothing but a transition initiated of the chain entropy. An increase of the temperature from T_1 takes the protein through a nearly unfolded state, whereupon the protein folds again at $T_2 = 1.8$. The temperature is now so high that the energy of water (Eq.3) is small (thermally excited) compared to the chain contact energy ϵ_0 , thus the protein prefers to fold again. Further increase of the temperature causes warm unfolding at $T_3 = 3.0$, because then the entropy of the chain again dominates the Gibbs free energy. It is interesting to note that the entropy of the chain causes two transitions, the cold folding and warm unfolding.

We now turn our interest to the sharpness of the transitions, i.e. the value of the parameter β in the van't Hoff enthalpy relation (Eq.1). For $M = 200$ is $\beta = 4$ both for the cold and warm unfolding. This means that the protein is thermodynamically regarded as a two-state system that folds in an "all-or-none" manner. Pivalov [14] has measured $\beta = 4.0$ for the warm unfolding. As far as we know there are no experiments on the sharpness of the cold unfolding, but Pivalov [10] indicates a sharpness for the cold unfolding as well, thus according to our model. The cold

folding transition has $\Delta G \approx 12$. This value is typical for a transition where one has small energy differences between the folded/unfolded states and the intermediate states. Remember that the "folded" state for $T < T_1$ is actually the first node unfolded, thus the unfolding will essentially depend on the polypeptide chain with the Hamiltonian in Eq.2, which can be shown corresponds to $\Delta G \approx 12$ [8,20].

Finally we note the consequence of a decreased ΔG is an increasing separation between the cold and warm unfolding as seen in Fig.4. This makes sense because a smaller ΔG is equivalent to a relatively smaller ΔG_w compared to ΔG_0 (see Eq.9), i.e. it is less favorable for the protein to be bounded to water. The consequence is that the protein prefers to be folded in a larger temperature interval, in where the water is expelled to the bulk. However, the transition temperature T_1 is not changed because this transition is given by the value $T_1 = 1/\ln f$. It is also seen that a smaller ΔG is qualitatively equivalent to a smaller a .

An increase in M is the same as a decrease in N , because then the water becomes more important relative to the chain, and will again allow a broader separation between T_2 and T_3 . The broader separation is also seen for a decreasing f , because this is equivalent to a larger M .

Further increase of ΔG will eventually merge peak 1 and 2. It is interesting to note that $\Delta G = 4$ for the merged peaks, because then the transition is energetically dominated of the M water molecules which caused the transition at T_2 in Fig.2.

In sum the qualitative change from Fig.3 to Fig.4, by a decreasing ΔG , is also obtained by an decrease of a , f or N or an increase of M .

4 Conclusion

We have in this paper studied a protein model with water interactions. The model is based on earlier models by Hansen et al. [5,6] and Bakke et al. [8,9]. In contrast to these similar models, where the water amount was supposed to increase linearly to the degree of unfolding of the polypeptide chain, we have, with a more speculative assumption, studied the situation where a macroscopic amount of water access the protein interior during unfolding of the last node is the only contribution to the water-protein Hamiltonian.

With reference to physiological temperatures we find that the protein exhibits cold and warm unfolding transitions, which is an experimental fact [10,11]. These transitions are associated by a sharpness indicating, from a thermodynamic point of view, a two-state system, which is also experimentally established [13,18]. Decreasing the temperature further below the cold unfolding region the protein folds again (cold folding). This folding, caused by the chain entropy, has a less sharp transition, which corresponds to a transition where the intermediate folding energies does not differ significantly from the folded/unfolded energies. In sum the model exhibits three unfolding/folding transitions.

It is interesting to note that both the cold folding and the warm unfolding is due to the polypeptide chain entropy.

Acknowledgements

A.B. thanks the Norwegian Research Council for financial support. A.H. thanks NORDITA and Niels Bohr Institute for warm hospitality and support. We thank J.S.H. Ye for enthusiastic and enlightening discussions.

References

- [1] C. B. Anfinsen. Principles that govern the folding of a protein chain (Nobel Lecture). *Science* 191, 223-230 (1973).
- [2] R. L. Baldwin, and G. D. Rose. Is protein folding hierarchic? I. Local structure and peptide folding. *TIBS* 24, 26-33 (1999).
- [3] R. L. Baldwin, and G. D. Rose. Is protein folding hierarchic? II. Folding intermediates and transition states. *TIBS* 24, 77-83 (1999).
- [4] Yi Wang, and D. Shortle. The Equilibrium Folding Pathway of Staphylococcal Nuclease: Identification of the Most Stable Chain-Chain Interactions by NMR and CD Spectroscopy. *Biochem.* 34, 15895-15905 (1995).
- [5] A. Hansen, M. H. Jensen, K. Sneppen, and G. Zocchi. Statistical mechanics of warm and cold unfolding in proteins. *The European Phys. J. B.* 6, 157-161 (1998).
- [6] A. Hansen, M. H. Jensen, K. Sneppen, and G. Zocchi. A model for the thermodynamics of globular proteins. *Physica A.* 270, 278-287 (1999).
- [7] A. Hansen, M. H. Jensen, K. Sneppen, and G. Zocchi. A hierarchical scheme for cooperativity and folding in proteins. *Physica A* 250, 355-361 (1998).
- [8] A. Bakk, J. S. H. Ye, A. Hansen, K. Sneppen, and M. H. Jensen. Pathways in Two-State Protein Folding. Submitted to *Biophysical Journal* (2000).
- [9] A. Bakk, J. S. H. Ye, A. Hansen, and K. Sneppen. Thermodynamic implications of a protein model with water interactions. Submitted to *Journal of Theoretical Biology* (2000).
- [10] P. L. Privalov. Cold Denaturation of Proteins. *Biochem. and Mol. Biol.* 25, 281-305 (1990).
- [11] B. Chen, and J. A. Schellman. Low-Temperature Unfolding of a Mutant of Phage T4 Lysozyme. I. Equilibrium Studies. *Biochem.* 28, 685-691 (1989).
- [12] T. E. Creighton. *Protein folding*. W. H. Freeman and Company, New York, Ch. 3 (1992).
- [13] P. L. Privalov, and N. N. Khechinashvili. A Thermodynamic Approach to the Problem of Stabilization of Globular Protein Structure: A Calorimetric study. *J. Mol. Biol.* 86, 665-684 (1974).
- [14] P. L. Privalov. Stability of Proteins, Small Globular Proteins. *Biochem. Adv. Protein Chem.* 33, 167-241 (1979).

- [15] G. I. Makhataдзе, and P. L. Privalov. Contribution of hydration to protein folding thermodynamics. I. The enthalpy of hydration. *J. Mol. Biol.* 232, 639-659 (1993).
- [16] S. E. Jackson, and A. R. Fersht. Folding of chymotrypsin inhibitor 2. Evidence for a two-state transition. *Biochemistry* 30, 10428-10435 (1991).
- [17] S. J. Bae, and J. M. Sturtevant. Thermodynamics of thermal unfolding of eglin c in the presence and absence of guanidinium chloride. *Biophys. Chem.* 55, 247-252 (1995).
- [18] Y. V. Grikko, G. I. Makhataдзе, P. L. Privalov, and R. W. Hartley. Thermodynamics of barnase unfolding. *Protein Sci.* 3, 669-676 (1994).
- [19] D. Shortle, Y. Wang, J. R. Gillespie, and J. O. Wrabel. Protein folding for realists: A timeless phenomenon. *Protein Sci.* 5, 991-1000 (1996).
- [20] P. G. Dommersnes, A. Hansen, M. H. Jensen, and K. Sneppen. Parameterization of Multiple Pathways in Proteins: Fast Folding versus Tight Transitions. Submitted to *Proc. Roy. Soc., London Ser. A*.
- [21] H. B. Callen. Thermodynamics and an introduction to thermostatistics (2. ed.). John Wiley & Sons, New York, Ch. 13 (1985).
- [22] J. D. Bryngelson, and P. G. Wolynes. Spin-glasses and the statistical mechanics of protein folding. *Proc. Natl. Acad. Sci. USA* 84, 7524-7528 (1987).
- [23] J. D. Bryngelson, and P. G. Wolynes. A simple statistical field-theory of collapse with application to protein folding. *Biopolymers* 30, 177-188 (1990).
- [24] P. L. Privalov, and G. I. Makhataдзе. Contribution of Hydration and Non-covalent Interactions to the Heat Capacity Effect on Protein Unfolding. *J. Mol. Biol.* 224, 715-723 (1992).
- [25] G. I. Makhataдзе, and P. L. Privalov. Energetics of Protein Structure. *Adv. Prot. Chem.* 47, 308-424 (1995).
- [26] P. L. Privalov. Intermediate States in Protein Folding. *J. Mol. Biol.* 258, 707-725 (1996).
- [27] E. J. Cohn, and J. T. Edsall. *Proteins, Amino Acids and Peptides*. Reinhold Publishing, New York (1943).
- [28] P. L. Privalov, Yu. V. Grikko, and S. Yu. Venyaminov. Cold Denaturation of Myoglobin. *J. Mol. Biol.* 190, 487-498 (1986).

Figure captions


Fig. 1. Schematic illustration of the heat capacity around an unfolding transition showing the parameters in the van't Hoff enthalpy relation (Eq.1). T_c is the transition temperature, Q (area of the peak) is the released energy (latent heat) and C is the peak height of the transition.

Fig. 2. Heat capacity $C(T)$ with the parameters $a = 0.077$, $\epsilon = 3.3$, $f = 2$, and $N = M = 200$, showing three peaks. With reference to the temperature region between peak 2 and 3 (physiological temperatures) we call the transitions: 1) cold folding, 2) cold unfolding and 3) warm unfolding.

Fig. 3. The corresponding order parameter $n(T)$ to Fig.2, describing the degree of folding. The chosen parameters are as in Fig.2. $n = 0$ corresponds to an unfolded protein, while $n = 1$ is interpreted as a completely folded protein.


Fig. 4. Heat capacity where the effective chemical potential is $\epsilon = 3.2$, slightly decreased from the value in Fig.2 ($\epsilon = 3.3$). All other parameters are chosen as in Fig.2. The qualitative picture is a broader separation between T_2 and T_3 compared to Fig.2.

Figure 1:




A .Bakk, A .Hansen and K .Sneppen
A protein model exhibiting three folding transitions

Figure 2:




A .Bakk, A .Hansen and K .Sneppen
A protein model exhibiting three folding transitions

Figure 3:



A .Bakk, A .Hansen and K .Sneppen
A protein model exhibiting three folding transitions

Figure 4:



A .Bakk, A .Hansen and K .Sneppen
A protein model exhibiting three folding transitions