

Comment on "A Kac-potential treatment of nonintegrable interactions" by Volm ayr-Lee and Luijten

Constantino Tsallis

Centro Brasileiro de Pesquisas Físicas, Rua Xavier Sigaud 150
22290-180 Rio de Janeiro-RJ, Brazil (tsallis@cbpfr)

We comment the recent manuscript by Volm ayr-Lee and Luijten [cond-mat/0009031] focusing on classical systems with nonintegrable interactions. The authors claim that they have proved that Boltzmann-Gibbs statistics suffices for describing the system in thermalequilibrium. We show that this statement is a misleading oversimplification since it only applies for the $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty}$ ordering, but certainly not for the $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty}$ one. The latter can even be the unique physically meaningful situation for thermodynamically large systems.

In a recent paper [1], Volm ayr-Lee and Luijten (VLL) present a Kac-potential approach of nonintegrable interactions. They consider a d -dimensional classical fluid with two-body interactions exhibiting a hard core as well as an attractive potential proportional to r^{-d} with $0 < d < 1$ (logarithmic dependence for $d = 1$) [2]. In their approach, they also include a Kac-like long-distance cutoff R such that no interactions exist for $r > R$, and then discuss the $R \rightarrow \infty$ limit. They show that the exact solution within Boltzmann-Gibbs statistical mechanics is possible and that (no surprise (see VLL Ref. [12] and references therein)) it exhibits a mean field criticality. Moreover, the authors argue that very similar considerations hold for lattice gases, $O(n)$ and Potts models.

Almost as an illustration of the fact that "It is the strange privilege of statistical mechanics to stimulate and nourish passionate discussions related to its foundations [...] [3], VLL state "Our findings imply that, contrary to some claims, Boltzmann-Gibbs statistics is sufficient for a standard description of this class of nonintegrable interactions.", and also that "we show that nonintegrable interactions do not require the application of generalized q -statistics.". VLL also inform us that "the main motivation for this [their] work stems from the considerable attention systems with nonintegrable interactions have received in the context of "nonextensive thermodynamics."".

It is the purpose of the present Comment to argue that VLL's (first two) above statements severely misguide the reader. Indeed, the interesting VLL discussion, along traditional lines, of their specific Kac-like model only exhibits that Boltzmann-Gibbs statistical mechanics is (as more than one century of brilliant successes guarantees!) necessary for obtaining thermal equilibrium properties without needing to do time averages; by no means it proves that it is sufficient, as we shall soon clarify. Neither it proves that wider approaches (such as, for instance, nonextensive statistical mechanics, VLL Refs.

[6,31] and present Ref. [4], or a similar formalism) are not required or convenient. The crucial point concerns time, a word that nowhere appears in the VLL paper. This is, in fact, rather surprising since the key role of t has been strongly emphasized in several occasions, for instance in VLL Ref. [31] (e.g., Fig. 4 of VLL Ref. [31] illustrates the expectation). For integrable (or "short-range", as frequently referred to in the literature focusing the present context [5]) two-body interactions in a N -body classical Hamiltonian system, i.e., for $d > 1$, we expect that the $t \rightarrow \infty$ and $N \rightarrow \infty$ limits are commutable in what concerns the equilibrium distribution $p(E)$, E being the total energy level associated with the macroscopic system. More precisely, we expect naturally that

$$p(E) = \lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty} p(E; N; t) = \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} p(E; N; t) / \exp[-E/kT] \quad (d > 1) \quad (1)$$

if the system is in thermal contact with a thermostat at temperature T . In contrast, the system is expected to behave in a more complex manner for nonintegrable (or "long-range") interactions, i.e., for $0 < d \leq 1$. In this case, no generic reasons seem to exist for the $t \rightarrow \infty$ and $N \rightarrow \infty$ limits to be commutable, and consistently we expect not necessarily equal results. The simplest of these results (which is in fact the one to be associated with the VLL paper, although therein these two relevant limits and their ordering are not mentioned) is, as we shall soon further comment,

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} p(E; N; t) / \exp[-(E/N^d)/(kT/N^d)] \quad (2)$$

where we have introduced $N^d = N^{1-d} = d = [1 - d]$ in order to stress the facts that generically

(i) E is not extensive, i.e., is not proportional to N (but is E/N^d instead; more precisely, E is extensive if $d > 1$, see [6] and VLL Refs. [4,5], and it is nonextensive if $0 < d \leq 1$),

and

(ii) T needs to be rescaled (a feature which is frequently absorbed in the literature by artificially size-rescaling the coupling constants of the Hamiltonian), in order to guarantee nontrivial finite equations of states. Of course, for $d = 0$, we have $N^d = N$, which recovers the traditional Mean Field scaling.

But, depending on the initial conditions, which determine the time evolution of the system if it is assumed isolated, quite different results can be obtained for the ordering $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty} p(E; N; t)$. This fact has been

profusely detected and stressed in the related literature (see, for instance, VLL Ref. [31], present Refs. [4,7{11] and references therein).

Let us discuss this point further. The impressive success of Boltzmann-Gibbs statistical mechanics relies on the fact that, for ubiquitous Hamiltonian systems, time averages (whose calculation is typically untractable) can be replaced by appropriate ensemble averages (by far less harder to calculate). More specifically, let us consider an isolated N -body system (microcanonical ensemble) for which we fix the total energy E (for some systems, other constants of motion need to be fixed as well). If we assume an initial condition for the system and let it evolve (following Newtonian mechanics if the system is a classical one), after sufficiently long time t (mathematically speaking, in the limit $t \rightarrow \infty$; practically speaking, for times well above the inverse maximum Lyapunov exponent, whenever it is positive) the distribution of energies is given by an expression which is the microcanonical equilibrium distribution. From this distribution, we can in principle calculate the marginal distribution associated to any one among the N particles. This equilibrium distribution typically is, as first perceived by Gibbs, a power-law (see, for instance, [7,12]). For fixed value of E properly scaled with N and $N \gg 1$, this distribution approaches the celebrated Boltzmann exponential factor, which corresponds in fact to the canonical ensemble, i.e., when the system is in thermal contact with a thermostat (which fixes T , of course). A beautiful illustration is explicitly worked out in [7] for the marginal distribution corresponding to the momentum of one particle: in the $N \gg 1$ limit, the celebrated Maxwellian velocity distribution is neatly recovered. By the way, in 1993 Plastino and Plastino remarked [12] that the microcanonical equilibrium power-law distribution just mentioned precisely is the one which emerges from nonextensive statistical mechanics where $1 - q$ plays essentially the role of $1/N$.

What we have just described corresponds clearly to the $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} p(E; N; t)$ ordering. What relevant modifications are expected to happen in the $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty} p(E; N; t)$ ordering? Typically none if no long-range interactions are involved. But, as mentioned above, the situation is expected to be much more subtle in the presence of long-range interactions. To be more precise, consistently with the available numerical results, we generically expect (as conjectured in Fig 4 of VLL reference [31], and exhibited in [7{10], among others) that, after some transient towards equilibration starting from certain classes of initial conditions (such as waterbag or double waterbag in velocities, for instance), the system achieves a quasi-stationary or meta-equilibrium state whose duration diverges with N . After this state, for finite N , the system typically relaxes to the Boltzmann-Gibbs equilibrium (i.e., $q = 1$). Let us stress at this point that, if $N \sim 10^{23}$, during times which could be longer than the age of the universe, it is the anomalous, non Boltzmann-Gibbsian, meta-equilibrium state which will be observed and not the standard one (focused

in the VLL paper). In other words, between the standard and such nonstandard macroscopic states, something analogous to an activation barrier exists, whose height diverges with N . A typical (very illustrative but by no means unique) such system is the classical dimensional model of localized planar rotators with two-body (attractive) interactions which decay as r^{-d} , and whose coupling constant does not (unphysically) depend on N . The following anomalies have been (either analytically or numerically) observed in the microcanonical molecular dynamics approach:

(i) Above a rescaled critical energy $u_c = U_c/(N \cdot N')$ where U denotes the fixed total energy ($u_c = 0.75$ for $d = 0$), the system exhibits a (conveniently scaled) Lyapunov spectrum whose largest Lyapunov exponent approaches, for increasingly large N , a finite positive value for $d > 1$, but vanishes instead for $0 \leq d \leq 1$ (it vanishes, in fact, as N^{-1} , where $(-d)$ seems to be an universal function, independent of both d and u , which decreases from $1/3$ to zero when $-d$ increases from zero to unity, see VLL reference [35] and present reference [13]);

(ii) Below u_c , long standing metaequilibrium states are observed everytime the simulation is started from single (or even double) waterbag distributions in velocities. Specifically, the time evolution of the (properly size-scaled) average kinetic energy (also averaged over many statistically equivalent sets of initial conditions), exhibits [10] plateaux corresponding to temperatures appreciably different from the one associated with the canonical Boltzmann-Gibbs distribution. The duration of these plateaux seems to diverge with N , while their associated height remains finite. This result carries a most important corollary: it strongly suggests that the zeroth principle of thermodynamics can be valid out from the usual ($q = 1$) statistical mechanics [14];

(iii) Consistently with point (ii), the one-particle distribution of momenta is not the Maxwellian one during the entire extent of the plateaux just mentioned. It does not change for all times included in the plateaux, it is more peaked than the Maxwellian one for low momenta (consistently with a nonextensive statistical mechanics canonical distribution for $q > 1$), and exhibits a cut-off for high momenta, as generically expected for any microcanonical distribution (indeed, since the total energy is finite, even if all the energy was momentarily concentrated in the kinetic energy of a single particle, the corresponding momentum could not be infinite);

(iv) Consistently with points (ii) and (iii), a Levy-type anomalous superdiffusion is observed during the plateaux, which makes a crossover to normal diffusion when the plateau ends, and the usual Boltzmann-Gibbs equilibrium is attained.

It is very clear that no reason at all exists for expecting the above list to be exhaustive in describing the anomalies associated with the meta-stable states. It has been given just as an illustration of the important features that the VLL paper has overlooked. It is in this sense

that we have argued in the beginning of this Comment that the VLL paper does not prove, contrary to what is therein stated, that Boltzmann-Gibbs is sufficient, it only provides one more illustration that it is necessary, as probably all mechanical statistical physicists of the world are convinced. Do we mean by this that we have herein proved that, for classical conservative many-body long-range interacting Hamiltonians, nonextensive statistical mechanics is also necessary? Certainly not! What we believe that we have shown is that another, nonstandard formalism is also needed: at the present stage, for the particular system we are focusing on, nonextensive statistical mechanics is but a candidate (perhaps the correct one), on which many scientists are currently working. In a kind of desperate last effort, one could easily imagine a devil's advocate arguing that Boltzmann-Gibbs statistical mechanics is the uniquely correct one for describing the ultimate equilibrium macroscopic state. On this we have no objection at all, it even strongly seems that this is a strictly correct statement. However, for long-range interacting macroscopic systems initially placed in a nonstandard basin of attraction (in the space of the distributions) of the initial conditions, this state might be achieved "after the end of the universe". This feature would make its physical utility a very controversial matter, of a byzantine style which is not much of our taste. Along these lines, the only thing that really matters is whether, yes or no, a neat connection can be established between the power-law distributions which extremize the nonextensive entropic form $S_q = -k \sum_i p_i^q = -k \ln_q[1]$ [4] and the long standing metaequilibrium states existing before the ultimate, Boltzmann-Gibbs equilibrium state is attained. To achieve this goal, a tractable way is to perform microcanonical molecular dynamics simulations in long-range-interacting systems which contain $N \gg 1$ particles, and then focusing on subsystems of them with $M \gg 1$ particles such that $N \gg M \gg 1$. In the $(M; N=M) \rightarrow (1; 1)$ limit the comparison is expected to make sense. Work along this possibility is in progress.

It is certainly relevant to remind at this point that the celebrated Boltzmann-Gibbs exponential distribution for Hamiltonian systems has been founded in the literature in at least four manners, namely the variational entropic principle [15], the steepest descent method [16], the laws of large numbers [17] and the microcanonical counting [18]. All four have been generalized now (respectively [4], [19], [20] and [21]), and systematically lead to the power-law distributions emerging within nonextensive statistical mechanics.

Although perhaps not totally transparent, the VLL paper and consistently the present Comment focus essentially on classical Hamiltonian many-body systems. However, the concepts involved in nonextensive statistical mechanics have been successfully applied to many other systems. These include fully developed turbulence [22,23] (Beck has recently developed a quite impressive theory with not a single free parameter, which compares remarkably well with the available experimental data), granu-

lar matter [24], electron-positron annihilation and other high energy systems [25], Levy and correlated types of anomalous diffusion [26], low-dimensional nonlinear dynamical systems [27], self-organized criticality [28], reassociation in folded proteins [29], quantum entanglement [30], to quote but a few. For these systems, it is allowed to think that the question which remains to be fully clarified is not whether the formalism works successfully, but why it does so. Given the wide diversity of the systems under focus, it is not yet totally clear what are the basic ingredients of the game. It is however already acquired that some type of (multi)fractality or hierarchical structure is apparently always present (the physical mechanisms capable of driving such fractality appear to be long-range interactions, strongly nonmarkovian processes, fractal boundary conditions, quantum nonlocality, mesoscopic dissipation, and others). This fact tends to generate a slow, power-law, mixing in phase and analogous spaces, as opposed to the usual, exponential mixing. The situation might well be, in some cases, more subtle than just this (for instance, in the system of long-range interacting rotators addressed above, the Lyapunov exponents in the $N \rightarrow 1$ limit vanish for $u > u_c$ but are finite for $u < u_c$; see [31]), but it is presently unavoidable to think that microscopic mixing issues lay at the heart of nonextensive statistical mechanics and thermodynamics. This possibility goes in fact very well along ideas of Krylov [32], Balescu [33], Dorfman [34] and others. Further studies are certainly needed and welcome.

It is with pleasure that I acknowledge useful remarks by D.H. Zanette, S. Abe and A.K. Rajagopal.

[1] B.P. Volnyayr-Lee and E. Luijten, cond-mat/0009031.

[2] In VLL Refs. [7-12,29,31,32,35] and elsewhere, d is noted (both cases $0 \leq d \leq 1$ and $d > 1$ are considered in those references; VLL focus on $0 \leq d \leq 1$). In [7] is noted (in [7], only $d > 0$ is considered). What VLL refer to as "nonintegrable" ("integrable") interactions is referred to as "long-range" ("short-range") interactions in VLL Refs. [7-12,29,31,32,35] and elsewhere, and corresponds to $0 \leq d \leq 1$ ($d > 1$).

[3] G.Nicolis and D.Daems, Chaos 8, 311 (1998).

[4] C.Tsallis, J. Stat. Phys. 52, 479 (1988); E.M.F. Curado and C.Tsallis, J. Phys. A 24, L69 (1991) [Corrigenda: 24, 3187 (1991) and 25, 1019 (1992)]; C.Tsallis, R.S.Mendes and A.R.Plastino, Physica A 261, 534 (1998); the bibliography of the subject is regularly updated at <http://tsallis.cat.cbpf.br/biblio.htm>; for recent reviews see C.Tsallis, in Nonextensive Statistical Mechanics and Thermodynamics, eds. S.R.A. Salinas and C.Tsallis, Braz. J. Phys. 29, 1 (1999) [accessible at <http://sbif.usp.br/WWW/pages/Journals/BJP/Vol29/Num1/index.htm>], C.Tsallis, Entropic nonextensivity: A possible measure of complexity, to appear in the Proc. of

- the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Solitons and Fractals* (2001) [Santa Fe Institute Working Paper 00-08-043 (2000); cond-mat/0010150], and C. Tsallis, in *Nonextensive Statistical Mechanics and its Applications*, eds. S. Abe and Y. Okamoto, *Series Lecture Notes in Physics* (Springer-Verlag, Berlin, 2000), in press.
- [5] The word integrable is used, in the present context, when d^3r^d / r^3 is finite. It is at this point worthy reminding that the expressions "short" and "long" range interactions are used in different senses in different chemical and physical areas. VLL address this fact stating that "a pervasive notational problem in the nonextensive thermodynamics literature is the use of "long-range interactions" to mean "nonintegrable interactions"; also, VLL use expressions such as "considerably more important" class of integrable interactions and "true" long-range interactions to communicate to the readers their preferences. They mention the VLL Refs. [1,27,28], to which one could add, along the same vein, M.E. Fisher and V.A. Privman, *Comm. Math. Phys.* 103, 527 (1986), among others. In the present context, we prefer to use instead the notation commonly adopted in the nonextensive thermodynamics literature, which is consistent with L. Tisza, *Annals Phys.* 13, 1 (1961) [or in *Generalized thermodynamics*, (MIT Press, Cambridge, 1966), p. 123]. [In his words: "The situation is different for the additivity postulate P_{a2}, the validity of which cannot be inferred from general principles. We have to require that the interaction energy between thermodynamic systems be negligible. This assumption is closely related to the homogeneity postulate P_{d1}. From the molecular point of view, additivity and homogeneity can be expected to be reasonable approximations for systems containing many particles, provided that the intramolecular forces have a short range character."] and with P.T. Landsberg, *Thermodynamics and Statistical Mechanics*, (Oxford University Press, Oxford, 1978; also Dover, 1990), page 102 [In his words: "The presence of long-range forces causes important amendments to thermodynamics, some of which are not fully investigated as yet."]. Ludwig Boltzmann himself must have had perceptive intuitions along related lines. Indeed, in the first page of the second part of his *Vorlesungen über Gastheorie*, he qualifies the concept of ideal gas by writing: "When the distance at which two gas molecules interact with each other noticeably is vanishingly small relative to the average distance between a molecule and its nearest neighbor (or, as one can also say, when the space occupied by the molecules (or their spheres of action) is negligible compared to the space filled by the gas { ...".
- [6] M.E. Fisher, *Arch. Rat. Mech. Anal.* 17, 377 (1964); *J. Chem. Phys.* 42, 3852 (1965); *J. Math. Phys.* 6, 1643 (1965).
- [7] Lj. Milanovic, H.A. Posch and W. Thirring, *Phys. Rev. E* 57, 2763 (1998).
- [8] M. Antoni and A. Torcini, *Phys. Rev. E* 57, R6233 (1998).
- [9] V. Latora, A. Rapisarda and S. Ruo, *Phys. Rev. Lett.* 80, 692 (1998); *Physica D* 131, 38 (1999); *Phys. Rev. Lett.* 83, 2104 (1999); *Physica A* 280, 81 (2000); M. Antoni, S. Ruo and A. Torcini, *Proc. of the Workshop "The Chaotic Universe"* (Rome-Pescara, February 1999) (World Scientific), in press [cond-mat/9908336].
- [10] V. Latora and A. Rapisarda, *Dynamical quasi-stationary states in a system with long-range forces*, to appear in the *Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics"* (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Solitons and Fractals* (2001) [cond-mat/0006112].
- [11] H. Koyama and T. Konishi, cond-mat/0008208 and cond-mat/0008507.
- [12] A.R. Plastino and A. Plastino, *Phys. Lett. A* 193, 140 (1994).
- [13] A. Campa, A. Giansanti, D. Moroni and C. Tsallis, *Long-range interacting classical systems: universality in mixing weakening*, cond-mat/0007104.
- [14] V. Latora, A. Rapisarda and C. Tsallis, in progress.
- [15] J.W. Gibbs, *Elementary Principles in Statistical Mechanics* (Yale University Press, New Haven, 1902).
- [16] C.G. Darwin and R.H. Fowler, *Phil. Mag. and J. Sci.* 44, 450 (1922); R.H. Fowler, *Phil. Mag. and J. Sci.* 45, 497 (1923).
- [17] A.I. Khinchin, *Mathematical Foundations of Statistical Mechanics* (Dover, New York, 1949).
- [18] R. Balian and N.L. Balazs, *Ann. Phys. (NY)* 179, 97 (1987).
- [19] S. Abe and A.K. Rajagopal, *Microcanonical foundation for systems with power-law distributions*, *J. Phys. A* (2000), in press [cond-mat/0002159].
- [20] S. Abe and A.K. Rajagopal, *Justification of power-law canonical distribution based on generalized central limit theorem*, *Europhys. Lett.* (2000), in press [cond-mat/0003380].
- [21] S. Abe and A.K. Rajagopal, *Phys. Lett. A* 272, 341 (2000).
- [22] C. Beck, *Physica A* 277, 115 (2000); C. Beck, *Non-extensive statistical mechanics approach to fully developed hydrodynamic turbulence*, to appear in the *Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics"* (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Solitons and Fractals* (2001) [cond-mat/0005408].
- [23] T. Arimitsu and N. Arimitsu, *Phys. Rev. E* 61, 3237 (2000); *J. Phys. A* 33, L235 (2000); Tsallis statistics and turbulence, to appear in the *Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics"* (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Solitons and Fractals* (2001).
- [24] Y.-H. Taguchi and H. Takayasu, *Europhys. Lett.* 30, 499 (1995); see also A. Kudrolli and J. Henry, *Phys. Rev. E* 62, R1489 (2000).
- [25] I. Bediaga, E.M.F. Curado and J. Miranda, *Physica A* 286, 156 (2000); C. Beck, *Physica A* 286, 164 (2000); D.B. Walton and J. Rafelski, *Phys. Rev. Lett.* 84, 31

- (2000); G. W. Iłk and Z. W. Łodarczyk, *Phys. Rev. Lett.* **84**, 2770 (2000); G. W. Iłk and Z. W. Łodarczyk, The imprints of nonextensive statistical mechanics in high energy collisions, to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. G. Rigolini, C. T. Sallis and B. J. West, *Chaos, Solitons and Fractals* (2001) [hep-ph/0004250]; F. S. Navarra, O. V. Utyuzh, G. W. Iłk and Z. W. Łodarczyk, Violation of the Feynman scaling law as a manifestation of nonextensivity, to appear in N. Cim ento (2000) [hep-ph/0009165]; D. B. Ion and M. L. D. Ion, *Phys. Rev. Lett.* **81**, 5714 (1998); M. L. D. Ion and D. B. Ion, *Phys. Rev. Lett.* **83**, 463 (1999); D. B. Ion and M. L. D. Ion, *Phys. Rev. E* **60**, 5261 (1999); D. B. Ion and M. L. D. Ion, Optimality, entropy and complexity for nonextensive quantum scattering, to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. G. Rigolini, C. T. Sallis and B. J. West, *Chaos, Solitons and Fractals* (2001); M. L. D. Ion and D. B. Ion, Strong evidences for correlated nonextensive quantum statistics in hadronic scatterings, *Phys. Lett. B* **482**, 57 (2000); G. Kaniadakis, A. Lavagno and P. Quarati, *Phys. Lett. B* **369**, 308 (1996); P. Quarati, A. Carbone, G. Gervino, G. Kaniadakis, A. Lavagno and E. Mirakli, *Nucl. Phys. A* **621**, 345c (1997); G. Kaniadakis, A. Lavagno and P. Quarati, *Astrophysics and space science* **258**, 145 (1998); A. Lavagno, Proc. of Baryons 98 (Bonn, 22-26 September 1998), eds. D. W. Menze and B. Metsch (World Scientific, Singapore, 1999), page 709; M. Corradu, G. Kaniadakis, A. Lavagno, M. Lissia, G. Mezzorani and P. Quarti, in *Nonextensive Statistical Mechanics and Thermodynamics*, eds. S. R. A. Salinas and C. T. Sallis, *Braz. J. Phys.* **29**, 153 (1999); W. M. A. Iberico, A. Lavagno and P. Quarati, *Eur. Phys. J. C* **12**, 499 (1999); A. Lavagno and P. Quarati, *Nucl. Phys. B, Proc. Suppl.* **87**, 209 (2000); A. Lavagno and P. Quarati, *Classical and quantum non-extensive statistics effects in nuclear many-body problems*, to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. G. Rigolini, C. T. Sallis and B. J. West, *Chaos, Solitons and Fractals* (2001).
- [26] D. H. Zanette and P. A. Alemany, *Phys. Rev. Lett.* **75**, 366 (1995); C. T. Sallis, S. V. F. Levy, A. M. C. de Souza and R. M. Aynard, *Phys. Rev. Lett.* **75**, 3589 (1995) [Erratum: **77**, 5442 (1996)]; M. Buiatti, P. G. Rigolini and A. Montagnini, *Phys. Rev. Lett.* **82**, 3383 (1999); D. Prato and C. T. Sallis, *Phys. Rev. E* **60**, 2398 (1999); C. Budde, D. Prato and M. Re, preprint (2000) [cond-mat/0007038]; A. Robledo, *Phys. Rev. Lett.* **83**, 2289 (1999); A. Robledo, *J. Stat. Phys.* **100**, 475 (2000); A. R. Plastino and A. Plastino, *Physica A* **222**, 347 (1995); C. T. Sallis and D. J. Bukman, *Phys. Rev. E* **54**, R2197 (1996).
- [27] C. T. Sallis, A. R. Plastino and W.-M. Zheng, *Chaos, Solitons and Fractals* **8**, 885 (1997); U. M. S. Costa, M. L. Lyra, A. R. Plastino and C. T. Sallis, *Phys. Rev. E* **56**, 245 (1997); M. L. Lyra and C. T. Sallis, *Phys. Rev. Lett.* **80**, 53 (1998); M. L. Lyra, *Ann. Rev. Com. Phys.*, ed. D. Staufer (World Scientific, Singapore, 1998), page 31; U. Timakli, C. T. Sallis and M. L. Lyra, *Eur. Phys. J. B* **10**, 309 (1999); U. Timakli, *Symmetric unimodal maps: Some results from q-generalized bit cumulants*, *Phys. Rev. E* (1 Dec 2000), in press [cond-mat/9911420]; M. Buiatti, P. G. Rigolini and L. Palatella, *Physica A* **268**, 214 (1999); C. R. da Silva, H. R. da Cruz and M. L. Lyra, *Low-dimensional non-linear dynamical systems and generalized entropy*, in *Nonextensive Statistical Mechanics and Thermodynamics*, eds. S. R. A. Salinas and C. T. Sallis, *Braz. J. Phys.* **29**, 144 (1999); R. S. Johal and R. Rai, *Physica A* **282**, 525 (2000); V. Latora, M. Baranger, A. Rapisarda and C. T. Sallis, *Phys. Lett. A* **273**, 97 (2000); U. Timakli, G. F. J. A. Nanos and C. T. Sallis, Generalization of the Kolmogorov-Sinai entropy: Logistic- and periodic-like dissipative maps at the chaos threshold, preprint (2000) [cond-mat/0005210]; J. Yang and P. G. Rigolini, *Phys. Lett. A* **263**, 323 (1999); S. Montanero, L. Fronzoni and P. G. Rigolini, The non-extensive version of the Kolmogorov-Sinai entropy at work, *cond-mat/9911412*; F. A. B. F. de Moura, U. Timakli and M. L. Lyra, Convergence to the critical attractor of dissipative maps: Log-periodic oscillations, fractality and nonextensivity, *Phys. Rev. E* **62** (1 November 2000), in press [cond-mat/0008130]; M. Baranger, V. Latora and A. Rapisarda, Time evolution of thermodynamic entropy for conservative and dissipative maps, to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. G. Rigolini, C. T. Sallis and B. J. West, *Chaos, Solitons and Fractals* (2001) [cond-mat/0007302].
- [28] F. A. Tamarit, S. A. Cannas and C. T. Sallis, *Eur. Phys. J. B* **1**, 545 (1998); A. R. R. Papa and C. T. Sallis, *Phys. Rev. E* **57**, 3923 (1998); P. M. G. Leiser, F. A. Tamarit and S. A. Cannas, *Physica A* **275**, 272 (2000).
- [29] C. T. Sallis, G. Bemski and R. S. Mendes, *Phys. Lett. A* **257**, 93 (1999).
- [30] S. Abe and A. K. Rajagopal, *Physica A* **289**, 157 (2001) [quant-ph/0001085]; C. T. Sallis, S. Lloyd and M. Baranger, Generalization of the Peres criterion for local realism through nonextensive entropy, *quant-ph/0007112*.
- [31] To illustrate the kind of subtleties that might be involved, let us consider a simple example. The solution of the linear ordinary differential equation $\dot{x} = -\lambda x$ (with $x(0) = 1$) is $x = e^{-\lambda t}$, where λ represents the sensitivity to the initial conditions and λ is assumed to be the Lyapunov exponent of some simple, one-dimensional, nonlinear dynamical system. If $\lambda = 0$, the appropriate equation might be the nonlinear one $\dot{x} = -q x^q$ ($q \geq 2$), whose solution is now a power-law, namely $x = [1 + (1-q)x_0]^{1/(1-q)}$. This solution recovers the previous one as the $q = 1$ instance. Let us now consider the more general case where both linear and nonlinear terms are present. The corresponding differential equation can be written as $\dot{x} = -\lambda x + (-q x^{q-1}) x^q$. The solution is now given by $x = [1 - \frac{\lambda}{1-q} + \frac{\lambda}{1-q} e^{(1-q)\lambda t}]^{1/(1-q)}$. The pure linear case is recovered in the $q = 1$ (8q) case as well as in the $q = 1$ cases. The pure nonlinear case is recovered in the $\lambda = 0$ particular case. An interesting situation

appears when $q < 1$ and $q_1 > 0$. In this case, for $t \ll 1/(1 - q_1)$ the solution practically coincides with the power-law one, and for $t \gg 1/(1 - q_1)$ the solution asymptotically reproduces the exponential behavior. Generally speaking, it is possible to imagine that some specific effects could depend basically only on the nonlinear term $(\dots)$ and practically not on the linear term $(\dots)$. If so, the relevant part of (t) would be basically related to the power-law behavior, whether q_1 is zero or not. Considerations of this sort might have relevance for deeply understanding what happens dynamically in macroscopic systems like the long-range interacting rotators one discussed in this Comment, where a value u_c exists such as at both sides of which unusual features are observed: vanishing Lyapunov exponents for $u > u_c$, and the existence of long standing metastable states coexisting with nonzero Lyapunov exponents for $u < u_c$. A situation analogous to this one does occur in standard critical phenomena. Indeed, a crucially important quantity (directly related to the correlation length, the temperature dependance of which defines in turn the critical exponent) is the two-body space correlation, not of the values of the local order parameter at different sites, but of the fluctuations, with regard to the average order parameter, of the values of the local order parameter at different sites. The distinction is inexistent for the disordered phase (typically at $T > T_c$), where the average order parameter vanishes, but it is most important for the ordered phase (typically at $T < T_c$), where the average order parameter is nonzero.

- [32] N. Krylov, Nature 153, 709 (1944) [In his words: \ In the present investigation, the notion of ergodicity is ignored. I reject the ergodical hypothesis completely: it is both insufficient and unnecessary for statistics. I use, as starting point, the notion of motions of the mixing type,

and show that the essential mechanical condition for the applicability of statistics consists in the requirement that in the phase space of the system all the regions with a sufficiently large size should vary in the course of time in such a way that while their volume remains constant (according to Liouville's theorem) their parts should be distributed over the whole phase space (more exactly over the layer, corresponding to given values of the single-valued integrals of the motion) with a steadily increasing degree of uniformity. (...) The main condition of mixing, which ensures the fulfillment of this condition, is a sufficiently rapid divergence of the geodesic lines of this Riemann space (that is, of the paths of the system in the n -dimensional configuration space), namely, an exponential divergence (cf. Nopf). ". For full details on this pioneering approach see N. S. Krylov, Works on the Foundations of Statistical Physics, translated by A. B. Migdal, Ya. G. Sinai and Yu. L. Zeman, Princeton Series in Physics (Princeton University Press, Princeton, 1979).

- [33] R. Balescu, Equilibrium and Non-equilibrium Statistical Mechanics (John Wiley, New York, 1975), page 727 [In his words: \ It therefore appears from the present discussion that the mixing property of a mechanical system is much more important for the understanding of statistical mechanics than the mere ergodicity. (...) A detailed rigorous study of the way in which the concepts of mixing and the concept of large numbers of degrees of freedom influence the macroscopic laws of motion is still lacking.].
- [34] J. R. Dorfman, An Introduction to Chaos in Nonequilibrium Statistical Mechanics, Cambridge Lecture Notes in Physics 14, eds. P. G. Oddard and J. Yeomans (Cambridge University Press, Cambridge, 1999), footnote in page 9 [In his words: \ It is worth mentioning that there are examples of mixing systems with no non-zero Liapunov exponents. The concepts of ergodicity, mixing, and chaos can be quite subtle"].