

Anisotropies of the Hamiltonian and the Wave Function: Inversion Phenomena in Quantum Spin Chains

Kiyomi OKAMOTO^{*)}

Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551

(Received)

We investigate the inversion phenomenon between the XXZ anisotropies of the Hamiltonian and the wave function in quantum spin chains, mainly focusing on the $S = 1/2$ trimerized XXZ model with the next-nearest-neighbor interactions. We have obtained the ground-state phase diagram by use of the degenerate perturbation theory and the level spectroscopy analysis of the numerical data calculated by the Lanczös method. In some parameter regions, the spin-fluid is realized for the Ising-like anisotropy, and the Néel state for the XY -like anisotropy, against the ordinary situation.

§1. Introduction

In the ground state problem of the antiferromagnetic quantum spin chains with the XXZ interaction anisotropy $\Delta \equiv J_z/J_\perp$, the Ising-like anisotropy $\Delta > 1$ is usually favorable to the Néel state and the XY -like anisotropy $\Delta < 1$ to the spin-fluid (SF) state. This can be seen, for instance, in the $S = 1/2$ XXZ chain and the $S = 1$ XXZ chain, although the Haldane phase exists between the Néel and the SF phase for the latter case. Very recently we have found that the *inversion phenomenon* between the anisotropies of the interaction and the wave function.¹⁾ Namely, the SF state is realized for the Ising-like case and the Néel state for the XY -like case in the $S = 1/2$ distorted diamond spin chain²⁾ with the XXZ anisotropy. This inversion phenomenon is considered to be due to the interplay among the frustration, the trimer nature and the XXZ anisotropy of the Hamiltonian. In case of strong frustrations, the spin system tends to form singlet pairs to avoid the energy loss by competing interactions³⁾. But the formation of singlet pairs is incompatible with the trimer nature. Thus, in the XY -like anisotropy case, the spins turn to the z -direction to avoid the energy loss, because interaction of the z -direction is weaker than that of the xy -direction in the XY -like case. Similarly, we can explain the existence of the SF state for the Ising-like anisotropy case.

If this physical explanation is to the point, the inversion phenomenon will be observed in many quantum spin chain models having the frustration, the trimer nature and the XXZ anisotropy. From this point of view, we have investigated the $S = 1/2$ trimerized XXZ chain with the next-nearest-neighbor interactions, which may be the most fundamental model with these three characteristics.

^{*)} E-mail: kokamoto@stat.phys.titech.ac.jp

§2. $S = 1/2$ Trimerized XXZ Spin Chain with Next-Nearest-Neighbor Interactions

Our model is the $S = 1/2$ trimerized XXZ chain with the next-nearest-neighbor interactions (Fig. 1), having the Hamiltonian

$$H = J_1 \sum_i \{h_{3i-1,3i}(\Delta) + h_{3i,3i+1}(\Delta)\} + J_2 \sum_i h_{3i-2,3i-1}(\Delta) + J_3 \sum_j h_{j,j+2}(\Delta), \quad (2.1)$$

where

$$h_{m,n} \equiv S_m^x S_n^x + S_m^y S_n^y + \Delta S_m^z S_n^z. \quad (2.2)$$

The quantity J_1 (thick lines) denotes the intra-trimer nearest-neighbor coupling, J_2 (thin lines) the inter-trimer nearest-neighbor coupling and J_3 (broken lines) the next-nearest-neighbor coupling. All the couplings are assumed to be positive (antiferromagnetic). Hereafter we focus on the $J_1 \gg J_2, J_3$ case for simplicity, where the inversion phenomenon




Fig. 1. The $S = 1/2$ trimerized XXZ chain with the next-nearest-neighbor interactions.

was observed in the on $S = 1/2$ distorted diamond chain with XXZ anisotropy.

Let us discuss the ground state of our model by use of the degenerate perturbation theory. First we consider the 3-spin problem of \mathbf{S}_{3i-1} , \mathbf{S}_{3i} and \mathbf{S}_{3i+1} . As far as $J_1 \gg J_3$, the ground states of the 3-spin cluster are

$$|\uparrow_i\rangle \equiv \frac{1}{A} (|\uparrow\uparrow\downarrow\rangle - a|\uparrow\downarrow\uparrow\rangle + |\downarrow\uparrow\uparrow\rangle) \quad (2.3)$$

$$|\downarrow_i\rangle \equiv \frac{1}{A} (|\uparrow\uparrow\downarrow\rangle - a|\uparrow\downarrow\uparrow\rangle + |\downarrow\uparrow\uparrow\rangle) \quad (2.4)$$

where $|\uparrow\uparrow\downarrow\rangle$ means $|\uparrow_{3i-1}\uparrow_{3i}\downarrow_{3i+1}\rangle$ for instance, $A = \sqrt{2 + a^2}$ and

$$a = \frac{-3\tilde{J}_3 + \Delta + r}{2 + \tilde{J}_3^2 + \tilde{J}_3\Delta - J_3r}, \quad r = \sqrt{8 + (\tilde{J}_3 + \Delta)^2} \quad (2.5)$$

where $\tilde{J}_3 \equiv J_3/J_1$. As far as $J_1 \gg J_2, J_3$, we can restrict ourselves to these two states, $|\uparrow_j\rangle$ and $|\downarrow_j\rangle$, for the j th trimer, neglecting other 6 states. For convenience we consider $|\uparrow_j\rangle$ and $|\downarrow_j\rangle$ as the up-spin and down-spin states of the pseudo-spin \mathbf{T}_j , respectively. The interactions between the trimers are expressed as the interactions between pseudo-spins. For instance, we have

$$\frac{J_3}{2} (S_{3j}^+ S_{3j+2}^- + S_{3j}^- S_{3j+2}^+) \Rightarrow -\frac{2aJ_3}{(a^2 + 2)^2} (T_j^+ T_{j+1}^- + T_j^- T_{j+1}^+) \quad (2.6)$$

In the lowest order with respect to J_2 and J_3 , a straightforward calculation leads to

$$H_{\text{eff}} = \sum_j \left\{ J_{\text{eff}}^\perp (T_j^x T_{j+1}^x + T_j^y T_{j+1}^y) + J_{\text{eff}}^z T_j^z T_{j+1}^z \right\} \quad (2.7)$$

where

$$J_{\text{eff}}^{\perp} = \frac{a}{(a^2 + 2)^2} \{4aJ_2 - 8J_3\} \quad (2.8)$$

$$J_{\text{eff}}^z = \frac{a}{(a^2 + 2)^2} \left\{ a^2(a^2 - 2)J_2 - 2(a^2 - 2)^2J_3 \right\} \quad (2.9)$$

In the effective theory of the lowest order with respect to J_2/J_1 and J_3/J_3 , we can let $J_3 = 0$ in a , that is

$$a = \frac{\Delta + \sqrt{\Delta^2 + 8}}{2} \quad (2.10)$$

When $\Delta = 1$, we see $J_{\text{eff}}^{\perp} = J_{\text{eff}}^z = (4/9)(J_2 - J_3)$. Then the ground state of H_{eff} is either the ferromagnetic state or the spin-fluid state depending on whether $J_2 < J_3$ or $J_2 > J_3$. The ferromagnetic state of the \mathbf{T} -system corresponds to the ferrimagnetic state of the \mathbf{S} -system with the magnetization of $M_s/3$, where M_s is the saturation magnetization.

In general case ($\Delta \neq 1$), the ground state of H_{eff} is known from J_{eff}^{\perp} and J_{eff}^z as

$$\begin{aligned} J_{\text{eff}}^z < 0 \text{ and } |J_{\text{eff}}^z| > |J_{\text{eff}}^{\perp}| &\Rightarrow \text{ferromagnetic state} \\ |J_{\text{eff}}^z| < |J_{\text{eff}}^{\perp}| &\Rightarrow \text{spin-fluid state} \\ J_{\text{eff}}^z > |J_{\text{eff}}^{\perp}| &\Rightarrow \text{Néel state} \end{aligned} \quad (2.11)$$

which leads to

	XY-like	Ising-like	
$J_3 < c_1 J_2$	SF	Néel	(2.12)
$c_1 J_2 < J_3 < c_2 J_2$	Néel	SF	
$J_3 > c_2 J_2$	SF	ferro	

where

$$c_1 = \frac{a^2 + 2a + 2}{2a(a + 2)} \quad c_2 = \frac{1}{2} \frac{a(a + 2)(a^2 - 2a + 2)}{a^4 - 4a^2 + 8} \quad (2.13)$$

We note that

$$c_1 \rightarrow 5/8 \quad c_2 \rightarrow 1 \quad (\text{as } \Delta \rightarrow 1) \quad (2.14)$$

The phase diagrams for the $\Delta = 1, 0.5$ and 2.5 cases near the truncation point $J_2 = J_3 = 0$ are shown in Figs. 2-4, where $\tilde{J}_2 \equiv J_2/J_1$, $\tilde{J}_3 \equiv J_3/J_1$.

In the Ising limit $\Delta \rightarrow \infty$, we see $c_1 = c_2 = 1/2$, resulting in the vanishing of the spin-fluid state in Fig. 4. The remarkable nature of Fig. 3 is the existence of the Néel region, although the interaction anisotropy is XY-like. Similarly, the SF state is realized in Fig. 4 in spite of the Ising-like anisotropy. These are the *inversion phases*.

We have also performed the numerical diagonalization by use of the Lanczös method and analyzed the numerical data by the level spectroscopy method.^{4), 3), 6), 7)} For instance, when $\tilde{J}_2 = 0.01$ and $\Delta = 0.5$, the critical values of the SF-Néel transitions are $(\tilde{J}_3)_{\text{cr1}} = 0.006610$ and $(\tilde{J}_3)_{\text{cr2}} = 0.009696$, showing good agreements with the theoretical values from Eqs.(2.12) and (2.13) ($(\tilde{J}_3)_{\text{cr1}} = 0.006609$ and $(\tilde{J}_3)_{\text{cr2}} = 0.009703$, respectively). For the $\tilde{J}_2 = 0.01$ and $\Delta = 2.5$ case, we have obtained $(\tilde{J}_3)_{\text{cr1}} = 0.005623$ (Néel-SF) and $(\tilde{J}_3)_{\text{cr2}} = 0.006863$ (SF-ferrimagnetic), also having good agreements with the theoretical values ($(\tilde{J}_3)_{\text{cr1}} = 0.006609$ and $(\tilde{J}_3)_{\text{cr2}} = 0.006850$, respectively).




Fig. 2. Phase diagram for the $\Delta = 1$ case.




Fig. 3. Phase diagram for the $\Delta = 0.5$ case.




Fig. 4. Phase diagram for the $\Delta = 2.5$ case.

§3. Concluding Remarks

We have shown that there exist inversion regions in the $S = 1/2$ trimerized XXZ chain with next-nearest-neighbor interactions by use of the degenerate perturbation theory as well as the level spectroscopy analysis⁴⁾ of the numerical data obtained by the Lanczös method. This strongly suggests that the inversion phenomenon is popular to the $= 1/2$ XXZ models with the frustration and the trimer nature. In fact, this phenomenon has been also found in the $S = 1/2$ frustrated 3-leg ladder with the XXZ anisotropy.⁵⁾

In this paper we have restricted ourselves to the $J_1 \gg J_2, J_3$ case for simplicity. In case of larger J_3 , the dimer ground state is expected. In fact, when $J_2 = J_1$, our model is reduced to the uniform XXZ chain with next-nearest-neighbor interactions, which shows the SF-dimer or Néel-dimer transitions.^{3), 6), 7)} The full phase diagram of the present model will be reported elsewhere.

References

- 1) K. Okamoto and Y. Ichikawa, to be published in J. Phys. Chem. Solids, cond-mat/0108528.
- 2) K. Okamoto, T. Tonegawa, Y. Takahashi and M. Kaburagi: J. Phys.: Cond. Matter **11** (1999), 10485.
- 3) K. Okamoto and K. Nomura, Phys. Lett. **A169** (1992), 433 and refereces therein.
- 4) K. Okamoto, Prog. Theor. Phys. in this issue.
- 5) K. Okamoto and T. Sakai, in preparation.
- 6) K. Nomura and K. Okamoto, J. Phys. Soc. Jpn. **62** (1993), 1123.
- 7) K. Nomura and K. Okamoto, J. Phys. A: Math. Gen. **27** (1994), 5773.