

2D RANDOM DIRAC FERMIONS: LARGE N APPROACH

D. Serban

Service de Physique Theorique, CE Saclay, F-91191 Gif-sur-Yvette, France
 serban@spt.saclay.cea.fr

Abstract We study the symmetry classes for the random Dirac fermions in 2 dimensions. We consider N_f species of fermions, coupled by different types of disorder. We analyse the renormalisation group flow at the order of one loop. At N_f large, the disorder distribution flows to an isotropic distribution and the effective action is a sigma model.

1. Introduction

Random Dirac fermions in 2 dimensions appear in various contexts in condensed matter physics. They were used to model disorder in systems with degenerate Fermi surface [2], in the d-wave superconductors [3, 4] or to investigate the plateau transition in the integer quantum Hall effect [5]. In all these problems, the main question concerns the localization of quasiparticles, often different from the generic Anderson localization.

The universal properties of a system where randomness is present depend largely on the discrete symmetries, like time reversal invariance or spin rotation symmetry. This principle governs the Wigner-Dyson classification of random hermitian matrices [1]. Considering other discrete symmetries, like particle-hole or chiral symmetry, led Altland and Zimbauer [6] to extend this classification. However, it was recognized that the physics of 2d Dirac fermions is richer than expected from the random matrix classification, in the sense that for the same symmetry class, some systems may be critical and other not. One reason is that in two dimension, the sigma models, which are supposed to describe the universal physics, can support a topological term. In the form of a theta or WZW term, they can render the model critical [7, 8]. Sometimes, the presence of such a term can be deduced from general considerations [7, 8], in other situations it can be obtained exactly [3, 16]. A possible strategy, adopted in [14, 15] and applied here to all classes, is to try to

derive the effective sigma model, topological term included, by chain of controlled approximations. The drawback is that we have to consider a large number of fermion species N_f in order to have a control parameter for the approximation. This means that the most interesting cases, where N_f is small, are out of reach of this method.

The symmetry classes for 2d random Dirac Hamiltonians, were recently analysed [9]. The constraint that the discrete symmetries should also preserve the form of the Dirac operator leads to a refinement of the Altland-Zimbauer classification, in which the classes A III, C I and D III, appear each in two different forms. Here, we pursue the characterization of the symmetry classes for the Dirac Hamiltonians of [9]. We compute the beta function for all the disorder perturbations. When the number of fermion components N_f is large the flow of the coupling constants generically closes on a single coupling constant, a fact which simplifies considerably the derivation of the effective action. This was done for classes D in [14] and for classes C and C I in [15], and we refer to these papers for the details of the method. Without surprise, we obtain the sigma models defined on the manifolds identified by Zimbauer [6]. The two different versions of classes A III, C I and D III give rise to sigma models with or without a WZW term.

We consider random Dirac Hamiltonians in 2d, in the representation

$$H = \begin{pmatrix} V + M & 2i\partial_z + A_z \\ 2i\partial_z + A_z & V - M \end{pmatrix}; \quad (1)$$

where $\partial_z = (\partial_x - i\partial_y)/2$, $\partial_{\bar{z}} = (\partial_x + i\partial_y)/2$ and V, M, A_z and $A_{\bar{z}}$ are random matrix fields of dimension N_f .

The discrete symmetries preserving the form of the Dirac Hamiltonian are [9]

$$H = P H P^{-1}; \quad P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \quad P P^\dagger = 1; \quad P^2 = 1 \quad (2)$$

$$H = C H^T C^{-1}; \quad C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \quad C C^\dagger = 1; \quad C^T = -C \quad (3)$$

The first condition will be called chiral symmetry, while the second one could be called particle-hole symmetry when $C = 1$ and time reversal symmetry (or spin rotation invariance) when $C = +1$. The symmetry conditions impose constraints on the matrix fields V, M, A_z and $A_{\bar{z}}$, which are listed in [9]. We consider gaussian probability distribution for the disorder, compatible with the symmetry conditions.

The spectral properties of the Hamiltonians 1 can be analysed using the supersymmetry [10] (or replica) method, in which products of Green functions are represented by gaussian integral over equal numbers of

fermionic and bosonic Dirac fields. Average over gaussian randomness introduce effective coupling between the bosonic and fermionic fields, the coupling constants being the disorder variances.

The first step of our analysis is the renormalisation group flow for the effective coupling constants. The analysis can be eventually performed at all orders [11]. In the section 3 we give the one loop beta functions for all the classes and for any N_f . Generically, at large N_f the flow closes on a single coupling constant, corresponding to equal variances for different types of disorder.¹ In this case, the effective interaction, quartic in the Dirac fields, can be decoupled via a Hubbard-Stratonovich transformation using a (super)matrix field Q whose dimension does not depend on N_f . The number of components of the Dirac fields appear as an overall factor and it can be used as control parameter for a saddle point approximation.

The effective action for the field Q , hence the saddle point equation, has global invariance, described by a group G . The solutions of the saddle point equation break the invariance to a subgroup H , so that the manifold of saddle points is given by G/H . The low energy behaviour is obtained by taking slowly varying configurations of the field on G/H (Goldstone modes) and evaluating the action for these modes. The result is a sigma model on some symmetric (super) space. Given the fact that in two dimensions sigma models can support a topological term, we may obtain a theta term or a WZW one. In the last section, we give a list of bilinears in Dirac fields to which couple the field Q , the invariance group G of Q , as well as a diagonal solution of the saddle point equation q_0 together with its stabiliser H . Together with the coupling constants, these are the basic ingredients to characterize the sigma model for each symmetry class.

2. Disorder perturbations

We use notations similar to [14]. To obtain a compact notation, we introduce supermultiplets

$$= \begin{pmatrix} 0 \\ \mathbf{B} \\ \mathbf{C} \end{pmatrix} + \begin{pmatrix} 1 \\ \mathbf{T} \\ \mathbf{A} \end{pmatrix} ; \quad t = \left(\begin{matrix} ; & \mathbf{T} & ; & \mathbf{T} \\ & + & & + \end{matrix} \right) \quad (4)$$

and similarly for ψ , t , where \mathbf{B} and \mathbf{C} are fermion multiplets with N_f components while \mathbf{T} and \mathbf{A} are bosonic ghosts with the same number of components. The superscript T denotes transposition.

In the absence of disorder, the action for the Dirac fields is written

$$S_{\text{eff}} = \int \frac{d^2x}{2} (\bar{\psi} \partial_z \psi + \bar{\psi} \partial_{\bar{z}} \psi); \quad (5)$$

and it corresponds to a conformal field theory with central charge $c = 0$. From the holomorphic (antiholomorphic) fields $\psi(z)$, one can construct the current algebra $OSp(2N_f | 2N_f)_{k=1}$ [16].

The perturbation induced by disorder is

$$S_{\text{pert}} = \int g_i \frac{d^2x}{2} O_i; \quad (6)$$

where g_i are the disorder variances associated to different disorder invariants and the operators O_i are defined as follows

$$\begin{aligned} O_A &= \sum_I (\bar{\psi}_I \psi_I) (\bar{\psi}_I \psi_I); & O_a &= (\bar{\psi}_0 \psi_0) (\bar{\psi}_0 \psi_0); \\ O_V &= \sum_I (\bar{\psi}_I \psi_I)^2; & O_v &= (\bar{\psi}_0 \psi_0)^2; \\ O_M &= \sum_I (\bar{\psi}_I^+ \psi_I^+)^2; & O_m &= (\bar{\psi}_0^+ \psi_0^+)^2; \end{aligned} \quad (7)$$

where

$$E_I = \begin{pmatrix} e_I & 0 \\ 0 & e_I^T \end{pmatrix};$$

e_I are generators of $gl(N_f)$ normalized by $\text{Tr}(e_I e_J) = \delta_{IJ}$ and $A; V; M$ are subsets of indices of $sl(N_f)$ which depend on the symmetry class. e_0 is the identity normalized by $\text{Tr} e_0^2 = 1$.

3. The one loop beta functions

To study the renormalisation group flow for the coupling constants, we need the corresponding beta function. They can be derived from the operator product expansion of the perturbing operators O_i

$$O_i(z; z) O_j(0) \sim \frac{1}{z^2} C_{ij}^k O_k(0) + \text{reg}; \quad (8)$$

The one loop beta functions are given by [13]

$$\beta_k = -\frac{1}{2} C_{ij}^k g_i g_j \quad (9)$$

The results for different classes are listed in the remaining of this section.

The non-chiral classes

Class 0 (A)

This class do not posses neither chiral invariance, nor particle-hole/time reversal symmetry. We set $N_f = N$ and consider only disorder which do not break parity invariance of the action. The disorder elds A , V and M belong to $gl(N)$; the $sl(N)$ part generates the perturbation operators O_A ; O_V ; O_M , while the $gl(1)$ part generates O_a ; O_v ; O_m . The corresponding beta functions are

$$\begin{aligned}
 A &= \frac{D}{2} g_A^2 - \frac{D}{8} (g_M^2 + g_V^2) + \frac{B}{4} g_M g_V + \frac{1}{N} g_m g_V + \frac{1}{N} g_m g_v ; \\
 a &= \frac{N Y}{4} g_M g_V + \frac{1}{N} g_m g_v ; \\
 M &= x g_M^2 - \frac{1}{N} (g_m + g_v) g_M + 2x \left(\frac{D}{2} g_A g_M + \frac{B}{2} g_A g_V \right. \\
 &\quad \left. + \frac{2}{N} g_A g_v + \frac{2}{N} g_a (g_v + g_m) \right) - x g_M g_v ; \\
 V &= x g_V^2 + \frac{1}{N} (g_m + g_v) g_V + 2x \left(\frac{D}{2} g_A g_V + \frac{B}{2} g_A g_M \right. \\
 &\quad \left. + \frac{2}{N} g_A g_m + \frac{2}{N} g_a (g_M + g_v) \right) + x g_M g_v ; \\
 m &= \frac{1}{N} g_m (g_m + g_v) - C g_m (g_M + g_V) + 2C g_A g_m \\
 &\quad + \frac{2}{N} g_a (g_m + g_v) + \frac{N Y}{2} g_A g_v ; \\
 v &= \frac{1}{N} g_v (g_v + g_m) + C g_v (g_V + g_M) + 2C g_A g_v \\
 &\quad + \frac{2}{N} g_a (g_v + g_m) + \frac{N Y}{2} g_A g_M ;
 \end{aligned}$$

where the constants, with group theoretical significance, are

$$\begin{aligned}
 D &= 2N ; \quad C = \frac{N^2 - 1}{N} ; \quad B = \frac{2(N^2 - 4)}{N} ; \\
 x &= \frac{1}{N} ; \quad Y = \frac{4(N^2 - 1)}{N^2} ;
 \end{aligned}$$

In the large N limit, the line $g_A = g_V = g_M = g_m = g_v = 2g_a = g^{-1} = N$ is preserved by the RG flow. It is interesting to note that we cannot obtain an invariant line with all the coupling constants positive.

Classes 3 and 4 (D and C)

These classes have particle-hole symmetry ($\epsilon_c = -1$), enforced by symmetric, respectively antisymmetric matrix $T = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. We set $N_f = 2N$, but for class D, N can be half integer. The random elds A and V belong to the algebra $so(2N)$ (case D) and $sp(2N)$ (class C). These algebra are spanned by a subset of the $sl(2N)$ generators e_I, f_I , selected by the condition $[T, e_I] = -e_I$, $[T, f_I] = f_I$. The eld M belongs to the complement of these algebras with respect to $sl(2N)$, spanned by T_I with $[T, T_I] =$

$T_I^T = 1$. Operators O_A ; O_V are not allowed by these symmetry classes. The beta functions are

$$\begin{aligned}
 A &= \frac{D}{2} g_A^2 - \frac{D_{TT}}{8} g_M^2 - \frac{D}{8} g_V^2 + \frac{B_{TT}}{4} g_M g_V + \frac{1}{2N} g_m g_V ; \\
 M &= x_{TTT} g_M^2 - \frac{1}{2N} g_m g_M + 2x_{TT} \left(\frac{D_{TT}}{2} g_A g_M + \frac{B}{2} g_A g_V \right) \\
 &\quad + x_{TT} g_M g_V ; \\
 V &= x_{TT} g_V^2 + \frac{1}{2N} g_m g_V + 2x_{TT} \left(\frac{D}{2} g_A g_V + \frac{B_{TT}}{2} g_A g_M \right) \\
 &\quad + \frac{1}{N} g_m g_A + x_{TT} g_M g_V ; \\
 m &= \frac{1}{2N} g_m^2 + 2C g_A g_m + N y g_A g_V - C g_V g_m - C^0 g_m g_M ;
 \end{aligned}$$

with the group coefficients (again, the upper sign is for class D and the lower one for class C) :

$$\begin{aligned}
 C &= \frac{2N-1}{2} ; \quad C^0 = C_{sl(2N)} \quad C = \frac{(2N-1)(N-1)}{2N} ; \\
 D &= 2(N-1) ; \quad D_{TT} = 2N ; \quad D_{TTT} = \frac{2N}{2(N-1)} ; \\
 x &= x_{TT} = \frac{1}{2} ; \quad x_{TTT} = \frac{1}{2N} ; \quad x_{TTT} = \frac{1}{2N} ; \\
 y &= \frac{(2N-1)}{N} ; \quad B = 2(N-1) ; \\
 B_{TT} &= 2 \frac{N^2-1}{N} ; \quad B_{TTT} = 2 \frac{(N-2)(N-1)}{N} ;
 \end{aligned}$$

At large N , an invariant line is given by $g_A = g_M = g_V = g_m = g$.

Classes 3_+ and 4_+ (AII and AI)

These classes have time reversal (spin rotation) symmetry, with $\epsilon_c = 1$ and $\epsilon_T = 1$. They are similar to classes 3 and 4, with the random fields V and M changing roles: now M belongs to $so(2N)$ and $sp(2N)$ respectively, while V belongs to the complement with respect to $sl(2N)$. The beta functions can be obtained from that of the classes 3 and 4 by changing $g_M \leftrightarrow g_V$ and $g_m \leftrightarrow g_V$.

The chiral classes

The first type of chiral classes corresponds to $\epsilon_c = 1$ in eq. 2 and it comprises three classes. Only disorder of the vector potential type is allowed, and these classes can be seen as limits of the previous ones when $g_V, g_M, g_V, g_m \rightarrow 0$.

Class 1 (AIII)

The random gauge potential A belongs to $gl(N) \times gl(1) \times sl(N)$ ($N_f = N$).

$$A = \frac{D}{2} g_A^2 ; \quad a = 0 ; \quad (10)$$

with $D = 2N$.

Classes 5 and 6 (D III and C I)

In addition to the chiral symmetry, these classes possess particle-hole symmetry. We set $N_f = 2N$. The random gauge potential A belongs to $so(2N)$ and $sp(2N)$ respectively

$$A = \frac{D}{2} g_A^2 ; \quad (11)$$

with $D = 2(N-1)$.

The second type of chiral classes correspond to $\beta = 3$ and it comprises five classes. We set $N_f = 2N$, but for classes 8 and 9, N should be understood as even integer.

Class 2 (A III)

The chiral symmetry imposes the constraint $[A;] = 0$ to the random gauge potential; this condition selects an algebra $gl(N) \oplus gl(N)'$ ($gl(1) \oplus sl(N)$)² out of $gl(2N)$. The two $gl(1)$ parts are generated by the identity and σ_3 and we denote the associated coupling constants by g_a and g_a . We denote by e_i the generators of $sl(N) \oplus sl(N)$. The two $sl(N)$ components can have different variances, g_{A1} and g_{A2} , with $g_A = (g_{A1} - g_{A2})/2$.

The fields V and M are constrained by $fV; g = 0, fM; g = 0$, so they belong to $gl(2N) \cap (gl(N) \oplus gl(N))$, generated by e_{\pm} . Operators O_v and O_m are not allowed by the chiral classes.

The beta functions are

$$\begin{aligned} \beta_A^+ &= \frac{D_{ii}}{2} (g_A^+{}^2 + g_A^2) - \frac{D}{8} (g_M^2 + g_V^2) + \frac{B}{4} g_M g_V ; \\ \beta_A &= D_{ii} g_A^+ g_A ; \quad \beta_a = \frac{N Y}{2} g_M g_V ; \quad \beta_a = \frac{D}{8} (g_M^2 + g_V^2) ; \\ \beta_M &= x g_M (g_M + g_V) - \frac{D_i}{2} g_A^+ g_M + \frac{B_i}{2} g_A^+ g_V + \frac{1}{N} g_a (g_M + g_V) ; \\ \beta_V &= x g_V (g_V + g_M) - \frac{D_i}{2} g_A^+ g_V + \frac{B_i}{2} g_A^+ g_M + \frac{1}{N} g_a (g_V + g_M) ; \end{aligned}$$

with the group coefficients

$$\begin{aligned} D_{ii} &= D_{sl(N)} = 2N ; \quad D = D_{gl(2N)} - D_{gl(N)} = 2N ; \\ D_i &= D_{gl(2N)} = 2 + \frac{2}{N} = 2N + \frac{2}{N} ; \quad Y = 2 ; \\ x &= 0 ; \quad B_i = \frac{2(N^2 - 1)}{N} ; \quad B = 2N ; \end{aligned}$$

At large N , the line $g_A = 0, g_A^+ = g_M = g_V = 2g_a = 4g_a$ is left invariant by the RG flow.

Classes 7 and 8 (D III and C I)

These classes have both chiral symmetry and particle-hole symmetry, with the matrices enforcing these symmetries commuting.

We split the generators of $gl(2N)$ in four groups $i; \quad ; T_i; T$, with $i; T_i$ commuting with \quad and $\quad ; T$ anticommuting with $i;$ also, $i; = \quad^T_{i;}$ and $T_i; = \quad^T_{T_i;}$. The random gauge potential A is an element of $so(N)$ or $so(N)$ or $sp(N)$ or $sp(N)$ generated by $i;$. If we choose the representation $\quad = 1, V$ belongs to $o(2N)$ or $o(N)$ or $sp(2N)$ or $sp(N)$ or $sp(N)$ respectively, generated by \quad , while the field M is generated by T .

The beta functions are

$$\begin{aligned} A &= D^{ii} g_A^+ g_A ; \\ A^+ &= \frac{D^{ii}}{2} (g_A^{+2} + g_A^2) - \frac{D_{TT}}{8} g_M^2 - \frac{D}{8} g_V^2 + \frac{B_T}{4} g_M g_V ; \\ M &= x_{TTT} g_M^2 + 2x_T^{ii} \frac{D^i_T}{2} g_A^+ g_M + \frac{B^i}{2} g_A^+ g_V - x_{T-T} g_M g_V ; \\ V &= x \quad g_V^2 + 2x^{ii} \frac{D^i}{2} g_A^+ g_V + \frac{B^i_T}{2} g_A^+ g_M + x_{T-T} g_M g_V ; \end{aligned}$$

where the group coefficients are

$$\begin{aligned} D^{ii} &= N - 2 ; \quad D^i = D_{TT} = N ; \\ D^i_T &= D^i_T = N - 1 ; \\ x_{TTT} &= x \quad = x_{T-T} = x_{T-T} = \frac{1}{2} ; \quad x_T^{ii} = x^{ii} = 0 ; \\ B_T &= N ; \quad B^i = B^i_T = N - 1 ; \end{aligned}$$

In the large N limit the RG flow preserve the line $g_A = 0, g_A^+ = g_M = g_V$.
Classes 9_+ and 9_- (D I and C II)

These classes have also both particle-hole and chiral symmetry, with the matrices enforcing these constraints anticommuting, $f ; g = 0$.

If we choose $\quad = 1$, the matrix defining the particle-hole symmetry is symmetric for class D I and antisymmetric for class C II, as for the cases 3 and 4 or 7 and 8. The opposite choice exchanges the symmetry of the matrix between the two classes.

As in the previous case, we split the $gl(2N)$ generators into four classes. Since $f ; g = 0$, i spans now $gl(N)$, to which the field A belong. V is generated by \quad , which span $o(2N)$ or $gl(N)$ and $sp(2N)$ or $gl(N)$ respectively. M is again generated by T .

$$\begin{aligned} a &= \frac{D_{TT}}{8} g_M^2 - \frac{D}{8} g_V^2 ; \quad A = D^{ii} g_A^+ g_A ; \\ A^+ &= \frac{D^{ii}}{2} (g_A^{+2} + g_A^2) - \frac{D_{TT}}{8} g_M^2 - \frac{D}{8} g_V^2 + \frac{B_T}{4} g_M g_V ; \end{aligned}$$

$$\begin{aligned}
g_M &= x_{TTT} g_M^2 + 2x_{Ti}^i \frac{D_{TT}^i}{2} g_A^+ g_M + \frac{B_{TT}^i}{2} g_A^+ g_V - x_{T-T} g_M g_V ; \\
g_V &= x_{VVV} g_V^2 + 2x_{Vi}^i \frac{D_{VT}^i}{2} g_A^+ g_V + \frac{B_{VT}^i}{2} g_A^+ g_M + x_{T-T} g_M g_V ;
\end{aligned}$$

with

$$\begin{aligned}
D_{ii} &= N ; D_{TT} = N - 2 ; D_{TTT} = N - 2 ; \\
D_{Ti}^i &= \frac{(N-1)(N-2)}{N} ; D_{VT}^i = \frac{(N-1)(N-2)}{N} ; \\
x_{TTT} &= x_{VVV} = x_{T-T} = x_{T-T} = 0 ; x_{Ti}^i = \frac{1}{2N} ; \frac{1}{2} ; \\
x_{Vi}^i &= \frac{1}{2N} ; \frac{1}{2} ; B_{TT} = N ; B_{Ti}^i = N - 1 ; B_{VT}^i = N - 1 ;
\end{aligned}$$

At large N the RG flow preserves the line $g_A = 0$ and $g_A^+ = g_M = g_V = 4g_A$.

We note that the equations listed above have interesting properties under the transformation $g_M \leftrightarrow g_V$ and $g_M \leftrightarrow g_V$. This operation leaves the equations for classes 0, 2, 7 and 8 invariant and exchanges the ones for classes 3, 4 and 9, therefore it is equivalent to changing the sign of c , when defined². Moreover, on the line $g_M = g_V = g$ ($= g_M = g_V$, when they exist), the flow for g and the flow for g_A decouple. This was noted in [18] for class 0 and was related to "spin-charge" separation, for class C in [12].

4. Hubbard-Stratonovich transformation and the sigma model

We consider products of n Green functions, n_R retarded and n_A advanced, and we need to introduce a copy of the multiplets ψ for each Green function. When the Hamiltonian has chiral or particle-hole invariance, the spectrum is symmetric with respect to $E = 0$ and we do not need to distinguish between retarded and advanced sector. In this case, the density of states can be singular at the point $E = 0$.

As already mentioned, when the coupling constants have equal values, the decoupling of the quartic interaction is particularly simple. At that particular point, the disorder perturbation can be written only in terms of singlets of $gl(N_f)$. The decoupling matrix Q couples to these singlets and it has a size independent of N_f . Up to an additive constant, the effective action for Q is

$$S[Q] = -N_f \int \frac{1}{g} \frac{d^2x}{2} \text{STr} Q^2 - \text{STr} \ln \begin{vmatrix} Q & 0 \\ 0 & Q \end{vmatrix} ; \quad (12)$$

where $\mathfrak{g} = N_f \mathfrak{g}$. Part of the invariance of the free action $S_0 \in \mathcal{O}Sp(2n_f, 2n_f)$, survives in the effective action 12. We denote the residual invariance group by G .

The saddle point equation $S_Q \models Q = 0$ can be solved making a diagonal ansatz $Q_0 = q_0$, where $q_0^2 = 1$ and q_0 has the dimension of a mass. As there is no scale in the problem, q_0 being dimensionless, is dynamically generated. The group G acts on Q by conjugation, and from a particular saddle point solution q_0 one generates a whole manifold by $q_0 \mapsto q = g q_0 g^{-1}$. This manifold is isomorphic to G/H , where H is the stabilizer of q_0 . The last step in deriving the sigma model is to evaluate the action 12 on slowly varying configurations $q(x)$ on G/H . An elegant way of doing it is to use non-abelian bosonisation [17], as in [4, 14] and to write the free action as a WZW model. The field $q(x)$ becomes a mass term for the WZW field, which is forced to follow it. At scales larger than q_0^{-1} , the effective action is

$$S[q] = \frac{1}{16\pi} \int d^2x \text{STr} \partial_\mu q \partial^\mu q + S_{\text{top}}[q] \quad (13)$$

In the following, we give the main characteristics of the sigma model manifold for the 13 classes.

Class 0 (A)

$$Q = \text{Tr}_{\mathfrak{gl}(N)} [\begin{smallmatrix} 3 & \\ & t+ \\ & & t \end{smallmatrix}]; \quad q_0 = \begin{smallmatrix} 3 & \\ & \\ & & 1 \end{smallmatrix};$$

$$G = GL(n_f); \quad H = GL(n_R, n_R) \times GL(n_A, n_A);$$

with $\text{diag}(1_{n_R}; 1_{n_A})$.

Classes 3 and 4 (D and C)

$$Q = \text{Tr}_{\mathfrak{gl}(2N)} \begin{smallmatrix} t+ & & \\ & t & \\ & & 1 \end{smallmatrix}; \quad q_0 = \begin{smallmatrix} 3 & \\ & \\ & & 1 \end{smallmatrix};$$

$$G = \mathcal{O}Sp(2n_f, n_f); \quad H = GL(n_f);$$

with $\text{diag}(1;)$

Classes 3+ and 4+ (A II and A I)

$$Q = \text{Tr}_{\mathfrak{gl}(2N)} \begin{smallmatrix} 3 & & t+ & & \\ & t & & & \\ & & & t & \\ & & & & 1 \end{smallmatrix}; \quad q_0 = \begin{smallmatrix} 3 & \\ & \\ & & 1 \end{smallmatrix};$$

$$G = \mathcal{O}Sp(2n_f, n_f); \quad H = \mathcal{O}Sp(n_R, n_R) \times \mathcal{O}Sp(n_A, n_A);$$

The effective action for the classes 1, 5 and 6 is invariant under multiplication at left/right by holomorphic/antiholomorphic group elements. This is a sign that the theory is conformally invariant and indeed, when deriving the sigma model we obtain a WZW term.

Class 1 (A III)

$$Q = \text{Tr}_{\mathfrak{gl}(N)} [\begin{smallmatrix} 3 & \\ & t \end{smallmatrix}]; \quad Q = 1 = 2 \text{Tr}_{\mathfrak{gl}(N)} [\begin{smallmatrix} 3 & \\ & t \end{smallmatrix}];$$

$$q_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \quad G = GL(n, \mathbb{H})_L \times GL(n, \mathbb{H})_R; \quad H = GL(n, \mathbb{H});$$

The level of the WZW action is $k = N_f$.

Classes 5 and 6 (D III and C I)

$$Q = Tr_{gl(2N)} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \quad Q = Tr_{gl(2N)} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \\ q_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \quad G = OSp(n, \mathbb{H})_L \times OSp(n, \mathbb{H})_R; \quad H = OSp(n, \mathbb{H});$$

The level of the WZW action is $k = 2N$, the change of sign indicating that the compact and non-compact sector switch from the orthogonal to the symplectic sector from one class to another.

To take into account the chiral symmetry for the last two classes, we double the supermultiplets by

$$\begin{aligned} \psi &= \frac{1}{\sqrt{2}} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}; \quad \psi^\dagger = \frac{1}{\sqrt{2}} (\psi_1^\dagger \quad \psi_2^\dagger); \\ \psi &= \frac{1}{\sqrt{2}} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}; \quad \psi^\dagger = \frac{1}{\sqrt{2}} (\psi_1^\dagger \quad \psi_2^\dagger) \end{aligned}$$

Class 2 (A III)

$$Q = Tr_{gl(2N)} [\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}]; \quad q_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \\ G = GL(n, \mathbb{H}) \times GL(n, \mathbb{H}); \quad H = GL(n, \mathbb{H});$$

where $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ is the Pauli matrix acting in the extra space introduced to take into account the chiral symmetry. G is selected from $GL(2n, \mathbb{H})$ by the condition $[\mathfrak{g}; \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}] = 0$.

Classes 7 and 8 (D III and C I)

$$Q = Tr_{gl(2N)} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \quad q_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \\ G = OSp(2n, \mathbb{H}) \times OSp(2n, \mathbb{H}); \quad H = OSp(2n, \mathbb{H});$$

G is selected from $OSp(4n, \mathbb{H})$ by the condition $[\mathfrak{g}; \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}] = 0$

Classes 9₋ and 9₊ (D I and C II)

$$Q = Tr_{gl(2N)} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \quad q_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \\ G = GL(2n, \mathbb{H}); \quad H = OSp(2n, \mathbb{H});$$

The group G is selected by from $OSp(4n, \mathbb{H})$ by $[\mathfrak{g}; \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}] = 0$.

The classes 2, 7, 8 and 9 do not allow a theta term; also, there is no WZW term. Note that the sigma model is defined on the same manifold for the pair of classes 1 and 2 (class A III), 5 and 7 (class D III) and 6 and 8 (class C I); the difference between the two members of a pair consists only on the presence or absence of the WZW term.

The coupling constant for the sigma model is given by the inverse of the number of the Dirac fermions, $f = 1/N_f$.

Notes

1. Exceptions may be the coupling constants corresponding to some $u(1)$ part of the disorder, specific to classes 0, 1, 2 and 9. This situation was analysed in the case 1 (A III) in [4] and in [18] in a non-hermitian version of case 0. For the moment, we ignore this point and analyse the case of equal disorder variances.

2. We thank to D. Bernard for pointing out that this can be realized on the Dirac Hamiltonian via a non-unitary transformation.

References

- [1] F. Dyson, J. Math. Phys. 3 (1962) 140;
- [2] F. Fradkin, Phys. Rev. B 33 3257;
- [3] A. A. Nersisyan, A. M. Tsvelik and F. Wegner, Phys. Rev. Lett. 72 (1994) 2628;
- [4] A. Altland, B. D. Simons, M. R. Zimbauer, cond-mat/0006362;
- [5] A. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, Phys. Rev. B 50 (1994) 7526
- [6] A. Altland and M. Zimbauer, Phys. Rev. B 55 (1997) 1142; M. Zimbauer, J. Math. Phys. 37 (1996) 4986;
- [7] N. Read and D. Green, Phys. Rev. B 61 (2000) 10267; T. Senthil and M. P. A. Fisher, Phys. Rev. B 61
- [8] P. Fendley and R. M. Konik, Phys. Rev. B 62 (2000) 9359,
- [9] D. Bernard and A. LeClair, "A classification of 2d random Dirac fermions", cond-mat/0109552.
- [10] K. B. Efetov, Adv. Phys. 32 (1983) 53;
- [11] B. Gerganov, A. LeClair and M. Moriconi, hep-th/0011189;
- [12] D. Bernard and A. LeClair, Phys. Rev. B 64 (2001), 045306.
- [13] J. Cardy, in Les Houches, Eds. E. Brezin and J. Zinn-Justin, North-Holland, 1998;
- [14] M. Bocquet, D. Serban, M. R. Zimbauer, Nucl. Phys. B 578 (2000) 628;
- [15] D. Bernard, N. Regnault, D. Serban, Nucl. Phys. B 612 (2001) 291;
- [16] D. Bernard, "(Perturbed) conformal field theory applied to 2d disordered systems: an introduction", hep-th/9509137
- [17] E. Witten, Commun. Math. Phys. 92 (1984) 455;
- [18] S. Guruswamy, A. LeClair and A. W. Ludwig, Nucl. Phys. B 583 (2000) 475;