

Is the mean-field approximation so bad?

A simple generalization yielding realistic critical indices for 3D Ising-class systems

A.N. Rubtsov*

Physical Department of Moscow State University, Moscow 119899, Russia

(Dated: March 22, 2022)

Modification of the renormalization-group approach, invoking Stratonovich transformation at each step, is proposed to describe phase transitions in 3D Ising-class systems. The proposed method is closely related to the mean-field approximation. The low-order scheme works well for a wide thermal range, is consistent with a scaling hypothesis and predicts very reasonable values of critical indices.

PACS numbers: 05.70.Fh

It is well understood since 1960's that an essential increase of fluctuations accompanies a second-order phase transition. Exact results for 2D Ising model [1] showed that Landau phenomenology [2] was too crude to describe the vicinity of the transition point. Ginzburg demonstrated that it is correlations what breaks Landau approach down [3]. He established the parameter which determines the thermal range for the so-called fluctuation region. According to the scaling hypothesis [4], the properties of a system in the fluctuation region are determined by the single quantity r_c , being the correlation length for fluctuations. This parameter shows a power-law dependence on the external field h and on the thermal interval t from the transition: $r_c \propto h^{-\mu}$, $r_c \propto t^{-\nu}$. Values of μ and ν are determined by the universal characteristics, like the dimensionality D , the number of components of the order parameter, and the type of the interaction in the system (long-range or short-range). Other critical indices can be expressed via μ and ν . For instance for the scalar order parameter \bar{x} one obtains $\bar{x} \propto t^{-\beta}$, $\beta = \nu D - \nu/\mu$.

The general framework to obtain critical indices is the renormalization-group (RG) approach [5, 6, 7]. The RG transformation consists of the two stages. First, one integrates out all x_k with wave vectors $k > k_0/s$, where k_0 is a cut-off and $s > 1$ is a parameter. The main assumption of the method is that the potential energy of thus renormalized system can be truncated to certain simplified form, for example to the form of ϕ^4 model. The second step is a scale transformation. It is aimed to restore the cut-off for k back to k_0 and preserve the original dispersion law for the long-wavelength modes.

The RG transformation should be applied recursively; its unstable fixed point corresponds to a phase transition. Lyapunov exponents for the vicinity of the fixed point determine the critical indices. The scaling hypothesis requires exactly two of the Lyapunov exponents to be positive, they correspond to the two independent critical indices.

With respect to the value of s , two modifications of the RG scheme can be distinguished. Wilson's ε expansion [5, 6] deals with $s \gg 1$. The method is formally valid near the upper critical dimension D_{up} (for the Ising-class classical systems $D_{up} = 4$). The "most divergent" dia-

gram series can be pointed out in this case. This gives the asymptotic expansion in terms of $\varepsilon \equiv D_{up} - D$. In the continuous version of the scheme (so-called exact RG approach) [7], s is infinitesimally close to unity. Certain decoupling (truncation) should be introduced in this case to solve the integro-differential equation for RG flow. An accuracy is then determined by a form of the decoupling used.

All the approaches mentioned above are very well-developed. They allow to calculate critical indices and universal combinations of amplitudes with a high accuracy. Progress is also achieved in the calculation of non-universal quantities; particularly a crossover between Gaussian (Landau-like) and critical behaviour is analyzed [8]. However RG is applicable only for long-wavelength excitations since they can be described as a fluctuating field. Therefore, in a practical calculation, short-wavelength excitations should be first integrated out by means of some perturbation expansion, and then RG should be applied. The short- and the long-wavelength excitations should be separated *ad hoc*. Such a two-step procedure is not very elegant. At least from the methodological point of view it is desired to develop a scheme which gives all the information in a uniform manner.

It is interesting to note that calculations are much simpler outside the critical region. In fact, even the mean-field approximation (MFA) is usually sufficient for the qualitative analysis of this region. In MFA, the non-local interaction in the system is replaced by an effect of a self-consistent average field. The MFA predictions for the transition temperature and thermal behaviour of the order parameter in 3D case are quite reasonable [9]. On the other hand MFA gives Landau set of indices and is not very accurate. Note that Landau approach itself has a mean-field nature.

In this paper, we present a simple generalization of MFA, performing well for the 3D Ising-class systems both nearby and far from the transition. Particularly the critical behaviour is reproduced correctly: the values of indices are predicted with a few-percent accuracy. Essentially, the present approach consists in the alternating application of the Stratonovich transformation and the most primitive version of the renormalization-group

transformation at a finite value of s .

We consider a classical scalar field with an anharmonic local potential and a short-range harmonic interaction at temperature T . The partition function of this system is

$$\begin{aligned} Z &= \int [Dx] e^{-V/T}; \\ V &= \int U(x_r + \bar{x}) d\mathbf{r} + \sum_k \frac{\Omega_k |x_k|^2}{2}. \end{aligned} \quad (1)$$

Here $U(x + \bar{x})$ is an even on-site potential, x_k is a Fourier transform of the displacement x_r from the average position \bar{x} ; $\bar{x} \neq 0$ below the transition temperature, and the average of x_r equals zero by definition. To make the definitions unambiguous, we put additional conditions $U(\bar{x}) = 0$; $\Omega_{k=0} = 0$. The wave vector k does not exceed the unitary cut-off: $k^2 \leq 1$. The case of a short-range interaction corresponds to the asymptotic behaviour

$$\Omega_k = \omega k^2, k \rightarrow 0. \quad (2)$$

The simplest way to integrate out the modes is to skip

x_k with $k > 1/s$. This corresponds to the zero-th approximation in the nonlinearity of U . The subsequent scaling results in the RG transformation $k \rightarrow sk$, $U \rightarrow s^3 U$, $x \rightarrow x/\sqrt{s}$. The defect of thus applied RG method comes from the increase of correlations (non-linear part of U grows). Therefore, after several iterative transformations, the RG procedure is not valid any more. Although the formal analysis of the fixed point $U(x) = 0$ can be done, it wrongly gives Landau-like critical indices.

To avoid the problem higher-order approximations and the truncation of the potential are used in known schemes described above (a very clear explanation is given in [10]). Here we use another procedure: the zero-th approximation is used, but the change of variables is made at each RG step. The change of variables reduces correlations and is aimed to compensate its increase at RG transformation.

At the first stage, we integrate out (omit) all the modes with $\Omega_k > \bar{\Omega}$, where $\bar{\Omega} = \frac{3}{4\pi} \int \Omega_k d^3k$ is an average of Ω_k . For the rest of the modes, the Stratonovich identity is utilized:

$$\exp\left(-\frac{(\Omega_k - \bar{\Omega})|x_k|^2}{T}\right) = \frac{\bar{\Omega} - \Omega_k}{\pi \bar{\Omega}^2} \int \exp\left(-\frac{1}{T} \left(2\bar{\Omega} \text{Re}(x_k f_k^*) + \frac{\bar{\Omega} \Omega_k |f_k|^2}{\bar{\Omega} - \Omega_k}\right)\right) df_k. \quad (3)$$

Here the integration over the complex variable $f_k = \Re(f_k) + i\Im(f_k)$ denotes the integration over the complex plane: $\int df_k \equiv \int_{-\infty}^{\infty} d\Re(f_k) \int_{-\infty}^{\infty} d\Im(f_k)$.

After (3) is substituted into (1), expressions $\sum_k \text{Re}(x_k f_k^*) = \int x_r f_r d\mathbf{r}$ and $\sum_k \bar{\Omega} |x_k|^2 = \int \bar{\Omega} x_r^2$ can be collected in the exponent. Therefore

$$Z \propto \int [Dx] \int [Df] \exp\left(-\frac{1}{T} \left(\int \left(U(x_r + \bar{x}) - \Omega_k f_r x_r + \frac{\bar{\Omega} x_r^2}{2}\right) d\mathbf{r} + \sum_k \frac{\bar{\Omega} \Omega_k |f_k|^2}{2(\bar{\Omega} - \Omega_k)}\right)\right), \quad (4)$$

where f_r is the inverse Fourier transform of f_k .

Now, we can integrate over x_r . Let us introduce the function $F(f)$ accordingly to the equation:

$$\exp\left(-\frac{F(f + \bar{x}) - F_0}{T}\right) = \int \exp\left(-\frac{1}{T} \left(U(x + \bar{x}) + \frac{\bar{\Omega}(x - f)^2}{2}\right)\right) dx, \quad (5)$$

where F_0 is a constant, defined from the condition $F(\bar{x}) = 0$.

With thus defined F , the partition function takes the form

$$\begin{aligned} Z &\propto \int [Df] e^{-W/T}; \\ W &= \int F(f_r + \bar{x}) d\mathbf{r} + \sum_k \frac{\bar{\Omega} \Omega_k |f_k|^2}{2(\bar{\Omega} - \Omega_k)}. \end{aligned} \quad (6)$$

Equation (6) is formally very similar to Eq.(1). Function F is even; the dispersion at $k \rightarrow 0$ coincides that

of Eq.(1). One can also guess from Eq.(3) that as the average of x_r equals zero, the average of f_r also vanishes. Indeed, if this average would not be zero, $f_{k=0}$ would fluctuate around a macroscopically large average value, and (3) would not be fulfilled.

We argue here that variables f_k are much less correlated, than x_k . Let us consider the parabolic approximation for F : $F(\bar{x} + f) \approx \frac{\partial F(\bar{x})}{\partial f} f + \frac{\partial^2 F(\bar{x})}{2\partial f^2} f^2$. Variables f_k are uncorrelated in this approximation. Obviously $\langle f \rangle >$

vanishes only if $\frac{\partial F(\bar{x})}{\partial f} = 0$, that gives

$$\frac{\bar{\Omega}\bar{x}}{T} = \frac{\int x \exp(-T^{-1}(U(x+\bar{x}) + \frac{1}{2}\bar{\Omega}x^2)) dx}{\int \exp(-T^{-1}(U(x+\bar{x}) + \frac{1}{2}\bar{\Omega}x^2)) dx}. \quad (7)$$

This is exactly the mean-field equation of state for the system (1). So, even the complete neglect of correlations in f_k still gives something reasonable. On the other hand, the parabolic approximation for U directly in Eq.(1) gives non-sense: since $U(x)$ does not depend on temperature, the system does not show a transition at all in this approximation. Such a comparison convinces us that correlations in f_k are less essential, than in x_k . As it was pointed above we expect that this can compensate an increase of correlations in RG transformation.

Now, we finish the RG transformation, operating with Eq.(6). All f_k with $k > 1/s$ are to be integrated out (omitted in our approximation). After the scale transformation $k \rightarrow sk, f \rightarrow x/\sqrt{s}$ is done, the potential energy of the system takes the form

$$\int U^{(1)}(x_r + \bar{x}^{(1)})d\mathbf{r} + \sum_k \frac{\Omega_k^{(1)}|x_k|^2}{2}, \quad (8)$$

where

$$\Omega_k^{(1)} = \frac{s^2\bar{\Omega}\Omega_k}{\bar{\Omega} - \Omega_{k/s}}, \quad (9)$$

$$U^{(1)}(x + \bar{x}^{(1)}) = s^3 F(x/\sqrt{s} + \bar{x}), \quad \bar{x}^{(1)} = \bar{x}\sqrt{s}. \quad (10)$$

Such a transformation should be applied recursively, like in the standard RG approach. The analysis can be easily performed numerically.

The sequence $\Omega_k, \Omega_k^{(1)}, \Omega_k^{(2)}, \dots$ is independent of U and T . At every step a new $\bar{\Omega}^{(n)}$ are to be calculated. After several transformations $\bar{\Omega}^{(n)}$ and $\Omega_k^{(n)}$ converge to certain limit. Calculation from formula (9) gives

$$\Omega_k^{(\infty)} = \frac{\omega\bar{\Omega}^{(\infty)}(s^2 - 1)k^2}{\bar{\Omega}^{(\infty)}(s^2 - 1) - \omega k^2}; \quad (11)$$

$\bar{\Omega}^{(\infty)}$ is implicitly defined from the equation $\bar{\Omega}^{(\infty)} = \frac{3}{4\pi} \int \Omega_k^{(\infty)} d^3k$.

Now, let us consider the properties of the sequence $U, U^{(1)}, U^{(2)}, \dots$ at $\bar{x} = 0$. Transformation $U^{(n)} \rightarrow U^{(n+1)}$ has a non-trivial unstable fixed point $U_f(x)$. This function is shown in the upper panel of Fig.1. Consider small deviations from this fixed point: $U^{(n)}(x) = U_f(x) + u_n(x)$. The linearization of the transformation (10) gives

$$u_{n+1}(y) = u^0 + s^3 \int u_n(x) A(x, y) dx; \quad (12)$$

$$A = \exp\left(-\frac{U_f(x) - s^{-3}U_f(y) + F + \frac{\bar{\Omega}}{2}(x - y/\sqrt{s})^2}{T}\right).$$




FIG. 1: Upper panel: the fixed point of the transformation (10) at $s = 2.5$. Lower panel: critical index ν vs. scaling factor s . Note a very weak dependence of $\nu(s)$.

where u^0 stands to fulfill the condition $u_{n+1}(0) = 0$. The analysis of this operator shows that among its eigenvalues only two exceed unity. Denote them s^{λ_e} and s^{λ_o} ; they correspond to an even and odd eigenvector, respectively. Thus the theory is consistent with the scaling hypothesis, as it was mentioned in an introductory part.

Exponents λ_e and λ_o are the inverse of the critical indices ν and μ , respectively [6, 10]. In the ideal case, critical indices should be independent of the particular $U(x)$ and Ω_k , as well as of the value of the factor s . Indeed, after a proper re-scaling of the units, the values of T and $\bar{\Omega}$ drop out from Eq.(10). Therefore in our model ν and μ can depend only on s . Further, it turns out that $\mu = 2/5$ at any s , as it should be for the local-potential approximation [7]. What is about ν , it depends on s very weakly. The numerical result for $\nu(s)$ is shown in the lower panel of Fig.1.

On the other hand it is important to point out that the crude universal properties of the model enter the transformation (10) essentially. The behaviour $\Omega_{k \rightarrow 0} \propto k^2$ corresponds to a short-range range interaction. Such behaviour and the dimensionality of the model determine the scale transformation, *i.e.* powers of s in Eq.(10). It is also directly reflected in the formula that the order parameter is a scalar.

For the more detailed analysis it is reasonable to consider an extreme value of $\nu(s)$, because in this case small

TABLE I: Values of the critical indices obtained from the Lyapunov exponents for the transformation (12) at $s = 2.5$. Results of the Landau theory and known values [6] for 3D Ising model are given for the comparison.

Index	Estimation from Eq.(12)	3D Ising	Landau theory
ν	0.652	0.63	1/2
μ	2/5	0.403	1/3
β	0.326	0.327	1/2

variations of s do not affect ν . This gives $s \approx 2.5$, $\nu \approx 0.652$. The obtained values of indices are collected in Table 1. An agreement with the numerical results for the 3D Ising model is even better than one could expect. Since the scaling hypothesis holds in the model, the values of all other critical indices can be determined and should have a small error bar. As an example, the calculated value of β is presented in Table 1.

The advantage of the present approach is that it allows to estimate not only critical indices, but also the (non-universal) thermodynamic quantities in a wide thermal range. Here we calculate the dependence of the order parameter \bar{x} on temperature for the 3D Ising model. The model is defined as a discrete cubic lattice with $U(x) = \delta(x-1) + \delta(x+1)$ and the nearest-neighbors interaction $\Omega_k = 3 - \cos \pi k_x - \cos \pi k_y - \cos \pi k_z$. The initial Brillouin zone is a cube $-1 < k_{x,y,z} < 1$; at the first step the value $\bar{\Omega} = 3$ is used and all the modes with $|k| > 1/s$ are integrated. The procedure for further steps is exactly as described above. To simplify the calculation, we put $\bar{\Omega}^{(n)} = \bar{\Omega}^{(\infty)}$ at these steps; this simplification almost does not change the result obtained. To determine the value of \bar{x} we use the condition

$$\frac{\partial}{\partial x} U_n(\bar{x}_n) = 0, n \rightarrow \infty \quad (13)$$

(compare with the derivation of formula (7)). Numerically, it is enough to calculate U_n up to $n \approx 8$. The result for $\bar{x}^2(T)$ is presented in Fig.2. The mean-field and experimental (Monte-Carlo) data are sketched in the same figure for comparison. Far from the transition $U^{(1)}$ is almost parabolic, therefore the theory passes into MFA. The critical behaviour occurs at the transition point. Note that Ising model is very non-linear, but the scheme still performs good.

We conclude that the presented low-order scheme describes correctly all the physics of the second-order phase transitions in 3D systems with a short-range interaction and a scalar order parameter. It gives a unified description both of the fluctuation region and of the system far

from the transition point. The method does not pretend to be very accurate; its advantage consists in simplicity. In principle, one can improve the accuracy by a higher-order procedure for the integration in RG transformation. An applicability of the method for another universality




FIG. 2: The dependence of the square of the order parameter on the temperature in the 3D Ising model. Solid curve: the present approach with $s = 2.5$; the dot curve: mean-field calculation; the dash curve: numerical results. Thin ticks show the tangents to $\bar{x}^2(T)$ in the transition point.

classes is the subject of further studies.

The work was supported by RFFI foundation (grant 00-02-16253) and by the "Russian Scientific Schools" program (grant 96-1596476).

* Electronic address: alex@shg.ru

- [1] L. Onsager Phys. Rev. **65** 117 (1944).
- [2] L.D. Landau Phys.Z. SU **11** 26 (1937), *ibid* **11** 545 (1937).
- [3] V.L.Ginzburg Sov.Phys.Solid State **2** 1824 (1960).
- [4] L.P. Kadanoff Physics **2** 263 (1966), A.Z. Patashinskii, V.L. Pokrovskii Sov. Phys. JETP **50** 439 (1966).
- [5] K.G. Wilson and M.E. Fisher Phys. Rev. Lett. **28** 240 (1972).
- [6] for a recent review and further references, see A. Pelissetto and E. Vicari e-print cond-mat/0012164.
- [7] F.J. Wegner and A. Houghton Phys. Rev. A **8**, 401 (1973), C. Bagnuls and C. Bervillier, Physics Reports, **348** 91 (2001).
- [8] C.Bagnuls and C. Bervillier Phys.Rev. B **32**, 7209 (1985).
- [9] S. Aubry J. Chem. Phys. **62** 3217 (1975), A.N. Rubtsov, J. Hlinka, and T. Janssen Phys. Rev. E **61** 126 (2000).
- [10] R.Shankar Rev. Mod. Phys. **66** 129 (1994).