

Heat conduction and Wiedemann-Franz Law in disordered Luttinger Liquids

Mei Rong Li¹ and E. O. Rignac²¹Department of Physics, University of Guelph, Guelph, Ontario, Canada N1G 2W1²LPTENS, CNRS UMR 8549, 24, Rue Lhomond - 75231 Paris Cedex 05, France

We consider heat transport in a Luttinger liquid (LL) with weak disorder and study the Lorenz number for this system. We start at a high- T regime, and calculate both the electrical and thermal conductivities using a memory function approach. The resulting Lorenz number L is independent of T but depends explicitly on the LL exponents. Lowering T , however, allows for a renormalization of the LL exponents from their bare values by disorder, causing a violation of the Wiedemann-Franz law. Finally, we extend the discussion to quantum wire systems and study the wire size dependence of the Lorenz number.

PACS Numbers: 71.10.Pm, 72.15.Eb, 72.15.Nj

The Wiedemann-Franz (WF) law states that for electrons elastically scattered by impurities the ratio of the thermal (κ) and electrical (σ) conductivities normalized by T is a universal number $L_0 = \kappa/\sigma T = 3e^2/4\pi^2$ called the Lorenz number. This law is obeyed for electrons in conventional metals [1], and inclusion of electronic interactions that lead to the standard Landau Fermi-liquid (FL) behavior does not modify L_0 [2], as their effect on κ and σ scales away when the particle-particle scattering rate is much smaller than the impurity scattering rate.

A fundamental question arises whether the WF law survives in strongly correlated electron systems with non-FL ground states. Interest in this question has been recently revived by an observation of violation of WF law in high- T_c cuprates [3] which are widely believed to be non-FL systems. An important and tractable example of a non-FL is a one-dimensional (1d) Luttinger liquid (LL) with separate charge (ρ) and spin (ρ_s) bosonic collective excitations [4,6] instead of fermionic quasiparticles. Heat being carried by both spin and charge excitations, there is a potential for violation of WF law in a LL. However, a potential complication in the LL case is that even a weak disorder can cause Anderson localization [7,11].

Experimentally, thermal transport in (quasi)-1d has been investigated in spin-chain, [12] spin-Peierls, [13] and spin-ladder materials [14]. There are also some early measurements of heat conduction in quasi-1d organic conductors [15]. Theoretically, although $\kappa(T)$ in the LLs has been extensively discussed [16-18] with memory function methods [19], the thermal conductivity and the WF law have only been addressed in two contexts. First, Kane and Fisher (KF) [20] examined spinless LLs with a single strong impurity for repulsive electronic interactions, and found a T -independent but nonuniversal Lorenz number $L^{(KF)}$. The generalization of this problem to the bulk disorder case is however not straightforward [21] (as discussed below). Second, Fazio et al. [22] and Kiviere [23] have studied thermal transport in a mesoscopic 1d wire connected to leads. They showed that even for a clean wire the contact with leads causes violation of the WF law except at sufficiently low T . However, the effect of backward scattering by finite-density impurities has not been considered in the latter case.

In this letter, we first examine the WF law for an infinite LL with a nonzero concentration of impurities. Unlike KF, we calculate $\kappa(T)$ and $\sigma(T)$ in the high- T regime where impurity scattering can be treated by the memory function method [19,16]. The resulting Lorenz number is independent of T but depends on the LL exponents. It differs from the one obtained by KF. As T goes down, we show that the Lorenz number acquires T dependence as the consequence of instability towards Anderson localized state. Such a violation of the WF law is in sharp contrast to the higher-d FL situation. Finally we examine the effects of contact in a realistic quantum wire measurement and discuss the wire-size dependence of the Lorenz number.

Conductivities from Memory function method. A LL of spin-1/2 fermions in the presence of impurities is described by the following Hamiltonian [11]

$$H = H_{LL} + H_{imp} = \int dx H(x); \quad (1)$$

$$H_{LL} = \sum_{i=1}^X \int_{-\infty}^{\infty} \frac{dx}{2} u_i K_i [\rho_i(x)]^2 + \frac{u_i}{K_i} \partial_x \rho_i(x)^2; \quad (2)$$

$$H_{imp} = \int dx \left[\frac{p}{2} \rho(x) \phi_x + \frac{(x)}{a} e^{i\frac{p}{2}\phi(x)} \cos\left[\frac{p}{2}\phi(x)\right] + H.c. \right]; \quad (3)$$

where K_i are the LL exponents, u_i the velocities of the excitations, a the lattice constant, and $\rho_i(x)$ and $\phi_i(x)$ are canonical momenta and coordinates, respectively. $\phi(x)$ in Eq. (3) is the $2k_F$ component of the impurity potential, assumed to have a Gaussian distribution with zero average, $\overline{\phi(x)\phi(x')} = D\delta(x-x')$: $\phi(x)$ is the forward scattering component. The charge and heat currents $J_c = \int dx J_c(x)$ and $J_Q = \int dx J_Q(x)$ are obtained from the continuity equations $\partial_x J_c(x) + \partial_t n(x) = 0$ and $\partial_x J_Q(x) + \partial_t H(x) = 0$, where $n(x) = \frac{p}{2} \partial_x \phi(x)$ is the electron density operator. We find

$$J_c(x) = \frac{p}{2} u K \phi(x); \quad (4)$$

$$J_Q(x) = u^2 \partial_x \phi(x) \partial_x \phi(x) + J(x); \quad (5)$$

where $J(x) = \frac{P}{2u} K(x)$ (x): We note that the transformation $\tilde{\psi}(x) = \psi(x) \frac{P}{2u} R_x(x^0) dx^0$ eliminates from both H_{imp} and J_Q , implying a null effect of (x) on c and q . Thus we set $\tilde{\psi}(x) = \psi(x)$ hereafter. Eqs. (4) and (5) also show that the heat and charge current operators are quadratic and linear in boson operators, respectively. This is crucial in resulting in deviation of the Lorenz number from the universal number at high T even for spinless fermion case as shown below.

In the absence of impurities, both J_c and J_Q are conserved currents. In fact, J_Q is (up to a prefactor) the total momentum of the system, and its conservation results from translational invariance. With impurities, neither J_c and J_Q are conserved and finite conductivities can be expected [18]. At sufficiently high T , quantum effects are cut off by inelastic thermal processes, and disorder scattering can be treated within the Born approximation. This yields conductivities to leading order in D^{-1} . A convenient formalism to do this is the memory function method [19,17,18], in which the finite-frequency electrical and thermal conductivities are expressed in terms of the memory functions $M_j(\omega) = \frac{1}{j!} \frac{d}{d\omega} \text{Im} F_j(\omega) |_{\omega=0}$ ($j = c, Q$) as

$$c(\omega) = ie^2 c[\omega + M_c(\omega)]^{-1}; \quad (6)$$

$$q(\omega) = i q_0 T^{-1} [\omega + M_Q(\omega)]^{-1}; \quad (7)$$

with $j = \frac{R_1}{T} \int_0^T d\tau h_j(0)^y J_j(i\tau)$ is the current static susceptibilities, $F_j(x) = [J_j(x); H]$, and $\text{Im} F_j(\omega) = \frac{1}{2\pi} \int_0^{\omega} d\omega' \frac{d}{d\omega'} \text{Im} F_j(\omega'; 0) |_{\omega=0}$. To leading order in D , both c and q take their pure LL values, $c = \frac{2}{3} u K$; $q = \frac{1}{3} (u + u) T^2$, and

$$\text{Im} F_j(\omega) \sim D^{-2K_t+2} (a)^{K_t-2} u^K u^K A_j(\omega)$$

$$\sin \frac{K_t}{2} B \frac{K_t}{2} \frac{i!}{2T}; 1 - K_t T^{-j}; \quad (8)$$

where $K_t = K + K$; $B(x; y)$ is the Euler Beta function, $c = 1 - K_t$, $q = c/2$, $A_c(\omega) = K^2 u^2 =$; and $A_Q(\omega) = [K^2 = 4 + \omega^2 = 2 T^2 (K u^2 + K u^2) = K_t (K_t + 1)]$: Inserting Eqs. (8) into Eqs. (6-7), and after some straightforward algebra, we find that the dc conductivities become

$$c(T) = \frac{2e^2 (K_t)}{2 (K_t=2)} \frac{a}{D} \frac{2 a T^{-2K_t}}{u} = e^2 l_{el}(T); \quad (9)$$

$$q(T) = \frac{2}{9} \frac{2T}{K_t} \frac{1 + K_t}{K_t} \frac{(u + u)^2}{u^2 K + u^2 K} \frac{(K_t)}{2 (K_t=2)} \frac{a}{D} \frac{2 a T^{-2K_t}}{u} / C_v(T) l_{el}(T); \quad (10)$$

where $D = (2D a = u^2) (u = u)^K$ is a dimensionless disorder parameter [11], $\Gamma(x)$ the Gamma function, $C_v(T)/T$ the specific heat of a pure LL, and $l_{el}(T)$ the elastic mean free path [8]. Both c and q exhibit power-law behavior in T with nonuniversal exponents.

Lorenz number at high T . From Eqs. (9) and (10), one has

$$L = \frac{c}{qT} = \frac{2}{9e^2} \frac{(u + u)^2}{u^2 K + u^2 K} \frac{1 + K_t}{K_t}; \quad (11)$$

K , K , u and u are bare intrinsic parameters, so L is independent of T , and a generalized Wiedemann-Franz law is obeyed. In the noninteracting case, $K = K = 1$, $u = u = v_F$, the universal Lorenz number $L = L_0$ is recovered from Eq. (11).

The result (11) can be reduced to the spinless LL case, by making $u = u = u$, $K = K = K$. We obtain

$$L^0 = \frac{2}{9e^2} (K^2 + 2K^{-1}); \quad (12)$$

It follows that $L^0 < L_0$ for an attractive interaction case $K > 1$, as a result of tendency towards a superconducting state with high electrical conduction but poor thermal conduction; While for a repulsion case $K < 1$, $L^0 > L_0$, indicating tendency towards a weakly pinned charge density wave (CDW) state with better heat transport than charge transport. It is instructive to compare L^0 with $L^{(KF)} = \frac{2}{9e^2} (K^2 + 2K)^{-1}$ obtained by KF for a single strong impurity at $K < 1$ [20]. We see that $L^0 - L^{(KF)} = 2(K - 1)^2 = 9K^2 (K + 2)^{-1} > 0$. From physical point of view, $L^{(KF)}$ and L^0 can be roughly understood as the results for a strongly pinned and a weakly pinned CDW phase, respectively, so that we should expect a larger $L^{(KF)}$ than L^0 . This puzzle can be resolved by noting that some Hamiltonian terms causing heat but no charge conduction are neglected in the derivation of the tunneling Hamiltonian used to calculate $L^{(KF)}$ [20]. Including these terms explicitly for $K = 1/2$ indeed leads to [20] $L^{(KF)} = \frac{3}{2} L^{(KF)} > L^0$.

We would also like to remark that the single strong impurity case considered by KF corresponds to a low energy fixed point which cannot be reached from the limit of weak impurity scatterers at high concentration c we consider [21]. The KF fixed point is realized when the renormalized impurity strength $t_B(\omega) = e^{(2-2K)\omega/t_B(0)}$ becomes of the order of the high energy cutoff $\omega = a$. This requires the renormalized length $a e^{\omega} = a [\omega = t_B(0)]^{\omega/(2-2K)}$ to remain still much smaller than the inter-impurity distance $l = c$. In the high impurity concentration limit, $c \rightarrow 1$, $t_B \rightarrow 0$ with $D = c t_B^2$ fixed, obviously such regime cannot be observed.

Lorenz number with decreasing T . The enhanced quantum interference effects, which are responsible for occurrence of the Anderson localization at $T < T_{loc} = u = l_{loc}$ ($l_{loc} = a D^{-1/(3-K_t)}$ [11]), lead to renormalization of K , K and D . Now we study the influence of such renormalization on L for T still much larger than T_{loc} , so that a perturbative renormalization group (RG) theory can be used. We first neglect renormalization of K and K by D . The RG flow equation determining the renormalization of D reads [11] $dD(l) = d l = (3 - K_t) D(l)$; which leads to

$$D(l) = D = e^{(3/K_t)l}. \quad (13)$$

It is clear that $K_t = 3$ describes a metal-insulator transition (MIT) line: for $K_t < 3$, $D(l)$ grows exponentially with renormalized length, and the Anderson localization will take place eventually; whereas for $K_t > 3$, $D(l)$ flows to zero, corresponding to a delocalized phase. By defining a T -dependent scale $l(T)$ from $ae^{l(T)} = u = (2T) = l_{th}(T)$ (l_{th} the thermal length), we see that Eq. (13) yields $(a/D)(2aT/u)^{2/K_t} = ae^{l(T)} = [D(l)]$; which allows for a simple RG interpretation for ϵ_c and ϵ_{in} in Eqs. (9-10): They can be obtained by applying the RG flow up to the scale $l(T)$ at which T becomes order of the energy cutoff $u = a$. At this scale, thermal effects suppress quantum effects, and the use of the memory function method is justified. One can then perform a memory function calculation in which the bare parameters D and a are replaced by $D(l)$ and ae^l , respectively.

A similar analysis can be made for inclusion of renormalization of K and K by D . We start from the bare parameters $K(0)$, $K(0)$ and $D(0)$ and follow their RG flow, being described by Eqs. (3.4) in Ref. [11], up to the scale $l(T)$. Since $D(l) \ll 1$ still holds for $T \gg T_{loc}$, but the quantum phase coherence has started to lose, Eqs. (9-10) are valid again [24], but with the bare K and K replaced by the renormalized ones, $K(l)$ and $K(l)$, respectively. The resulting RG-improved L acquires T dependence through K and K , and the generalized WF law holding at high T breaks down. This is in contrast to a higher-d FL case, in which such a renormalization is always negligible [2].




FIG. 1. Lorenz number as a function of temperature close to the MIT line for spinless fermions. $W = a$ is a high energy cutoff. The initial conditions are $K(0) = 3/2$; $D(0) = 10^{-2}$.

We proceed with a discussion of the spinless fermion case for simplicity. The above analysis suggests that L^0 in Eq. (12) as a function of the initial parameter K becomes

$$L(T) = (e^2/9e^2) f[K(l)]^2 + 2[K(l)]^1 g; \quad (14)$$

where $K(l)$ can be obtained from the RG flow equation

[11], $dK(l) = dl = K^2(l)D(l) = 2$. For a generic initial K away from the MIT line, the RG flow line in the K - D plane is almost vertical, and the deviation of L from its high- T value is in order of D which is insignificant unless T becomes close to T_{loc} . Alternatively, considering the second order terms in D in ϵ_c and ϵ_{in} one also obtains a T -dependent power-law correction term to $L(T)$ with the exponent predicted by the RG method.

However, in the vicinity of the MIT line, the RG flow is no longer vertical, leading to a faster variation of L with T . $K(l) = K_j$ is found to be proportional to $D(l) = D$, indicating that $L(T)$ shown in Eq. (14) is systematic. $L(T)$ for this case is shown in Fig. 1. We emphasize that such a behavior is not expected in a higher-d FL case, in which the Lorenz number is constant even close to the MIT point [2].

Lorenz number in dirty quantum wires. An experimental realization for the measurement of the transport properties of a LL is through a finite size wire connected to leads [25] or a carbon nanotube [26]. The theory presented here for a bulk material needs to be modified, as the boundary condition imposed by the contact is shown [27,22] to drastically influence the transport through the wire. Charge crosses the contact in terms of pieces of fractional charges [28], whereas heat is carried by plasmon modes and they cross the contact like waves passing through barriers [22]. The Lorenz number will obtain a nontrivial wire size dependence as shown below. For simplicity we only discuss about spinless LL wires, and leave the discussion of the spinful case for a future publication [29]. Besides, we follow Ref. [27] and model the leads as 1d noninteracting FLs.

We first briefly recall the clean wire case. The electrical conductance G is purely from the contact and is quantized [27], $G = G_0 = e^2/2$ (for $\hbar = 1$), which implies a perfect transmission of the total charge after a long time (dc) measurement [28]. The mismatch of the plasmon wave velocities at the two sides of the contact leads to T -dependent thermal conductance K_{th} normalized by T [22], which, at low temperatures $T \ll d$ (d the wire size), asymptotically approaches to a constant $K_{th}/T = 6$ [22]. The WF law is typically violated for a generic T and is restored at $T \gg d$.

Let us now turn to the case of a wire with impurities. We consider the case of l_{loc} being much larger than d and/or l_{th} so that we can neglect Anderson localization. The total electrical resistance is the sum of the contact resistance G_0^{-1} and the wire resistance G_{wire}^{-1} , resulting in the following total conductance,

$$G = (G_0^{-1} + G_{wire}^{-1})^{-1}; \quad (15)$$

G_{wire} depends on the relative values of the two length scales d and $l_{th}(T)$.

1) At $d \ll l_{th}(T \gg d)$, a simple Ohm law holds for the electrical conductance, i.e., $G_{wire} = (T)/d = G_0/2$ with (T) and l_{th} being found in Eq. (9). Eq. (15) reduces, for $d \ll l_{th}(T)$, to Maslov's perturbation

result [30], $G \propto G_0 [1 - d/(2l_{el})]$; and, for $d \ll l_{el}(T)$, crosses to the Dnude formula. In this regime, the thermal conductance of even a clean wire, K_{th} , from Eq. (9) of Ref. [22], depends on the details of the barrier, so that the thermal conductance of the dirty wire becomes a rather non-universal function of disorder, wire size, and temperature [29].

2) At $d \ll l_{th}(T \rightarrow d)$, Eq. (9) of Ref. [22] indicates that heat is carried by plasmon modes of wavelength much larger than the distance between the contacts, and thermal conductivity of the clean wire becomes universal. In this regime, we can make use of the $T \rightarrow d$ equivalence [31,30] to obtain an explicit d -dependent Lorenz number. According to Ogata and Fukuyama, the finite size effect can be taken into account within the memory function formalism by replacing l in Eqs. (6) and (7) by $id \rightarrow d$. This leads to

$$G(d) = \frac{ue^2}{2u + 2u C_c(u;K)} \frac{1}{(d)}; \quad (16)$$

$$K_{th}(d) = \frac{uT}{3} \frac{1}{2u + 2u C_{th}(u;K)} \frac{1}{(d)}; \quad (17)$$

with $(d) = (d - l_{oc})^{3/2K}$, and $C_c(u;K)$ and $C_{th}(u;K)$ being easily found by comparing Eqs. (16) and (17) with Eqs. (9) and (10). In writing down Eq. (16) we have carefully taken into account the screening of the electric field coming from the leads [32]. We recover a universal Lorenz number in this regime in the absence of impurities in agreement with [22]. In the presence of impurities, Eqs. (16) and (17) immediately yield

$$L(d) \propto L_0 \frac{1 + C_c(u;K)}{1 + C_{th}(u;K)} \frac{1}{(d)}; \quad (18)$$

which is a function of the wire size since $C_c(u;K) \neq C_{th}(u;K)$ in the presence of interactions. Thus, deviation from WF law is obtained in finite size systems with impurities even for $T \rightarrow d$.

Conclusion. We have investigated the WF law in a disordered LL system. At high T where thermal effects cut off Anderson localization, the Lorenz number is constant in T , and, in the spinless fermion case, smaller (larger) than the universal L_0 for attraction (repulsion) between the fermions. Its dependence on the LL exponents is different from the one obtained in a single strong impurity case [20]. As T goes down, Anderson localization effects induce an interplay between disorder and electronic interactions, which is responsible for the violation of the WF law. When a dirty 1d wire is connected to leads we find the Lorenz number becomes a function of the wire length.

Acknowledgments. We are indebted to N. Andrei, P. J. Hirschfeld, A. Rosch, and Y.-J. Wang for valuable discussion. We also thank T. Giamarchi, Y. Suzumura, and I. Vekhter for comments on the manuscript.

- [1] J.M. Ziman, *Electrons and Phonons* (Clarendon, Oxford, 1962); A.A. Abrikosov, *Fundamentals of the theory of metals* (Elsevier Science, Amsterdam, 1988).
- [2] C. Castellani et al., *Phys. Rev. Lett.* 59, 323 (1987).
- [3] R.W. Hill et al., *Nature* 414, 711 (2001).
- [4] V.J. Emery, in *Highly Conducting One-Dimensional Solids*, edited by J.T. Devreese, R.P. Evrard, and V.E. van Doren (Plenum Press, New York and London, 1979).
- [5] J. Solym, *Adv. Phys.* 28, 209 (1979).
- [6] J. von Delft and H. Schoeller, *Ann. Phys. (Leipzig)* 7, 225 (1998).
- [7] W. Apel, *J. Phys. C* 15, 1973 (1982).
- [8] W. Apel and T.M. Rice, *J. Phys. C* 16, L271 (1982).
- [9] W. Apel and T.M. Rice, *Phys. Rev. B* 26, 7063 (1982).
- [10] Y. Suzumura and H. Fukuyama, *J. Phys. Soc. Jpn.* 52, 2870 (1983); *ibid.* 53, 3918 (1984).
- [11] T. Giamarchi and H.J. Schulz, *Phys. Rev. B* 37, 325 (1988), and references therein.
- [12] A.V. Solgubenko et al., *Phys. Rev. B* 62, R6108 (2000); *Phys. Rev. B* 64, 054412 (2001).
- [13] Y. Ando et al., *Phys. Rev. B* 58, R2913 (1998); J. Takeya et al., *Phys. Rev. B* 61, 14700 (2000); *ibid.* 62, R9260 (2000); *ibid.* 63, 214407 (2001).
- [14] A.V. Solgubenko et al., *Phys. Rev. Lett.* 84, 2714 (2000); C. Hess et al., *Phys. Rev. B* 64, 184305 (2001); K. Kudo et al., *J. Phys. Soc. Jpn.* 70, 437 (2001).
- [15] M.Y. Choi, P.M. Chaikin, and R.L. Greene, *Phys. Rev. B* 34, 7727 (1986).
- [16] A. Luther and I. Peschel, *Phys. Rev. Lett.* 32, 992 (1974).
- [17] T. Giamarchi, *Phys. Rev. B* 44, 2905 (1991).
- [18] A. Rosch and N. Andrei, *Phys. Rev. Lett.* 85, 1092 (2000).
- [19] W. Gotze and P. Wölfe, *Phys. Rev. B* 6, 1226 (1972).
- [20] C.L. Kane and M.P.A. Fisher, *Phys. Rev. Lett.* 76, 3192 (1996).
- [21] T. Giamarchi and H. Maurey in *Correlated fermions and transport in mesoscopic systems* edited by T. Martin, G.M. Montambaux and J. Tran Thanh Van (Editions Frontieres, Gif sur Yvette, 1996).
- [22] R. Fazio, F.W.J. Hekking, and D.E. Khmelnitskii, *Phys. Rev. Lett.* 80, 5611 (1998).
- [23] I.V. Krive, *Fiz. Nizk. Temp.* 24, 498 (1998), [*Low Temp. Phys.* 24, 377 (1998)].
- [24] Certain type of interactions in the spin channel may cause complication. This is left for a future publication. [29]
- [25] S. Tanucha, T. Honda, and T. Saku, *Sol. State Comm.* 94, 413 (1995).
- [26] P. Kín, L. Shi, A. Majumdar, and P.L. McEuen, *Phys. Rev. Lett.* 87, 215502 (2001).
- [27] D.L. Maslov and M. Stone, *Phys. Rev. B* 52, R5539 (1995).
- [28] I. Sa and H.J. Schulz, *Phys. Rev. B* 52, R17040 (1995).
- [29] M.-R. Li and E. Ognac, in preparation (unpublished).
- [30] D.L. Maslov, *Phys. Rev. B* 52, R14368 (1995).
- [31] M. Ogata and H. Fukuyama, *Phys. Rev. Lett.* 73, 468 (1994).
- [32] A. Kawabata, *J. Phys. Soc. Jpn.* 65, 30 (1996).