

Comment on "Exact Spectral Functions of a Non-Fermi Liquid in 1 Dimension"

A. A. Zvyagin

B. I. Verkin Institute for Low Temperature Physics and Engineering,
Ukrainian National Academy of Sciences, 47 Lenin Avenue,
Kharkov, 61164, Ukraine

March 22, 2024

Abstract

We point out the incorrect statement of the recent manuscript by K. Penc and B. S. Shastry.

In a recent manuscript [1] K. Penc and B. S. Shastry wrote in the first paragraph: "Schulz and Shastry [2] have introduced a new class of gauge-coupled one-dimensional (1D) Fermi systems that are non-Fermi liquid in the sense that their momentum distribution function has a cusp at the Fermi momentum k_F rather than a jump as in a Fermi liquid [3]. This behavior is of the sort first found by Luttinger in the context of his study of a one-dimensional model that is popularly known as the Luttinger model [4]. The model introduced by Schulz and Shastry (SS) is in fact intimately connected to the Luttinger model, and is best viewed as a reinterpretation of Luttinger's original model as a gauge theory. Particles of different species exert a mutual gauge potential on each other, and this is sufficient to destroy the Fermi liquid. This model has the added property that the charge and spin correlations are unaffected by the interaction, owing to the 'gauge' nature of interaction."

I would like to point out that the main statement of this paragraph is incorrect. In fact, this can be seen from the second reference from [2] (which is the Reply to my previous Comment [5]) and that Comment itself. It turns out that all the properties, mentioned in the cited paragraph, namely |

The gauge coupling between 1D Fermi systems, which produces the non-Fermi liquid behavior (the Luttinger liquid behavior);

Particles of different species in that class exert a mutual gauge potential on each other (which is sufficient to destroy the Fermi liquid);

The charge and spin correlations are unaffected by the interaction, owing to the "gauge" nature of interaction

It had been already known for the class of models, introduced by us in Refs. [6, 7] in 1992. This is why the priority of introducing the models with these properties belongs to us. One can check that particles in [6, 7] of different species are connected with each other via 'gauge' potentials (reminiscent of the Peierls phase factors), similar to the 'gauge' potentials in [2], and that we emphasized on the Luttinger liquid behavior of our 1D models with this 'gauge' couplings already in 1992, much earlier than [2]. Even the title of [7] is "Exactly solvable models of an effectively two-dimensional Luttinger liquid". Notice that in [6, 7] we interpreted the additional index, which distinguishes species of particles, as a number, which enumerates 1D chains, coupled with each other via gauge potentials. By the way, SS in their first paper of [2] never used this interpretation, but now, in [1], the authors already imply that the class of models, introduced by SS described coupled 1D chains.

According to the Reply [2], SS introduced some class of models, which had all mentioned above properties of the class, introduced in [6, 7], but with the additional constraint: Some recursion relations for 'gauge' potentials have to be satisfied. SS especially pointed out that difference, see, please, the footnote [5] of the Reply [2], where they wrote: "We use the term 'class of models' in the specific sense that the members share a common method of solution, rather than a vague sense in which many models share certain physical properties." Obviously, only the properties, mentioned in the first paragraph of [1], cannot properly define the class of models, introduced by SS, because they belong to both classes: [6, 7] and [2].

Actually, there are two alternatives:

(1) Either SS introduced a subclass (with some specific, additional properties) of the class of models [6, 7] (which had been earlier introduced in our papers) with the common properties, mentioned in the disputed paragraph. Certainly, one cannot introduce any new class of models in 1998 with the same properties as the models, introduced in 1992.

(2) Or SS (cf. their Reply [2]) introduced some new class of models, different from ours. However in this case the authors of the manuscript [1] could clearly define the properties, which determine only the models, introduced by SS, but which are not present in the class, introduced earlier in [6, 7]. At least they could carefully distinguish between those two classes, and not to emphasize on the properties, which belong to the other class of models, introduced in [6, 7] in the definition of the models, introduced by SS.

In both cases the statements of the disputed paragraph [1] are wrong. I do not imply that the model, studied in [1] belongs to the class of models, introduced in [6, 7] in the sense (2). However, clearly, when writing about the properties of the models, introduced in [2], the authors of [1] could properly write about the features, which pertain only to the class introduced by SS, but not about the ones, which had been known for the other class of models, introduced earlier by us in [6, 7]. It turns out that according to the definition of the Reply [2], all mentioned in the disputed paragraph of [1] properties have namely "a vague sense in which many models share certain physical properties".

This is why, immediately after the manuscript [1] appeared in the Archive, I asked the authors of [1] to correct the statement of the disputed paragraph. However the authors did not agree, i.e., they insist that the class of models with the properties, mentioned in the cited paragraph of their work, was introduced in [2]. But then a contradiction exists: If the statements of the Reply [2] are correct, then the statements of the first paragraph of the manuscript [1] are obviously wrong.

It is the goal of this my Comment to point out this contradiction.

References

- [1] K . Penc and B . S. Shastry, preprint cond-mat/0109339.
- [2] H . J. Schulz and B . S. Shastry, Phys. Rev. Lett. 80, 1924 (1998); Phys. Rev. Lett. 82, 2410 (1999).
- [3] I. E. D zyaloshinskii and A . I. Larkin, Zh. Eksp. Teor. Fiz. 65 411 (1973) [Sov. Phys. JETP 38, 202 (1974)]; H . Shiba and M . Ogata, Int. J. Mod. Phys. B 5, 31 (1991).
- [4] S. Tom onaga, Prog. Theor. Phys. 5, 349 (1950); J. M . Luttinger, J. Math. Phys. 4, 1154 (1963); D . C . Mattis and E . Lieb, ibid. 6, 304 (1965).
- [5] A . A . Zvyagin, Phys. Rev. Lett. 82, 2409 (1999).
- [6] A . E. Borovick, A . A . Zvyagin, V . Yu. Popkov and Yu. M . Strzheim echnyi, Pis'ma Zh. Eksp. Teor. Fiz. 55, 293 (1992) [JETP Lett. 55, 292 (1992)].
- [7] A . A . Zvyagin, Fiz. N izk. Temp. 18, 1029 (1992) [Sov. J. Low Temp. Phys. 18, 723 (1992)].