

Fractal Plate Tectonics

Didier Somette

Laboratoire de Physique de la Matière Condensée, CNRS UMR 6622 and Université de Nice-Sophia Antipolis, Nice, France and Department of Earth and Space Sciences and Institute of Geophysics and Planetary Physics, University of California, Los Angeles, CA

Vladilen Pisarenko

International Institute of Earthquake Prediction Theory and Mathematical Geophysics, Russian Ac. Sci. W arshavskoye sh., 79, kor. 2, Moscow 113556, Russia

Abstract. We analyze in details the statistical significance of the claim by Bird [2002] of a power law distribution of plate areas covering the Earth and confirm that the power law with exponent 0.25 ± 0.05 is the most robust and parsimonious model for all plates, including the very largest plates, when taking into account the constraint that the plates areas must sum up to 4π steradians. We propose a general class of fragmentation models that rationalize this observation and discuss the implications for the earth dynamics and the general self-organization of tectonic deformations at multiple scales.

Introduction

According to the theory of plate tectonics, the outermost 100 km of the solid Earth is composed of a relatively small number of internally rigid plates (lithosphere) that move over a weak substrate (asthenosphere). The plate sizes and positions change over time and are driven by the internal mantle convection. Dewets et al. [1990] performed a global inversion to determine the relative rotation rates of the 12 largest plates (the NUVEL-1 model) later refined into the NUVEL-1A solution [Dewets et al., 1994]. The edges of these plates, where they move against each other, are sites of intense geologic activity, such as earthquakes, volcanoes, and mountain building. These plate edges are not sharp narrow boundaries but are often constituted of complex systems of competing faults and other geological structures of width extending over several hundreds of kilometers for transform and subduction boundaries and up to thousands kilometers for continental collisions. It is now common lore to view such tectonic deformation as possessing some kind of self-similarity or fractal properties [Scholz and Mandelbrot, 1989], or better a hierarchical structure [Duillbn et al., 1996]. In addition, several researchers have repeatedly proposed to describe this multi-scale organization of faulting by a hierarchy of blocks of multiple sizes [Gelfand et al., 1976; Gorshkov et al., 2001]. In these models, blocks slide against each

other, rotate, lock at nodes or triple junctions which may represent the loci of major earthquakes, in a way similar to (but at a reduced scale compared to) the relative motion of the major tectonic plates. An example of a complex network of faults is observed in the broad San Andreas transform boundary between the Pacific and the North American plates in California, for which Bird and Rosenstock [1984] have suggested 21 possible microplates within southern California alone.

Keeping in mind these ingredients of a few major plates at large scales on one hand and a hierarchical self-similar organization of blocks at the boundary scale on the other hand, the recent reassessment of present plate boundaries on the Earth by Bird [2002] is particularly inspiring: taking into account relative plate velocities from magnetic anomalies, moment tensor solutions, and/or geodesy, to the 14 large plates whose motion was described by the NUVEL-1A poles, model PB2001 [Bird, 2002] includes 28 additional small plates, for a total of 42 plates. Bird [2002] suggests that the cumulative-number/area distribution for his model follows a power-law for plates of less than 1 steradian of area.

Statistical analysis of the size distribution of the 42 plates

Analysis with rank-order statistics and the Pareto law

The statistical analysis of such a data set is very difficult due to its small size $N = 42$ but is not impossible. Figure 1 shows the complementary cumulative number $N(A)$ of plates as a function of area A in steradians, i.e., the number of plates with an area equal to or larger than A . Most of the data except for a few largest plates follow a linear dependence in the double log-scale of the figure. The slope of this straight line is close to 0.25. We first compare this sample with a pure power law (the Pareto distribution). For this purpose we use the rank-ordering method (see [Somette et al., 1996]). We put the sample into descending order $A_1 \geq A_2 \geq \dots \geq A_N$. The probability density function (PDF) of the n -th rank, denoted $p_N(x)$, is well-known

$$p_N(x) = (N - n + 1) \frac{1}{N} F^{N-n}(x) (1 - F(x))^{n-1} f(x); \quad (1)$$

where $F(x)$; $f(x)$ are the distribution function (DF) and PDF of the random values in question. Putting the Pareto law $F(x) = 1 - (a/x)^b$, $x \geq a$, into (1), we get

$$p_N(x) / (1 - (a/x)^b)^{n-1} x^{(N-n+1)b} = \quad (2)$$

The mode $M_{n,N}$ of the PDF (2) (i.e., the maximum of the PDF) is the most probable value of the random variable A_n :

$$M_{n,N} = a \frac{N+1}{n+1}^{1-\alpha} \quad (3)$$

Besides, an interval around the mode containing some prescribed probability (say, 90%) can be derived from the density $g_{n,N}(x)$. The dependence (3) with $\alpha = 1/4$ is shown as the long-dashed line in Fig. 1. The two short-dashed lines represent the upper/lower limits of the 90%-confidence interval. The data are well accounted for by the power law, except, perhaps, for the three smallest ranks, i.e. the three largest plates, the Pacific, the Antarctica, and the Africa plates, which fall outside the confidence interval.

"Finite-size" constraint on the PDF of plate sizes

From a visual inspection of figure 1, it might be argued that the deviation from the power law prediction (3) occurs somewhat earlier, say, at rank $n = 7$, i.e., the seven largest plates with area more than 1 steradian belong to a different population than the rest of the plates. The simplest possibility is that the largest plates are sensitive to a "finite-size" effect, namely that the sum of areas over all plates must sum up to 4π . The largest continent and ocean plates are commensurable with the size of the convection cells in the upper mantle and they would thus feel the constraints due to the finite size of the Earth surface. In contrast, one could argue that the smaller plates appeared as a result of the interaction and collision between the larger plates and do not feel any special restriction. More precisely, the smaller plates may be the result of fragmentation/creation processes in the neighborhood of the borders and especially near triple points of the seven largest plates.

The distribution $g_{N,C}(x)$ of sample values x_1, \dots, x_N conditioned by the constraint

$$S_N = x_1 + \dots + x_N = C \quad (4)$$

where C is a constant (4π for the plates) is modified from its unconditional Pareto density expression $f(x) = a x^{1-\alpha}$ for $x \geq a$ and $f(x) = 0$ for $x < a$. The lower threshold (the minimum value) for the plate data is $a = 0.00178$ steradian and corresponds to the smallest documented plate in [Bird, 2002]. Denoting the unconditional density of the sum $S_k = x_1 + \dots + x_k$; $k = 1, \dots, N$ by $s_k(x)$, we have $g_{N,C}(x) = s_{N-1}(C-x) f(x) / s_N(C)$, for $a \leq x \leq C$.

Thus, the constraint (4) decreases the unconditional Pareto density $f(x)$ by a factor $s_{N-1}(C-x)/s_N(C)$ which acts as a "taper."

In order to use the Maximum Likelihood (ML) method for the estimation of the exponent α , we need the vectorial distribution of conditional sample to take into account the interdependence between the different variables (areas of the plates) $x_1; \dots; x_{42}$ induced by the constraint (4). The corresponding likelihood function is therefore

$$f(x_1; \dots; x_{42}) = \frac{(x_1 + \dots + x_{42})^{4-1}}{s_{42}(4-1)} f(x_1) \dots f(x_{42}) : \quad (5)$$

The resulting ML estimate is $\alpha = 0.25 \pm 0.05$. With this value, we generate an artificial sample of 1045 conditional 42-dimensional (42D) vectors with the condition (4) with $C = 4$. Rank-ordering each of these 1045 vectors, we determine their sample medians $M_1; \dots; M_{42}$, where M_j is the median of the j -th rank. These conditional medians are slightly smaller than given by (3) for the unconditional Pareto distribution. The conditional distribution (5) allows us to construct a confidence domain for the 42D random vectors, defined as a "corridor" of the form $[cM_j; (1+c)M_j]$, the constant $c = 0.244$ being chosen such that 95% of vectors fall within this corridor. The medians $M_1; \dots; M_{42}$ and their corridor are shown in Fig. 2: all samples of the tectonic plates falls within the 95% confidence corridor, showing that (in contrast with the "pure" Pareto used in Fig. 1) the Pareto model together with the total area constraint (4) accounts satisfactorily for all the data, including the largest plates.

We have performed many other tests which all confirm the goodness of fit of the constrained Pareto distribution (or of its approximation by the truncated Pareto distribution). Using Wilks' theorem [Rao, 1965], we used the so-called Gamma-distribution with complementary cumulative distribution function $G(x)$, truncated from both sides $G(x; a; b) = C(a; b) \frac{e^{-ax}}{x^{1+b}}$ for $u_0 < x < u_1$ and find that (1) the introduction of the additional parameter a in the exponential does not increase significantly (probability of rejection 38%) the likelihood as compared with the truncated Pareto and (2) the truncated Pareto distribution is much preferable compared with the truncated exponential distribution (significance $1 \cdot 10^{-8}$). This conclusion is strongly confirmed by tests using so-called sufficient statistics for both the power law and exponential models. We also tested the popular lognormal DDF, which is a natural model for fragmentation since it is the limit distribution for processes of subdivision of rock under rather

general assumptions (basically multiplicative processes) [Kolmogorov, 1941]. Using the Kolmogorov test and the bootstrap method, we estimate that the data deviates significantly from the lognormal DF at the 95% significance level.

Discussion

Our main result is that Bird's suggestion [2002] of a power-like distribution of plates areas is confirmed up to the largest plates, when taking into account the very natural constraint that the plates areas must sum up to 4π steradians.

Given this strength of the validity of the Pareto distribution for the 42 plates documented in [Bird, 2002], one can expect that it can be extrapolated beyond this range to smaller yet undetected plates. Using the complementary Pareto distribution ($a=x$) with $a = 0.00178$ steradian and $\alpha = 0.25$, this extrapolation predicts 100 plates larger than $1.35 \cdot 10^{-4}$ steradian ($5,500 \text{ km}^2$ or $74 \cdot 74 \text{ km}$) [Bird, 2002]. The total area needed to define another 58 plates ranging from this size up to the smallest size in PB2001 would be only about 0.028 steradians, which could be taken from large plates like EU and NA without materially affecting their areas. As discussed by Bird [2002], the places where additional small plates are most likely to be recognized is within the zones of distributed deformation identified in [Bird, 2002], which have total area of 0.838 steradians (6.7% of Earth).

The main suggestion inspired by Bird's discovery is that the separation of scales between a plate tectonics involving a few major plates at large scales and tectonic deformation at smaller scales may be an artifact that resulted historically from insufficient statistics and more recently from a not quite correct interpretation of "visual" deviations from the pure power law. With the improved extended compilation of [Bird, 2002] which will probably still improve in the years to come, we conjecture that there is a hierarchy of plates and block sizes from the largest ones to the smallest scales, which may be the result of the self-organization of the plate motions, creation and destruction over millions of years. In other words, we suggest that our view of plate tectonics should be revised in terms of a dynamical model of plates with creation, fragmentation and destruction that may occur at all scales. This view may provide a link between the large organization of major plates down to the block structures at scales of tens of kilometers or less discussed in the introduction.

What could be the mechanism(s) at the origin of the observed power law distribution of plate areas? We pro-

pose a fragmentation model with source terms, which can be progressively enriched by taking into account the kinematic compatibility constraints between plates. There is a large class of such fragmentation models that can be solved explicitly within the scaling theory of [Cheng and Redner, 1988]. In a nutshell, their approach is to write down a linear integro-differential coupled equation for the time evolution of fragment populations. The fragment distribution $p(l)$ is characterized by the moments $m = \int_0^{\infty} l p(l) dl$. Note that calculating a moment corresponds to taking the Mellin transform of $p(l)$. The moments m give access to $p(l)$ by taking the inverse Mellin transform. The equations of evolution for the fragment population then transform into the following recurrence equation for the moments: $m_{k+1} = \frac{1}{L-1} m_k$, where L is the exponent describing the overall rate of breakup (assumed to be proportional to l), $\frac{1}{L-1}$ is a normalizing constant and $L = \int_0^1 x b(x) dx$, where $b(x)$ is the probability that the fragment ratio be x . From the factor $\frac{1}{L-1}$, if there is a value $L > 1$ such that $L = 1$, then all moments with $k > 0$ will become infinite. Provided reasonable analyticity conditions hold, it follows that the value is a pole of m . Taking the inverse Mellin transform of m then allows us to get $p(l)$ and, using the existence of the pole at $L = 1$, this immediately predicts that $p(l) \sim l^{-(1+\alpha)}$ is a power law distribution. This model was used by Ouyllon et al. [1996] to account for the power law distribution of joint spacing in a granitic massif in Saudi Arabia, with exponent $\alpha = 1/2$. Note that block sizes are proportional to spacing taken to the second power, which then predicts $\alpha = 1/2 = 1/4$, which is in agreement with our best estimate $\alpha = 0.25$ for the plate area distribution. This agreement between two very different systems may be fortuitous but is worth keeping in mind, especially in view of our claim of a continuity of the physics of fragmentation from the large plates to the blocks. We note also that the fragmentation model of [Cheng and Redner, 1988] predicts generically log-periodic corrections to the main power law since, for a large variety of $b(x)$, there are solutions of $L = 1$ with complex exponents. This has been shown to signal the existence of discrete scale invariance [Ouyllon et al., 1996]. It is tempting to interpret in this way the three approximately regular oscillations observed in figure 1, with a preferred scaling ratio approximately equal to 10. This value is in good agreement with the square of the preferred scaling ratio 3.5 found in the discrete hierarchy of linear sizes in fracture of rocks by Sadovskiy et al. [1984].

Our present conclusion on a universal fractal char-

acter of plate fragmentation does not prevent the existence of some underlying distinctions. For instance, P. Bird (private communication and [Bird, 2001]) proposes three main tectonic origins for plates: plume-assisted rifting for the larger plates (ranks 1-8 in Figs. 1 and 2), continental collision stress for the intermediate plate sizes (ranks 8-20), and back-arc spreading for the smallest plates (ranks 21-42). In Figs. 1 and 2, one can discern slightly different patterns of the tail behavior of these three subgroups. However, any formal statistical analysis of these possible distinctions would be, to our opinion, excessive due to the extreme smallness of the data set.

Acknowledgments. We acknowledge stimulating discussions with P. Bird and Y.Y. Kagan and thank P. Bird warmly for sharing his data.

References

- Bird, P., and R.W. Rosenstock, Kinematics of present crust and mantle flow in southern California, *Geol. Soc. Am. Bull.*, 95, 946-957, 1984.
- Bird, P., Plate Boundary Model PB2001, in press, 2002.
- Cheng, Z., and S. Redner, Scaling theory of fragmentation, *Phys. Rev. Lett.*, 60, 2450-2453, 1988.
- Gelfand, I.M., Guberman, Sh.A., Keilis-Borok, V.I., Knopov, L., Press, F., Ranzman, E.Ya., Rotwain, I.M., and A.M. Sadovsky, Pattern recognition applied to earthquake epicenters in California, *Physics of the Earth and Planetary Interiors*, 11, 227-283, 1976.
- Gorshkov, A.I., Kossobokov, V.G., Ranzman, E.Ya. and Soloviev, A.A., Recognition of earthquake prone areas: validity of results obtained from 1972 to 2000, *Computational Seismology*, 32, 48-57, 2001.
- Kolmogorov, A.N., On the logarithmically normal law of distribution of the size of particles under pulverisation, *Dokl. Akad. Nauk. SSSR* 31, 99-101, 1941.
- Ouillon, G., C. Castaing and D. Sornette, Hierarchical scaling of faulting, *J. Geophys. Res.*, 101, 5477-5487, 1996.
- Ouillon, G., D. Sornette, A. Gentier and C. Castaing, The imaginary part of rock jointing, *J. Phys. France*, 6, 1127-1139, 1996.
- Rao, C., Linear statistical inference and its applications, Wiley, N.Y., 522 p. (Chapter 6, Section 6e.3), 1965.
- Sadovskiy, M.A., T.V. Golubeva, V.F. Pisarenko and M.G. Shnirman, Characteristic dimensions of rock and hierarchical properties of seismicity, *Izvestiya Earth Physics*, 20, 87-96, 1984.
- Scholz C.H. and Mandelbrot B.B., eds., *Fractals in Geophysics*, Birkhäuser, 1989.
- Sornette, D., L. Knopov, Y.Y. Kagan and C. Vanneste, Rank-ordering statistics of extreme events : application to the distribution of large earthquakes", *J. Geophys. Res.* 101, 13883-13893, 1996.

D. Somette, Laboratoire de Physique de la Matière Condensée, CNRS UMR 6622 and Université de Nice-Sophia Antipolis, Parc Valrose, 06108 Nice, France, and Department of Earth and Space Sciences and Institute of Geophysics and Planetary Physics, University of California, Los Angeles, California 90095-1567. (email: somette@unice.fr)

V.F. Pisarenko, International Institute of Earthquake Prediction Theory and Mathematical Geophysics, Russian Ac. Sci. W arshavskoye sh., 79, kor. 2, M oscow 113556, Russia. (email: vlad@sirius.m itp.ru)

(Received X X X ; revised X X X ;
accepted X X X .)




Figure 1. Complementary cumulative distribution of the areas of tectonic plates (open circles) compared to the fit with the formula (3) for a power law (central long-dashed line) with exponent $\gamma = 0.25$ and $a = 0.002$. The small-dashed line and medium-dashed line provide a confidence interval defined in the text.




Figure 2. Medians $M_1; \dots; M_{42}$ (continuous line) and their corridor at the 95% confidence level delimited by the two dotted lines of the conditional Pareto distribution $(x_1; \dots; x_{42})$ given by (5) compared with the empirical cumulative distribution of the 42 plate areas (circles linked by straight segments), documented by Bird [2002].

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS

SORNETTE AND PISARENKO: FRACTAL PLATE TECTONICS