

Generalized contact process on random environments

Gyorgy Szabo¹, Hajnalka Gergely², and Beata Oborny²

¹Research Institute for Technical Physics and Materials Science
P.O. Box 49, H-1525 Budapest, Hungary

²Department of Plant Taxonomy and Ecology, Eötvös University
H-1117 Budapest, Pazmany Peter setany 1/C, Hungary

November 10, 2021

Spreading from a seed is studied by Monte Carlo simulation on a square lattice with two types of sites affecting the rates of birth and death. These systems exhibit a critical transition between survival and extinction. For time-dependent background, this transition is equivalent to those found in homogeneous systems (i.e. to directed percolation). For frozen backgrounds, the appearance of Gri ths phase prevents the accurate analysis of this transition. For long times in the subcritical region, spreading remains localized in compact (rather than ramified) patches, and the average number of occupied sites increases logarithmically in the surviving trials.

PACS numbers: 02.50.-r, 05.50.+q, 87.23.Cc

I. INTRODUCTION

Contact process (CP) was introduced by Harris [1] to describe competition between death and birth events for a spatially distributed species. In this model, the organisms localized at the sites of a lattice can die with a death rate or can create additional offspring in one of their (empty) neighboring sites. When increasing the ratio of birth to death rates, this system exhibits a critical phase transition from empty to the active state as the system size approaches infinity. This transition belongs to the so-called directed percolation (DP) universality class [2,3]. Similar transitions have been observed for some other, related models (for recent reviews see the Refs. [4,5]).

The above mentioned CPs applied homogeneous backgrounds. Numerical investigations on two-dimensional random environments have been restricted to diluted lattices [6-10]. Noest [6] have shown that, although the transition remains continuous, the extinction process is modified drastically on the randomly diluted lattice above the percolation threshold. Recently Dickman and Moreira [9] have reported that the scaling (a fundamental feature of critical transitions) is violated in this case. According to a field-theoretic analysis, similar result is predicted by Janssen [11].

Below the percolation threshold on a diluted lattice, the available sites form isolated (finite) clusters. In any particular cluster the species is expected to die out within finite time and the cluster remains empty afterwards. A simple mathematical description suggested by Noest [7] indicates that the average concentration of species vanishes algebraically with time and this phenomenon is analogous to the relaxation observed for the "Gri ths phase" in disordered spin models [12].

Field-theoretic arguments [5] support that the temporally quenched disorder itself, as well as the spatially quenched disorder, disturbs crucially the DP transition.

However, DP transition is expected on large scales for those random environments where the spatio-temporal disorder is uncorrelated. For two-dimensional systems this expectation is supported only by a small number of numerical evidences. For example, in an evolutionary game, extinction of one of three competing strategies represents a DP transition on a background whose time dependence is maintained by the competition of the remaining two strategies [13].

In the present paper, we study a generalized CP on a square lattice with two types of sites that provide different conditions for survival and reproduction for a hypothetical species. Our analysis is not restricted to frozen backgrounds. We give numerical evidence that the extinction process becomes analogous to DP transition if the random environment involves uncorrelated time dependence. Using numerical simulations, we investigate the main characteristics of spreading when initially there is only a single individual in the system. This method was suggested by Grassberger and de la Torre [14] to study DP transition (for homogeneous background) in the close vicinity of the critical point. Moreira and Dickman [8] have demonstrated that this technique is also efficient for the investigation of inhomogeneous systems. Now we have adopted this method to study the above-mentioned generalized CP in random environments. The results confirm the theoretical expectations [4,5,11,15] as well as the previous observations based on MC simulations [6-10]. We have also studied some geometrical features of the patches formed by occupied sites in the subcritical region.

II. THE MODEL

Space is represented by a square lattice. Each lattice site, $r = (x; y)$ (x and y are integers), is assumed to represent a microhabitat for one individual of a species. Quality

of a site, $g(r)$, determines the probabilities of survival and reproduction of the inhabitant individual. We assume two microhabitat types: good and bad (where $g(r) = 1$ or 0 , respectively). Site qualities are distributed randomly in space. Initially, each site is chosen to be good (bad) with a probability P ($1 - P$). Distribution of individuals over this random background is described by a state variable $\langle r \rangle$ that is 1 for occupied and 0 for empty state. Time evolution of the system is governed by three elementary processes affecting the values of $\langle r \rangle$ and $g(r)$. For an occupied site, extinction (death) can occur with a death rate $d[g(r)]$. For an empty site, colonization (birth) can take place with a birth rate $b[g(r)]$, provided that a randomly chosen nearest-neighbor site is occupied. It is supposed that good sites provide better conditions for living, that is $b(1) > b(0)$ and $d(1) < d(0)$. Furthermore, redrawing the value of $g(r)$ we change the environment with a rate f . The new state is independent of the actual values of $g(r)$ and $\langle r \rangle$. This occasional modification results in stationary distribution of good or bad sites over time, with probabilities P and $1 - P$, respectively, and leaves their distribution uncorrelated.

In the special case when every site is good ($P = 1$), the present model is equivalent to the well investigated contact process on homogeneous background [4,5]. Previous studies have shown that in this case, the species dies out if $\beta = b(1)/d(1) < c_c = 1.6488(1)$. Concentration of population vanishes algebraically in the active stationary state, that is $c \propto (\beta - c_c)^{\nu}$ where $\nu = 0.575(3)$ if $\beta > c_c$ [4,5,16,17]. This critical transition is accompanied with diverging fluctuations and correlation length.

The conditions of CP on diluted lattice can be reproduced by allowing the species to stay only in good sites ($b(0) = 0$ and $d(0) = 1$) for $P < 1$ and $f = 0$.

A mean-field approximation for frozen backgrounds ($f = 0$) predicts different concentration of species on good and bad sites: c_1 and c_0 , respectively. Neglecting the details of a straightforward calculation, the concentrations obey the following expressions:

$$c_1 = \frac{1C}{1 + (1-P)C} \quad \text{and} \quad c_0 = \frac{0C}{1 + (1-P)C}; \quad (1)$$

where $\beta = b(1)/d(1)$ ($\beta = 0$ or 1), and the average concentration, $c = P c_1 + (1 - P) c_0$ is obtained as

$$c = \frac{P}{A + \frac{P}{A^2 + 4(1-P)(\beta - 1)}} \quad (2)$$

where $A = \beta - 1$ and $\beta_{av} = P\beta + (1 - P)0$. This mean-field approximation predicts that concentration vanishes linearly when $\beta_{av} \rightarrow 1$. More precisely, for low concentrations the above expression can be approximated as

$$c = \frac{P(1 + (1 - P)0)}{\beta_{av} + 1} \quad (3)$$

if $\beta_{av} > 1$. Obviously, the trivial ($c = 0$) solution will be stable for $\beta_{av} < 1$.

In the limit $f \rightarrow 1$ death and birth occur with averaged rates: $d = P d(1) + (1 - P) d(0)$ and $b = P b(1) + (1 - P) b(0)$, respectively. Thus the system becomes equivalent to a homogeneous system (where $c_0 = c_1 = c$), and the average concentration obeys the traditional form [4,5]:

$$c = \frac{1}{1 + \beta} \quad (4)$$

if $\beta = b/d > 1$ and $c = 0$ otherwise. The difference between the Eqs. (2) and (4) refers to the f dependence.

III. SIMULATION OF SPREADING

The Monte Carlo (MC) simulations are performed on a lattice with $L \times L$ sites under periodic boundary conditions. Following the method suggested by Grassberger and de la Torre [14] spreading is investigated by averaging over many trials M when the initial state is close to the absorbing state. More precisely, each run is started with a single individual [at position $r = (0; 0)$] on a new, random, uncorrelated background. We have determined the survival probability

$$S(t) = \langle n(t) \rangle_i; \quad (5)$$

where the number of individuals at a given time t is defined as

$$n(t) = \sum_r \langle r \rangle; \quad (6)$$

and $\langle \cdot \rangle_i$ means the average over M trials and $\langle z \rangle = 1$ (0) for $z > 0$ ($z = 0$). We have also evaluated the average number of surviving individuals,

$$N(t) = \langle n(t) \rangle_i; \quad (7)$$

and the mean-square distance of individuals from the origin,

$$R^2(t) = \frac{1}{N(t)} \sum_r \langle r^2 \rangle; \quad (8)$$

M is varied from 10^4 to 10^8 in the systematic investigations. The system size L is adjusted to exceed significantly the average radius of occupied sites, namely, $L > 15R(t_{max})$, where t_{max} indicates the time limit. As usual, time is measured in Monte Carlo Steps (MCS). During one time unit, each site has a chance to modify the value of $\langle r \rangle$ (approximately once on average).

The efficiency of these simulations can be greatly improved by labeling the individuals, recording their coordinates, and picking up one individual randomly for updating the state of occupancy [4]. In the present case, a chosen individual residing at site r dies with a probability $d(g(r)) = (b(1) + d(0))$ or creates an offspring on

one of its nearest-neighboring sites r^0 with a probability $b(g(r^0)) = (b(1) + d(0))$. These attempted events take time $t = 1/n$ on average, therefore time is increased by t . Variation of the background can be handled in the same way within a smaller $(1 - l)$ region whose sites affect the CP at the given time. Namely, the values of $g(r)$ are redrawn at randomly chosen l^2 sites.

The average values defined above [see Eqs. (5)–(8)] are determined for discrete time values chosen equidistantly in a logarithmic scale.

IV. RESULTS FOR FROZEN BACKGROUND

First we concentrate on a system where the random background is frozen, i.e. $f = 0$. Varying the proportion of good sites (P), we have made a systematic MC analysis for fixed death and birth rates. For the present parameters (see Fig. 1) the good sites form isolated clusters [having $P < P_{\text{perc}} = 0.5926$ (percolation threshold)] and infinite spreading is possible across the bad sites. The results summarized in the subsequent figures agree quantitatively with those found by Moreira and Dickman [8] on a diluted lattice for $P > P_{\text{perc}}$.




FIG. 1. Survival probabilities as functions of time at different P values (indicated by labels) for $b(1) = 1$, $b(0) = 0.5$, $d(1) = 0.25$, and $d(0) = 0.5$.

Figure 1 shows some typical time-dependence functions of survival in a log-log plot. The statistical errors are comparable to the line width due to the large number of trials (e.g. $M = 10^8$ for $P = 0.22$). In the active region ($P > P_c$), $S(t)$ tends to a constant value as expected [14]. In the subcritical region, $S(t)$ can be well approximated by a power law. Unfortunately, convergency toward the limit value becomes extremely slow in the vicinity of the critical point, and this prevents the accurate determination of P_c . Division between the active and inactive regions becomes more visible when considering the function $N(t)$ (see Fig. 2).




FIG. 2. Average number of individuals vs. time for the same simulations as in Fig. 1.

In the active region, the surviving individuals occupy a compact patch whose radius increases linearly with time. Consequently, $N(t)$ becomes proportional to t^2 for sufficiently long times as indicated in the Fig. 2. By contrast, when $P < P_c$, then $N(t)$ tends to a power law decay with an exponent depending on P .




FIG. 3. Log-log plot of mean-square distance of individuals from the origin as a function of time. Parameter values are the same as in Fig. 1.

Figure 3 shows the time evolution of the mean square distance of surviving individuals as defined by Eq. (8). In agreement with the expectations $R^2(t) / t^2$ for sufficiently long times if $P > P_c$. In the subcritical region, the increase of $R^2(t)$ becomes significantly slower than that found for the homogeneous system. In the homogeneous system, the surviving individuals perform random walks independently of each other, therefore $R^2(t) / t$ in the subcritical region. In the present case, however, our

data indicate a power law increase: $R^2(t) / t$, where the exponent $(P) < 1$. This indicates localization of spreading as, it will be explained in Section VI.

Evaluation of $N(t)$ involves those (stopped) trials which have reached the empty state before the given time t . Consequently, the ratio $N(t)=S(t)$ expresses the average number of individuals in the surviving trials. In the subcritical region, the time-dependence of this quantity tends to logarithmic increase as shown in a lin-log plot (see Fig. 4).




FIG. 4. Time dependence of the average number of individuals in the surviving trials at $P = 0.22, 0.23$, and 0.235 .

In the classical theory of percolation (reviewed in Refs. [18,19]), a geometrical feature of the clusters of size s is characterized by an average cluster radius $R_{\text{perc}}(s)$. Below the percolation threshold $R_{\text{perc}}(s) / s$ for the large s limit. According to MC simulations [18,20,21], $\nu = 2/3$, while the "self-avoiding walk" prediction [22] gives $\nu = 3/4$. In general, $\nu > 1/2$ values are characteristic for ramified clusters [18]. On the contrary, above the percolation threshold the finite size clusters are compact, that yields $\nu = 1/2$.

Following this analogy, we examine the relation between $N(t)=S(t)$, the actual size of population in the surviving trials, and the average radius of the area they occupy at a given time in the subcritical region. Figure 5 shows this relation on a log-log plot. In the close vicinity of the critical point (at $P = 0.239$) we find power law behavior with an exponent $1.66(6)$. This behavior indicates that the species spreads over a ramified cluster in the whole time interval we could study the system.

In the subcritical region, however, we can observe two different regions. In the transient (short-time) region, the increase of $R^2(t)$ is faster than at the critical point. This indicates that the offspring move away from the origin along preferred paths formed by good sites. Conversely, the long-time behavior is dominated by those trials where the individuals stay on a compact patch.




FIG. 5. $R^2(t)$ vs. $N(t)=S(t)$ for $P = 0.22$ (dashed-dotted), 0.23 (dashed), 0.235 (solid), and 0.239 (dotted line).

In the next section we show that these features fundamentally change when we allow the background change over time.

V. RESULTS FOR TIME-DEPENDENT BACKGROUND

Applying time-dependent backgrounds, we determined the same quantities $[S(t), N(t), \text{ and } R^2(t)]$ as before. We use the same birth and death rates when varying the value of P for a fixed rate (f) of background change. Our results estimate the critical value for the ratio of good sites at $P_c = 0.2680(1)$.




FIG. 6. Log-log plot of survival probabilities versus time at different P values (as indicated by labels) for $b(1) = 1$, $b(0) = 0.5$, $d(1) = 0.25$, $d(0) = 0.5$, and $f = 0.01$.

Figure 6 shows that the survival probability $S(t)$ vanishes exponentially when $P < P_c$. In the active region

($P > P_c$), however, $S(t)$ tends to a constant value. Notice, that the convergence is significantly faster here than at the frozen background (see Fig. 1).

Data for $P = 0.268$ represent the behavior of the system at the critical point. A numerical analysis confirms that the survival probabilities can be well approximated by a power law, $S(t) / t$ for sufficiently long times. Numerical fitting gives $\alpha = 0.45(2)$ in agreement with the exponent found in DP transitions in homogeneous systems [4,5].

When considering $N(t)$, we can also distinguish between three different behaviors (see Fig. 7). Above the critical point ($P > P_c$), $N(t)$ tends towards quadratic time dependence. Below the critical point, $N(t)$ decreases exponentially. At the critical point, however, our MC data indicate power law behavior, i.e., $N(t) / t$ with $\alpha = 0.23(1)$. This numerical value of α agrees with the characteristic exponent of DP transitions [4,5].




FIG. 7. Average number of individuals vs. time for the same simulations as in Fig. 6.

In summary, all the above-mentioned numerical results support that contact process exhibits the same features in time-dependent random environments as in homogeneous background. Similarly to $S(t)$ and $N(t)$, $R^2(t)$ also supports this conclusion (data not shown).

These results have been gained at a fixed rate of changes ($f = 0.01$). Preliminary results suggest, however, that the same conclusions hold for other values of f as well. General features of spreading remain unchanged due to the robustness of DP transition.

Here it is worth mentioning that we have also studied the concentration of individuals (as well as its fluctuation) in the stationary state for some parameter values. These analyses support that the variation of both quantities is consistent with the expectations, i.e., the species concentration and its fluctuation exhibit power law behavior with exponents characteristic for the two-dimensional DP systems [16,17,4,5].

VI. CONCLUSIONS AND SUMMARY

We have studied a generalized version of contact process on a square lattice. The background in the present model provides a continuous transition from a homogeneous system to a randomly fluctuating environment, including (frozen) diluted lattices. The environment is inhomogeneous, consisting of two types of sites (good and bad) that affect the rates of birth and death of the species. In this model, spreading from a cluster of good sites to another one is permitted across the bad sites and/or bridges (of good sites) created occasionally by background fluctuations.

Our analyses have been restricted to the consideration of some typical features, because the simulations are rather time-consuming. The results confirm the theoretical expectations as well as the general picture drawn by previous authors [6-9] who considered CP on diluted lattices.

Obviously, an inhomogeneous environment contains areas that provide better (or worse) conditions for the survival relative to the average quality of the habitat. This variation becomes particularly important in the vicinity of the transition point that separates the active and inactive regions for long-time behavior. In the subcritical region, the surviving individuals are constrained to remain within isolated patches. Conversely, in the active region, favorable areas can form a percolating cluster, that has infinite extension, and sustains infinite survival.

To explain the main results, we briefly remind the reader of a simple calculation suggested by Noest [7]. In this description we characterize the favorable patches by size s and assume that their probability decreases exponentially, i.e., n_s / e^{-As} . Furthermore, it is also assumed that the average over many trials (or patches of size s) yields an exponentially decreasing number of individuals over time, that is, $m_s(t) / e^{-t/\tau_s}$ where the average survival time can be approximated as τ_s / e^{Bs} if initially all sites are occupied. Noest has shown that long-time behavior can be well approximated in this case by a power law decrease, that is, $m(t) = \sum_s n_s m_s(t) / t^{A=B}$. This prediction is derived from a maximum likelihood estimation that implies the existence of a typical cluster size giving the dominant contribution to $m(t)$. According to this approach, the size of dominant clusters increases logarithmically with time t .

The adoption of this approach is not straightforward to the present situation where we consider the spreading from a single seed. In our case, the species can die out before achieving homogeneous distribution, while homogeneous initial state is assumed in the former calculation. We can assume, however, that the species spreads over the whole cluster or dies out within a transient time, and the probabilities of these outcomes are independent of cluster size. If this hypothesis holds, then the approach of Noest can be applied for the analysis of long-time be-

havior.

Some predictions of this rough method have been confirmed by our MC simulations. Namely, $S(t)$ and $N(t)$ exhibit a dominant power law decrease in the subcritical region, while $N(t)=S(t)$ (in proportion to the dominant cluster size) increases logarithmically. Furthermore, in the active region, the algebraically decaying contribution of $S(t)$ comes from extinction on (compact) finite size clusters whose probability distribution satisfies the above assumption [21,19].

Noest's approach is focused on compact clusters where s/R^2 . Consequently, $N(t)=S(t)/R^2(t)$ is expected when the system behavior is dominated by a typical cluster size. This behavior can be observed for long times far below the critical point. Figure 5 indicates, however, that these predictions (and assumptions) fail in the close vicinity of the critical point. In other words, the ramified clusters give relevant contribution to $S(t)$ and $N(t)$ during a transient time that increases when approaching the critical point. In this case, we need a more sophisticated description allowing the emergence of ramified clusters and the shape-dependence of s .

Numerical analysis of the extinction process is difficult (in a quenched environment) because decrease of $S(t)$ and $N(t)$ slows down in the vicinity of the critical point. In practice this implies persistence over long time periods before extinction. The slow extinction process, however, requires rather specific environmental conditions. Even a small degree of temporal fluctuation results in an extinction process that is analogous to DP transition on large scales. Our mean-field calculations and preliminary simulations have indicated that the transition point is strongly affected by the rate of background change. Further analyses are required to quantify this phenomenon for different birth and death rates.

ACKNOWLEDGMENTS

Support from the Hungarian National Research Fund (T-33098) is acknowledged.

[1] T. E. Harris, Ann. Prob. 2, 969 (1974).
 [2] H. K. Janssen, Z. Phys. B 42, 151 (1981).
 [3] P. Grassberger, Z. Phys. 47, 365 (1982).
 [4] J. Marro and R. Dickman, Nonequilibrium Phase Transitions in Lattice Models, (Cambridge University Press, Cambridge USA, 1999).
 [5] H. Hinrichsen, Adv. Phys. 49, 815 (2000).
 [6] A. J. Noest, Phys. Rev. Lett. 57, 90 (1986).
 [7] A. J. Noest, Phys. Rev. B 38, 2715 (1988).
 [8] A. G. Moreira and R. Dickman, Phys. Rev. E 54, R3090 (1996).

[9] R. Dickman and A. G. Moreira, Phys. Rev. E 57, 1263 (1998).
 [10] T. Tao, K. Tainaka, and H. Nishimori, J. Phys. Soc. Jap. 68, 326 (1999).
 [11] H. K. Janssen, H. K. (1997). Phys. Rev. E 55, 6253 (1997).
 [12] R. B. Griffiths, Phys. Rev. Lett. 23, 17 (1969).
 [13] G. Szabo, T. Antal, P. Szabo, and M. Droz, Phys. Rev. E, 62, 1095 (2000).
 [14] P. Grassberger and A. de la Torre, Ann. Phys. (N.Y.) 122, 373 (1979).
 [15] M. C. Marques, J. Phys. A 23, 3389 (1990).
 [16] J. Adler and J. A. M. Duarte, Phys. Rev. B 35, 7046 (1987).
 [17] R. C. Brower, M. A. Fuman, and M. Moshe, Phys. Lett. 76B, 213 (1978).
 [18] D. Staufer, Phys. Rep. 54, 1 (1979).
 [19] J. W. Essam, Rep. Prog. Phys. 43, 833 (1980).
 [20] P. L. Leath, Phys. Rev. Lett. 36, 921 (1976); Phys. Rev. B 14, 5046 (1976).
 [21] D. Staufer, Phys. Rev. Lett. 41, 1333 (1978).
 [22] D. Staufer, Z. Phys. B 30, 173 (1978).