

# New approach to the superconductivity problem

I. M. Yurin and V. B. Kalinin

Institute of Physical Chemistry, Leninskiy prosp. 31, GSP-1, Moscow, 119991, Russia

In this work, a question is tackled concerning the formation of a superconducting condensate in an earlier proposed model of "elastic jelly", in which phonons of the ion system play the part of initiating ones. It was shown that in distinction from the BCS theory, the momenta of forming electron pairs are different from zero. This fact changes the pattern of the description of the superconductivity phenomenon in the proposed model. First, the gap in the one-electron spectrum appears due to the effect of a "mean field" on the energy of electron state from the side of occupied states, the nearest neighbors over the momenta grid. Second, the condensate is formed by electron states with energies below that of the gap edge, this is why the Fermi-condensation arises in the system. Third, anomalous expectation values are strictly equal to zero in the proposed model.

PACS number(s): 74.20.Mn, 74.72.-h

## I. INTRODUCTION

A considerable number of fundamental works have been devoted to tackling the superconductivity (SC) problem and, in particular, a high temperature SC one (HTSC). The opinion on the present state of the problem may be formed considering the works [1-3]. Most relevant publication are to some extent associated with the BCS theory [4]. Now, for metals enumerated in the Periodic System of elements (PSE), the SC theory is considered to be more or less completed and the validity of the BCS theory leaves no room for doubt on the part of the majority of authors. At the same time in [5-8] a conclusion was drawn that the BCS theory may be not the only possible way to explain the phenomenon of SC.

Outstanding results [9] obtained by Bednorz J.G. and Müller K.A. in 1986 for  $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$  caused the great sensation among chemists, physicists and material researchers. Since that time the highest value for  $T_s$  at about 164K was observed in  $\text{HgC}_2\text{Ba}_2\text{Cu}_3\text{O}_{8+x}$  ceramics.

There are grounds available highlighting a possibility of observing high  $T_s$  in intercalates and nanotubes [3]. In our opinion, an increase in the value of  $T_s$  and the observation of HTSC are very plausible in high-pressure (HP) phases. Almost twenty-five years ago, SC was reported to have been observed in  $\text{AuGa}_2$  [10]. Comparatively high values of  $T_s$  were observed in fullerenes, HP phases with respect to graphite, and their derivatives fullerides  $\text{C}_{60}\text{M}$  [11,12], where M stands for K, Rb, Cs. The idea of obtaining HTSC in HP phases of  $\text{Li}_3\text{P}$  or  $\text{Li}_3\text{N}$  is at present at the stage of being accepted [13,14]. In the last publications on this topic, SC is investigated in a HP phase of  $\text{MgB}_2$  [15-17]. In conjunction with the problem under consideration, we also can't help mentioning a hypothetical phase of metallic hydrogen [18] and nitrogen (a private communication).

The obtained results, especially the identification of HTSC with the values of  $T_s > 100\text{K}$ , are rather inconsistent with both the predictions of the BCS theory proper [19], which had been made before discovering HTSC, and any of its complementary modifications, among which the theory of bipolarons has to be singled out [2]. In connection with this, we note that most HTSC materials in stoichiometric phase are, strictly speaking,

dielectrics, whose conductivity is fully determined by low concentrations of electroactive defects, whereas the theory of bipolarons gives low values of  $T_s$  as the Fermi level approaches the edges of zones. Thus, it does not explain HTSC for the aforementioned class of compounds.

In Russia, one usually considers the results of microscopic calculations are in good agreement with the predictions of the BCS theory, including the case of HTSC [20] as well. There is also another viewpoint which is doubtful about the reliability of any one calculation performed by and large within the scope of the zone theory [21].

The problem is that a lot of poorly substantiated approximations did appear for the years of the existence of the zone theory. Among these, for instance, the procedure of accounting for the screening effects seems to be quite harmless. These approximations were introduced by virtue of quite relevant reasons associated with an insufficient quick operation of computer equipment to perform strict calculations of the zone structure of metals, including that of the Periodic System. The application of these procedures, however, renders the calculation of the electron-phonon interaction (EPI) constants quite an unpredictable operation, depending only on theoretical preferences of a specialist in the field of zone computations.

From our viewpoint, the main source of difficulties consists in that all the calculations associated with EPI pursue the goal of reducing the Hamiltonian of the system to the Frohlich form. It has been known for a long time [19] that the Frohlich model does not allow a consistent accounting for the electron-electron interaction (EEI), because the Coulomb interaction cannot still be reasonably separated into that already accounted for in the initial values of Frohlich Hamiltonian constants, and that which will thereafter show up in the renormalization of the initial phonons owing to EPI. Accordingly, all the microscopic calculations of the EPI constants are, strictly speaking, inconsistent.

Just these considerations underlied the need of a revision of the fundamentals of the SC theory, which was undertaken in studies [22-24]. It was shown the attraction between electrons exists in a long-wave range due to an exchange of virtual phonons. This question was not discussed in earlier works (for example, [25]) devoted to the problem being tackled. We believe this was due to the following two basic causes.

First, in the framework of "procedure to account for the screening effects" the function of the permittivity of electronic plasma was incorrectly used for the removal of "undesirable" singularities from the matrix elements of EPI in the long-wave limit. Second, the simplified form of the effective EEI used in study [25] did not suppose substantial differences in the electron interaction both in the vicinity of the Fermi surface and far off it.

The authors of [25] well understood the disadvantages of the aforementioned approaches. Remind that in [19] serious forebodings were caused just by these approximations, which gave rise to a detailed discussion. Moreover, an idea was put forward that in a further development of the "jelly" model one should have refrained from the use of ion-plasma oscillations as initiating phonons.

The accounting for just these remarks underlied the investigation in the long-wave range of EEI, which had been undertaken in our studies [22-24], where the elastic oscillations of the ion system serve as initiating phonons. It was shown the adequate inclusion of the phonon-phonon interaction in the examination process enables one to avoid the application of poorly substantiated procedures to "account for the screening effects". And finally, the EEI potential was calculated without any limitations as to its form.

It was shown that the unitary transformation effected in the framework of the model suggested in [23,24]

can reduce the Hamiltonian of the electronic system of an infinite monoatomic metal to the following form ( $\hbar = 1$ ):

$$H_{\text{tot}} = \sum_p \sum_{\sigma} E_p C_{p\sigma}^\dagger C_{p\sigma} + H_{ee} \quad (1)$$

$$H_{ee} = \sum_p \sum_k \sum_q \sum_{\sigma} \sum_{\sigma'} U(p; k; q) C_{p+q\sigma}^\dagger C_{k\sigma}^\dagger C_{k\sigma} C_{p\sigma} \quad (2)$$

and for  $U(p; k; q)$  at  $p, k \ll K_F$  we have

$$U(p; k; q) = -4 \frac{zm}{3M} \frac{G_{ee}}{q^2} \frac{K_F^2}{q^2} \frac{K_F^2}{1 - q^2} \quad (3)$$

where  $E_p = \frac{p^2}{2m}$ ,  $G_{ee} = \frac{e^2}{4\pi\epsilon}$ ,  $e$  is the electron charge,  $C_{p\sigma}^\dagger$  and  $C_{p\sigma}$  are operators of the creation and annihilation of electrons with a momentum  $p$ , respectively;  $\sigma, \sigma'$  are spin indices,  $\epsilon$  is the permittivity of the ion system,  $M$  is the mass of ion,  $m$  is the zone mass of electron,  $z$  is the number of conductivity electrons per elementary cell;  $\epsilon = 4\pi m^2 S^2 \frac{zm}{3M} K_F^2$ ,  $1 - q^2 = 1 + \frac{zm}{3M} K_F^2 - q^2$ ,  $q^2 = (4e^2/\epsilon) m K_F$ ,  $K_F$  is the value of Fermi vector, and  $S$  is the observable sound velocity in the system.

The potential of the form (3) leads to an instability relative to the formation of electron pairs. In distinction from the BCS theory, the momenta of electron pairs are different from zero, while the pairs bonding energy  $E_b$  if a typical relationship  $\frac{E_b}{2m} \ll E_b$  for SC systems is fulfilled, may be estimated as follows:

$$E_b = \frac{zm}{M} \frac{G_{ee} K_F^4}{Q^3} \quad (4)$$

where  $Q = 1 - q^2$ .

It is obvious that  $E_b \propto M^{-1/2}$ , and this is consistent with the observations of an isotopic effect in a series of metals [26]. We also note that the coherence length  $l_{\text{coh}} \propto Q^{-1}$ .

## II. SC CONDENSATE IN THE MODEL OF ELASTIC JELLY

The further discussion will be carried out for the 3D system - a cubic crystal  $L \times L \times L$  with periodic boundary conditions.

Consider in the framework of Eqs. (1)–(3) a wave function of the bound state of an electron pair with maximum bonding energy (MBE). This state will obviously be the state with zero spin. With increasing  $S$ , its wave function will first become hydrogen-like, and then delta-like, so that at  $S \rightarrow 1$  it proves to be possible to derive for the state  $|R\rangle$  of an electron pair with a momentum  $R$  at an accuracy of up to the normalizing factor the following expression:

$$|R\rangle = \sum_{p+k=R} \tilde{\psi}(k-p) C_{p\sigma}^\dagger C_{\#k\sigma}^\dagger |0\rangle \quad (5)$$

where  $\tilde{\psi}(k-p)$  is a discrete delta-shaped function, which is determined as follows:

$$\tilde{\psi}(r) = \begin{cases} \frac{8}{L^3} < 1; & \text{if } |r| \leq \frac{L}{2}; \\ 0 & \text{in the rest cases.} \end{cases} \quad (6)$$

Hence, it follows that in finite crystal, at least in the limit under consideration  $S \rightarrow \infty$ , the following Hamiltonian (7) well describes the statistics and energy spectrum of MBE-states:

$$H = \sum_p E_p \hat{n}_p - E_b \sum_p \hat{n}_p \hat{n}_{\#p} - \frac{E_b}{4} \sum_{p \neq k} \tilde{n}(p-k) \hat{n}_p \hat{n}_k + \frac{E_b}{4} \sum_{p \neq k} \tilde{n}(p-k) \hat{C}_p^y \hat{C}_{\#k}^y \hat{C}_k^y \hat{C}_{\#p}^y \quad (7)$$

where  $\hat{n}_p = \hat{C}_p^y \hat{C}_p$  and the operators  $\hat{C}_p^y$  and  $\hat{C}_p$  are suggested to be linked with operators  $C_p^y$  and  $C_p$  from Eq. (1) by the unitary transformation, whose derivation is postponed for further publication. Obviously the wave function of the ground state of the Hamiltonian (7) expressed in terms of operators  $\hat{C}_p^y$  does not differ from the analogous one in the case of normal metal:  $|\psi_0\rangle = \prod_{k < K_F} \hat{C}_k^y \hat{C}_{\#k}^y$ .

The quasi-particle energy  $\tilde{E}_k$  may be determined in random phase approximation from the equation  $H |\hat{C}_k^y \psi_0\rangle = \tilde{E}_k \hat{C}_k^y \psi_0$  and thus one can get:

$$\tilde{E}_k = E_k - 2\tilde{n}_k \epsilon_0 \quad (8)$$

where  $\epsilon_0 = 7E_b$  and

$$\tilde{n}_k = \frac{1}{14} \langle \hat{n}_k \rangle + \frac{1}{2} \sum_p \tilde{n}(p-k) \langle \hat{n}_p \rangle \quad (9)$$

For the large systems ( $L \rightarrow \infty$ ) in the absence of any spin order in the electron system we have  $\tilde{n}_k = \langle \hat{n}_k \rangle$ . The averaged occupation number  $\tilde{n}_k$  may be expressed through the quasi-particle energy  $\tilde{E}_k$  by means of Fermi-Dirac distribution function ( $k_B = 1$ ,  $\mu$  is the chemical potential of electron system):

$$\tilde{n}_k = \frac{1}{1 + \exp \frac{\tilde{E}_k - \mu}{T}} \quad (10)$$




FIG. 1. Temperature dependence of one-electron energy spectrum in the model of elastic jelly (schematic)

The evolution of a one-electron spectrum of the system with its temperature varying becomes obvious from Fig. 1. At high temperatures, this spectrum is described by an ordinary parabola; and at  $T = T_s$  the spectrum exhibits a peculiar feature which, at  $T < T_s$ , transforms into a gap. So, the SC condensate is formed by electrons with the energy  $E_k < 0$  at  $T < T_s$ .

Taking into account the Pauli principle, two versions of scatter are possible in the system at lowest temperatures. In the first version, the state of electron with momentum  $k > K_F$  or hole with  $k < K_F$  changes (electron and hole concentrations in this case  $\exp(-\epsilon_0/T)$ ). In the second version, an electron is removed from SC condensate with the formation of an electron-hole pair, or vice versa an electron-hole pair annihilates. The contributions of both channels to collision integrals are "frozen out" as temperature lowers according to the exponential law. This circumstance combined with that a shift of SC condensate as a whole in the momenta space leads to the formation of a state with nonzero current is convincing in that a SC transition is actually observed in the system under investigation.

The transition temperature  $T_s$  may be estimated by the equation  $\partial \sim \partial E_{\vec{k}} = 1$ . Then we derive for  $T_s$

$$T_s = \frac{0}{2} \quad (11)$$

This estimation well agrees with the well known relationship  $T_s = 2 \epsilon_0/3.5$ , coming from the BCS theory in the case of weak coupling. We suppose the deviations from the relationship (11) are due to correlation and many-particle effects.

The remaining bound, but not MBE pair states may be accounted for by inserting additional terms in the Hamiltonian (7). Yet this does not substantially distort the microscopic picture of the phenomenon by virtue of two interrelated causes. Firstly, these terms are small in their absolute values, compared with the bound energy of MBE-state, which has already been accounted for. Secondly, the state of the SC condensate is separated by an energy gap from the other states. Therefore, in considering the state of SC condensate the arisen terms may be accounted for in the framework of the perturbation theory without any substantial changes in its wave function.

The behaviour of SC condensate in the electromagnetic field may be described by the set of wave functions of MBE pair states due to the fact, that the unitary transformation, linking operators  $C^y$  and  $C^y$  involves these wave functions as transformation parameters. This circumstance is helpful for the explanation of flux quantization and Josephson effects in SC systems. The latter fact needs a detailed consideration and is postponed for further publication.

Finally, let consider the aforesaid limit  $S \rightarrow 1$ . The energies of the bound states of the starting Hamiltonian (1) are of an analytical character in relation to the parameter  $S$  throughout the whole range of its admissible values. This is why the analyticity of the energy parameters of the Hamiltonian (7) relative to the same parameter  $S$  cannot raise doubts. Then the presence of the delta-shaped terms in the Hamiltonian (7) for all the admissible values of  $S$  seems to be obvious.

### III. THE CONDITION FOR THE ONSET OF HTSC

A particular interest is spurred by the divergence of the bonding energy  $E_b$  at  $Q \rightarrow 0$ , following from Eq. (4). In real systems, this divergence should not reveal itself in the observed  $T_s$  by virtue of symmetry reasons put

forward in [23]. Simultaneously, with varying parameters of specimens, there should be observed a maximum of the transition temperatures if the following condition were fulfilled:

$$S^2 - \frac{zm}{3M} V_F^2 \quad (12)$$

This condition can be with a satisfactory accuracy fulfilled in systems with variable physical parameters, in which, provided Eq. (12) is obeyed, HTSC should be observed.

Let consider a semiconductor with a rather low concentration of electroactive defects, undergoing a phase transition with a change in volume. In both phases, the material is stable, and, as a rule,  $S^2 - \frac{zm}{3M} V_F^2$ . In unstable phases  $S^2 < 0$ . It is possible to assume HTSC is observed in nonequilibrium systems, where the phase transition is decelerated  $S^2 - \frac{zm}{3M} V_F^2 > 0$  (the diamond-graphite transition is not realized exclusively due to kinetic reasons).

One may expect the fulfillment of relationship (12) with good accuracy in the vicinity of the boundaries of stability of HP phases. From this standpoint, it seems to be very promising to investigate the kinetics of the phase transitions at HP with a view to searching for new HTSC materials.

#### IV. CONCLUSIONS

The basic qualitative differences of the BCS theory from the suggested SC model are listed below.

1. In accordance with the BCS theory, the SC condensate is formed by the pairs of electrons with zero momenta, and is therefore called Bose condensate.

In the proposed theory, the SC condensate is formed by electrons, and, accordingly, transforms into the Fermi condensate.

2. In the BCS theory, the ground state of the system is obtained through the superposition of the states with different total numbers of electrons. It is this property that underlies one of the formulations of the BCS theory [27] using anomalous expectation values  $\langle \psi^\dagger \psi_i \rangle$ , where  $\psi^\dagger$  are electron creation operators, as order parameters.

Now, in the proposed theory these expectation values are strictly equal to zero.

3. It is easy to show each of the states with a fixed total number of electrons, forming the superposition of the ground state in the BCS theory, is stationary state according to charge conservation law. The energies of all these states are the same value  $E_{min}$  due to the fact, that the wave function of the ground state  $|j\rangle$  satisfies the following equation  $H|j\rangle = E_{min}|j\rangle$ . Thus, a conclusion may be drawn that the removal of electrons from the superconductor does not change its energy. Hence, the work function of electrons in SC at lowest temperatures must be equal to zero.

Now, in the proposed model such a requirement is irrelevant.

Therefore, the investigation of the work function of electrons in SC materials seems to be a main practicable method for the experimental comparison of the theories.

## V . A C K N O W L E D G E M E N T S

We are grateful to Dr. N.V. K lassen, Dr. V.V. G rom ov, and Prof. Allan Solom on for their interest of our work and support of it.

---

- [1] J.G. Bednorz, K.A. M uller, "Perovskite-type O xides – the New Approaches to H igh – T<sub>c</sub> superconductivity" (The Nobel Foundation, New York, 1988).
- [2] A.S.A lexandrov, A.B. K rebs, U sp.F iz.N auk (Rus) 162 (5), 1 (1992).
- [3] V.L.G inzburg, U sp.F iz.N auk (Rus) 170 (6), 619 (2000); *ibid.* Conf.: M ajor Trends in Superconductivity in New M illenium , 31m .-6 apr. K lsters, Switzerland (2000).
- [4] J.Bardeen, L.N.Cooper, and J.R.Schrie er, Phys.Rev.106, 162 (1957); *ibid.* 108, 1175 (1957).
- [5] V.L.G inzburg, Progr.Low Tem p. Phys. 12, 1 (1959).
- [6] N.N.Bogolubov, JETP (Rus) 34, 41 (1958).
- [7] J.Valatin, Nuovo C im ento 7, 843 (1958).
- [8] C.K uper, Adv. Phys. 8 (29), 1 (1959).
- [9] J.G. Bednorz, K.A. M uller, Z.Phys.B 64, 189 (1986).
- [10] M .K .Palistrant, A.T.T rifan, Superconducting alloys and their properties, InstApplPhys., K ishinev, 1977.
- [11] V.H.C respi, Phys.Rev.B 60, 100 (1999).
- [12] W .Neuerberger, V.B.K alinin, Pat.USA accepted, (Appl.N 09/450.633 PCT 11/22/00).
- [13] V.B.K alinin, A.A.D onchak, A.M .G olubev, S.A K ulinich, K .P.Burdina, T.P.Puryaeva, V.D.Butskii, M aterialvedeniye (Rus) 2, 40 (1999).
- [14] V.B.K alinin and A.M .G olubev, M aterialvedeniye (Rus) 4, 26 (1999).
- [15] A.F.G oncharov, V.V.Stuzhkin, E.G regoryanz, J.Hu, R.J.H em ley, H o-kwangM ao, G .Lapertot, S.L.Bud'ko, P.C.C an eld, cond-m at/0104042.
- [16] S.L.Bud'ko, G .Lapertot, C.Petrovic, C.E.Cunningham , N.Anderson, P.C.C an eld, Phys.Rev.Lett. 86, 1877 (2001).
- [17] D.K.F innem ore, J.E.O stenson, S.L.Bud'ko, G .Lapertot, P.C.C an eld, Phys.Rev.Lett. 86, 2420 (2001);
- [18] E.G .M aksim ov, U sp.F iz.N auk (Rus) 170 (10), 1033 (2000).
- [19] Eds V.L.G inzburg, D.A.K irzhnits, Problem of H igh-Tem perature Superconductivity, M oscow , 1977; *ibid.*, H igh-Tem perature Superconductivity, New York: Consultants Bureau, 1982.

- [20] E. G. Maksimov, D. Y. Savrasov, S. Y. Savrasov, *Usp. Fiz. Nauk* (Rus) 40 (4), 337 (1997).
- [21] M. R. Norman, *J. Phys. Chem. Sol.* 54, 1165 (1993).
- [22] V. B. Kalinin, A. M. Golubev, I. P. Shilov, I. M. Yurin, *Conf.: Inst. Phys. Chem. at the Transition of Centuries*, Moscow, March, poster, 103, 2000.
- [23] I. M. Yurin, V. B. Kalinin, *Materialovedeniye* (Rus) 1, 2 (2001).
- [24] I. M. Yurin, V. B. Kalinin, *cond-mat/0102407*.
- [25] D. Bohm, D. Pines, *Phys. Rev.* 82, 625 (1951).
- [26] C. A. Reynolds, B. Serin, W. A. Wright, and L. B. Nesbitt, *Phys. Rev.* 78, 487 (1950).
- [27] G. M. Eliashberg, *JETP* (Rus) 38, 966 (1960); *ibid.*, 39, 1437 (1960).