A nalog of K elvin-H elm holtz instability on super uid liquid free surface

S.E.Korshunov¹

L.D. Landau Institute for Theoretical Physics RAS, Kosygina 2, 117940 Moscow, Russia

(April 19, 2002)

We analyse the analog of the Kelvin-Helm holtz instability on free surface of a super uid liquid. This instability is induced by the relative motion of super uid and norm alocom ponents of the same liquid along the surface. The instability threshold is found to be independent of the value of viscosity, but turns out to be lower than in absence of dissipation. The result is similar to that obtained for the interface between two sliding super uids (with di erent mechanism of dissipation) and con med by the rst experimental observation of the Kelvin-Helm holtz instability on the interface between ${}^{3}\text{He-A}$ and ${}^{3}\text{He-B}$ by B laauw geers et al. (cond-mat/0111343).

PACS num bers: 47.20 M a, 68.03 K n, 67.40 Pm

I. IN TRODUCTION

The K elvin-H elm holtz instability [1] is a dynam ic corrugation instability of the interface separating two liquids sliding with respect to each other. The concept of such instability has been orginally introduced when considering ideal liquids and in presence of dissipation becomes ill de ned, because a relative motion of two liquids in contact with each other is no longer a solution of the hydrodynam ic equations.

The simplest situation, when an equilibrium di erence in velocities can be maintained at a surface of a liquid, is the relativem otion of super uid and norm alcomponents (a counter ow) in super uid He. The corrugation instability of a super uid liquid free surface in presence of a counter ow along the surface was studied in Ref. [2] (in relation with experiments of Egolf et al. [3]). It can be considered as an example of the K elvin-Helm holtz instability, in which both liquids are located on the same side of the interface. A nalogous instability can appear when super uid ⁴He is sliding along the atom ically-rough interface separating it from solid ⁴He [4]. Such interface is known to allow for equilibrium melting and crystallization of ⁴He [5,6], and, as a consequence, its behavior resembles that of free surface of a liquid.

Recently the interest to surface instabilities in superiuds has been revived [7{9] in relation with experiments on laser manipulated Bose gases and the rst experimentalobservation of the K elvin-H elm holtz instability on the interface between two super uids,³He-A and ³He-B [10]. In particular, it has been demonstrated [9] that addition of a friction related to the motion of the interface with respect to container walls shifts the point of instability from the well known classical threshold [1] to another value. This value does not depend on the strength of dissipation and can be reproduced in the fram ework of the therm odynam ic analysis by looking for the instability of the free energy calculated in the reference fram e of norm al component, which in equilibrium is at rest with respect to container walls. The appearance of the sam e threshold in dynam ic analysis has been ascribed in Ref. [9] to the sym m etry breaking related with the violation of the G alilean invariance by the considered friction force.

In the present work we return to investigation of the corrugation instability on free surface of a supe uid liquid in presence of a counter ow [2], taking into account the viscosity of the norm alcom ponent, and show that for any nite value of viscosity the instability threshold is shifted to viscosity independent value, which is in agreem ent with the results of R ef. [9]. How ever, in our analysis this phenom enon appears in absence of the friction force violating the G alilean invariance. Therefore, the m odi – cation of the instability criterion in presence of dissipation is not a consequence of the sym m etry breaking form of the friction, but has a m ore general nature.

II.D ISPERSION RELATION

The calculation of the surface oscillations spectrum in super uid liquid in presence of a counter ow can be performed in the same way as the calculation of the spectrum of gravitational wave in norm al liquid with nite viscosity [11]. For frequencies sm all in comparison with that of the rst and the second sound, the mass and the entropy densities can be assumed to be constant. A coordingly, the conservation laws for mass and entropy are reduced to constraints:

$$div v_{s} = div v_{n} = 0; \qquad (1)$$

where v_s and v_n are super uid and normal velocities. In that limit the Navier-Stokes equation for super uid liquid can be written as [12]

¹em ail: serkor@ landau.ac.ru

$$s \frac{\partial v_s}{\partial t} + (v_s r)v_s + n \frac{\partial v_n}{\partial t} + (v_n r)v_n =$$
$$= rp \quad g + v_n; \quad (2)$$

where s and n are super uid and normal densities (= s + n being the total density), p is the pressure, g is the free fall acceleration, and is the viscosity.

Solution of Eqs. (1)-(2), satisfying the constraint $rotv_s = 0$ and corresponding to a small amplitude surface wave with frequency ! and wavevector q parallel to the surface (we assume that in the equilibrium the liquid is situated at z < 0), can be chosen in the form

$$v_{s}^{k}(\mathbf{r};\mathbf{t}) = v_{s}^{0} + i\mathbf{q} e^{qz}\mathbf{A} ; \qquad (3)$$

$$v_s^z(r;t) = q e^{qz}A; \qquad (4)$$

$$v_{n}^{k}(r;t) = v_{n}^{0} + iq (e^{qz}B + e^{kz}C);$$
 (5)

$$v_n^z(r;t) = [qe^{qz}B + (q^2=k)e^{kz}C];$$
 (6)

$$p(\mathbf{r};\mathbf{t}) = g\mathbf{z} + (7)$$

+i
$$e^{qz}$$
[_s(! $v_s^0 q)A$ + _n(! $v_n^0 q)B$];

where superscript k refers to the component of a vector parallel to the surface, $= \exp i(qr !t)$,

$$k = q_1^2 \quad i - \frac{n}{(! \quad v_n^0 q)}; \quad Rek > 0;$$
 (8)

A, B and C are (yet arbitrary) constants, and the possibility of an equilibrium counter ow (characterized by $v_s^0 \in v_n^0$) is taken into account.

Substitution of Eqs. (3)-(7) into the boundary conditions describing the conservation of m ass and entropy

$$v_{s}^{z}$$
 $(v_{s}^{k}r^{k}) = v_{n}^{z}$ $(v_{n}^{k}r^{k}) = \frac{\theta}{\theta t}$; (9)

and mechanical equilibrium

$$r^{z}v_{n}^{k} + r^{k}v_{n}^{z} = 0$$
; (10)

$$p+2 r^{z}v_{n}^{z} = (r^{k})^{2}$$
 (11)

at the surface (whose deviation from the plane z = 0 is denoted by and surface tension by) shows that they are compatible with each other when

$$s ! v_{s}^{0}q^{2} + n ! v_{n}^{0}q + i\frac{2 q^{2}}{n}^{2} + \frac{4 ^{2}q^{3}k}{n} = gq + q^{3}:$$
(12)

The derivation of Eq. (12) have not required to assume the viscosity sm all, so it is applicable for arbitrary value of viscosity.

III. IN STABILITY THRESHOLDS FOR ZERO AND FINITE VISCOSITY

For $_{s} = 0$ and = 0 Eq. (12) is transformed into the dispersion relation of gravitational wave on free surface

of norm al liquid [11], whereas in the lim it of = 0 it is reduced to equation

$$(! vq)^2 = gq + -q^3 - \frac{n s}{2} (wq)^2$$
 (13)

describing the spectrum of surface wave in super ind with a counter ow [2] derived in the framework of the non-dissipative two- uid description. Here $v = ({}_{s}v_{s}^{0} + {}_{n}v_{n}^{0})=$ is the mass velocity and $w = v_{n}^{0} + v_{s}^{0}$ the relative velocity in the super uid. The form of Eq. (13) shows that the roots with positive and negative in aginary parts (the form er corresponds to growing corrugation) exist only when the right-hand side can be negative, that is when the absolute value of w exceeds w_{c0} de ned by

$$w_{c0}^{2} = \frac{2({}^{3}g)^{1=2}}{{}_{n} {}^{s}};$$
 (14)

the instability taking place at q = $(w = w)q_c$, where $q_c^2 = q = .$

On the other hand, for any nite > 0 one of the roots of Eq. (12) crosses the real axis already when

S (q)
$$gq + -q^3 - (wq)^2$$
 (15)

touches zero, that is at

$$j_{w} j = w_{c}$$
 $\frac{2(g)^{1=2}}{s} = \frac{n}{2} w_{c0};$ (16)

the instability appearing at the lower value of relative velocity than in absence of dissipation, although at the same value of q. Note that in the lim it of zero tem perature (when $_{\rm s}$!) the criterion (16) coinsides with the Landau criterion for creation of ripplons in the reference frame of container walls.

For S (q) su ciently close to zero the value of the root which crosses real axis is given by

$$! (q) v_n^0 q = \frac{1}{2} \frac{S(q)}{_{sw} q + i q^2} :$$
 (17)

This shows that for small viscosity and w just above w $_{\rm c}$ the rate of the developm ent of the instability decreases with decreasing contrary to what is natural to expect.

By looking when the free energy of a corrugation, calculated in the reference fram e of the norm al component, is no longer positively de ned (such approach can be considered as a macroscopic generalization of the Landau criterion), the threshold for the instability of the interface between two di erent super uids has been found [9] to be given by

$$v_{s1}^{0} (v_{s1}^{0} - v_{n}^{0})^{2} + v_{s2}^{0} (v_{s2}^{0} - v_{n}^{0})^{2} = 2 (F)^{1=2}$$
; (18)

where F is a generalized restoring force, the role of which in the case of free surface is played by g. In the limit when the density of one of the liquids goes to zero, Eq. (18) is reduced to our criterion (16) obtained for a free surface of a single super uid liquid.

IV.CONCLUSION

In the present work we have investigated the dynam ic instability of a super uid liquid free surface caused by the relative motion of super uid and norm alcom ponents along the surface. The value of the instability threshold for nite viscosity, given by Eq. (16), has been found to be independent of viscosity, but lower than in absence of dissipation. The same criterion can be obtained by looking for the therm odynam ic instability in the reference frame of the normal component (which in equilibrium is xed by container walls).

Analogousmodi cation of the instability threshold has been found [9] to take place on the interface between two super uids in presence of a friction with respect to the reference fram e related with container walls [13], which leads to violation of the G alilean invariance. Note that in our problem the same phenom enon appears in situation when the form of dissipation does not imply the explicit selection of the particular reference fram e. Nonetheless, the presence of dissipation (a nite value of viscosity) turnes out to be su cient to produce the sam e criterion for surface instability as in the case when the form of the friction leads to the direct violation of the Galilean invariance.

The rst experimental observation of the Kelvin-Helm holtz instability on the interface between ³He-A and ³He-B by B laauw geers et al. [10] have unam bigously dem onstrated that it indeed takes place not at the classical, but at the modi ed value of the threshold. A coording to our results, the same can be expected from the instability on free surface of super uid⁴He.

The author is grateful to G E. Volovik for useful discussion. This work has been supported by the Program "Quantum Macrophysics" of the Russian Academy of Sciences, by the Program "Scienti c Schools of the Rus-

sian Federation" (grant No. 00-15-96747), by the Swiss National Science Foundation and by the Netherlands Organization for Scienti c Research (NWO) in the fram ework of Russian-Dutch Cooperation Program .

- [1] L.D. Landau and E.M. Lifshitz, Gidrodinamika, Nauka, Moscow (1986) Fluid Mechanics, Pergamon Press (1989)], Sec. 62, problem 3.
- [2] S.E.Korshunov, Europhys. Lett. 16, 673 (1991).
- [3] P.W. Egolf, J.L. Olsen, B. Roericht and D.A. Weiss, Physica B 169, 217 (1991).
- [4] M.Yu.Kagan, Zh.Eksp.Teor.Fiz. 90, 498 (1986) [Sov. Phys.-JETP 63,288 (1986)].
- [5] A.F. Andreev and A.Ya. Parshin, Zh. Eksp. Teor. Fiz. 75,1511 (1978) [Sov.Phys.-JETP 48,763 (1978)].
- [6] K.O. Keshishev, A.Ya. Parshin and A.V. Babkin, Pis'ma ZhETF 30, 63 (1979) [JETP Lett. 30, 56 (1979)]; Zh. Eksp. Teor. Fiz. 80, 716 (1981) [Sov. Phys. - JETP 53, 362 (1981)].
- [7] K.W. Madison, F. Chevy, V. Bretin and J. Dalibard, Phys.Rev.Lett.86,4443 (2001).
- [8] S. Sinha and Y. Castin, Phys. Rev. Lett. 87, 190402 (2001).
- [9] G E. Volovik, cond-m at/0202445 (2002); P is'm a ZhETF 75,491 (2002).
- [10] R. Blaauw geers, V. B. Eltsov, G. Eska, A. P. Finne, R. P. Haley, M. Krusius, J.J. Ruohio, L. Skrbek, and G.E. Volovik, cond-m at/0111343 (2001).
- [11] Ref. 1, Sec. 25, problem 1.
- [12] Ref. 1, Sec. 140, problem 1.
- [13] The same type of dissipation has been taken into account by K agan [4] when studying the instability of the quantum interface between super ind and solid ⁴He.,