Number Fluctuation in an interacting trapped gas in one and two dim ensions

R.K.Bhaduri¹, M.V.N.Murthy and MuoiN.Tran¹ The Institute of Mathematical Sciences Chennai 600 113, India 1. Permanent Address: Department of Physics and Astronomy, M dMaster University, Hamilton, Ontario, Canada L8S 4M 1

Abstract

It is well-known that the number uctuation in the grand canonical ensemble, which is directly proportional to the compressibility, diverges for an idealbose gas as T ! 0. We show that this divergence is removed when the atom s interact in one dimension through an inverse square two-body interaction. In two dimensions, similar results are obtained using a self-consistent Thom as-Ferm i (TF) model for a repulsive zero-range interaction. Both models may be mapped on to a system of non-interacting particles obeying the Haldane-W u exclusion statistics. We also calculate the number uctuation from the ground state of the gas in these interacting models, and compare the grand canonical results with those obtained from the canonical ensemble.

Typeset using REVT_EX

I. IN TRODUCTION

Consider an idealbose gas at low tem peratures. In the grand canonical ensemble (GCE), the compressibility and the number uctuation of this system diverge at low tem peratures [1]. To quote Landau and Lifshitz [1]:

...in a bose gas at tem perature $T < T_c$, the pressure is independent of the volume, i.e. the compressibility becomes in nite. A coordingly...this would imply that the uctuations of the number of particles also become in nite. This means that, in calculating uctuations in a gas obeying bose statistics, the interactions between the particles cannot be neglected at low tem peratures, how ever weak this interaction m ay be. W hen the interactions, which m ust exist in any actual gas, is taken into account, the resulting uctuations are nite.

One purpose of this paper is to demonstrate the validity of this statement in one and two dimensional traps for a special class of interactions.

In one dimension, the quantum many-body problem of particles in a harmonic oscillator interacting with an inverse square two-body potential is exactly solvable [2,3]. Moreover, it is known that the global properties of these interacting bosons are the same as those of non-interacting particles obeying the Haldane-W u generalized exclusion statistics (ideal haldons) [4,5]. U sing this mapping, we show that the number uctuation is nite as $T \, ! \, 0$, no matter how weak the interaction strength is. In this model, since the exact correlation function is known for some speci c strength parameters of the interaction [3], it is also possible to verify explicitly the well-known relation between its integral and the number uctuation.

In two dimensions, there is no suitable exactly solvable model for our purpose. We therefore consider the mean-eld model of bosons trapped in a harmonic oscillator, and interacting pair-wise with a zero-range repulsive pseudo-potential [6]. In the absence of this interaction, the number uctuation diverges at $T = T_c$. When the interaction is present, however weak, we show that the compressibility and hence the number uctuation of the system is nite right down to T = 0. Moreover, they are shown to be identical to those of a collection of non-interacting haldons.

A nother quantity of som e interest is the number uctuation of particles from the ground state of the system, which is present even when the total number of particles in the trap is xed. Consider a dilute gas of bosons in a trap at T = 0. The system is in its ground state. When a certain amount of excitation energy is given to the system, it may be absorbed in many possible ways, so that the number of bosons remaining in the ground state is not xed. This number uctuation for non-interacting bosons in a harmonic trap has been calculated by a number of authors [9] as a function of the excitation energy or temperature. Therm odynam ic identities and particle number uctuations in weakly interacting BEC have also been analysed when the particle number is xed [10]. For ideal bosons, the number uctuation from the ground state diverges at low temperatures in GCE, but this can be avoided using more careful canonical, orm icrocanonical treatment. When the inverse square pairwise interacting model, we can also perform the canonical ensemble (CE) calculations, and com pare with the GCE results. We not that even though the ground state number

uctuation goes to zero as T ! 0 in both GCE and CE, the very low tem perature behaviors are rather di erent. At higher tem peratures the results from CE and GCE tend to coincide. Sim ilar studies can be made for ferm ionic system sat low tem peratures. Since the interactions that we have used are of statistical character [5,11], our results interpolate from bosonic to ferm ionic behavior with the variation in the strength of the interaction.

The outline of the paper is as follows: Sect. II deals with the total number uctuation in GCE for interacting models. Sect. II (a) is devoted to a discussion of the uctuations in one dimension, and II (b) to uctuation in two dimensions. In Sect. III, the ground state number uctuations for the one-dimensional interacting model in GCE and CE are calculated and compared. Particular attention is paid to the low temperature behavior (see also Appendix). We conclude the paper with a brief discussion of the results.

II.FLUCTUATIONS IN GCE

The number uctuation in a gas in the GCE formalism is defined by $(N)^2 = (< N^2 > < N >^2)$, where the angular brackets denote ensemble averaging. For an ideal bose gas,

$$(N)^{2} = \int_{k=0}^{k} hn_{k}i(hn_{k}i+1);$$
 (1)

where the single particle occupancy $hn_k\, i$ at a given tem perature $T\,$ and energy $_k\,$ are given by the bose distribution function

$$hn_{k}i = \frac{1}{\exp[(k_{k})] - 1};$$
(2)

Here = 1=T with the Boltzm ann constant $k_B = 1$, and denotes the chemical potential. The isotherm alcompressibility _ of a gas of density _ 0 is related to the number uctuation of the system in GCE :

$$\frac{(N)^2}{N} = T_{0} T$$
(3)

The problem with the number uctuation in a bose gas in GCE is obvious from Eq. (1). If there is BEC, then a macroscopic fraction of the particles occupy the ground state for $T < T_c$, so that $(N_0)^2 ' N^2$. Even if there is no BEC, the same is the case as T ! 0. In the therm odynam ic lim it, therefore, the uctuation diverges below the critical tem perature, or in any case at T = 0. This is manifestly incorrect, since at T = 0, all the bosons are in the ground state, and the number uctuation should vanish. In the presence of interactions, how ever, this number uctuation is expected to be nite. We demonstrate this by analyzing two models in one and two dimensions.

A.Fluctuations in a one dim ensionalm odel

We rst consider the exactly solvable one dimensional Calogero-Sutherland M odel (CSM) of a system of interacting particles with the H am iltonian [2,3]

$$H = \frac{X^{N}}{\sum_{i=1}^{m}} \frac{h^{2}}{2m} \frac{\theta^{2}}{\theta x_{i}^{2}} + \frac{1}{2}m !^{2} x_{i}^{2} + \frac{h^{2}}{m} \frac{X^{N}}{\sum_{i< j=1}^{m}} \frac{g(g \ 1)}{(x_{i} \ x_{j})^{2}};$$
(4)

with the dimensionless coupling parameter g 0. The particles are conned in a harmonic well and the thermodynamic limit is obtained by taking !! 0 as N ! 1, with !N = constant. In the thermodynamic limit, the properties of the system are translationally invariant, and would be the same if the particles were on a line, or a circle, instead of a harmonic connement. To make the problem well de ned quantum mechanically, we have to dem and that the wave functions go to zero as $jx_i = x_j f$ whenever two particles i and j approach each other. Since the particles cannot cross each other, we may choose the wave function to be either symmetric (bosonic) or antisymmetric (fermionic). For g = 0 and 1, the model describes free bosons and free fermions respectively.

Using the relation between the integral of the correlation function and the num ber uctuation, we now show that the number uctuation vanishes at zero temperature in the above interacting model unlike the ideal boson result in GCE. If (r) denotes the two-particle ground state density-density correlation function in the ground state, with $r = jx_1 - x_2 j$, then the number uctuation is related to the correlation function as [1]

$$\frac{(N)^{2}}{N} = \frac{1}{1} = \frac{2}{1} (x) dx:$$
(5)

Note that the ground state correlation function (r) is defined only for r 0. However, in computing the above integral it is necessary to assume (r) to be even function, and extend the domain of integration to negative values of r [12]. Unlike the one-particle of -diagonal density matrix, (r), by definition, is related to the diagonal element of the the two-particle density matrix, and is the same in CSM for bosons or fermions. In this section we work in the bosonic basis. The correlation functions are known exactly in the CSM for three values of g independent of whether the particles are bosons are fermions and are given (in the therm odynam ic limit) by β

g = 1 : (r) = s(r)² =
$$\frac{\sin(r)}{r}^{\#_2}$$
 (6)

$$g = 1=2$$
: $(r) = s(r)^{2} + \frac{ds}{dr} \int_{r}^{Z_{1}} dt [s(t)]$ (7)

$$g = 2$$
: $(r) = s(2r)^2 - \frac{ds(2r)}{dr} \int_{0}^{2} dt [s(t)];$ (8)

where the Fermin on entum k_F is set equal to so that the maximum central density is unity. For all three forms of (r) given above, explicit calculations show that

Ζ

$$\int_{1}^{1} (x) dx = \int_{1}^{2} [s(r)]^{2} dx = 1;$$
(9)

independent of the value of g. Substituting this result in Eq.(5) it follows that for interacting bosons in CSM the uctuation vanishes identically at T = 0.

W hile we cannot obtain the exact (r) in CSM for all g, the same m ay be calculated for all values of g in the harm onic lattice approximation. The correlation function so obtained compares very well with the exact correlation functions for g = 1=2;1;2 and is given by [13]

$$(x) = \int_{n \in 0}^{X} \frac{1}{4 F(n;0)} \exp^{h} \frac{(x_{0} n)^{2}}{4F(n;0)} \int_{0}^{1} (10)$$

where $_0 = N = L$ is the average density and

$$F(n;0) = \frac{1}{2^2} \int_{0}^{Z} dy \frac{1}{y} \frac{\cos(yn)}{y^2=2} = \frac{1}{2^2} \int_{0}^{Z} dy \frac{1}{y} \frac{\cos(yn)}{y} : \quad (11)$$

The above expression is given for completeness and its exact form is not needed for further calculations. Again integrating over the real line we get a result identical to that obtained using the exact correlation functions in CSM. Thus the uctuation vanishes identically for all g in this approximation at zero temperature. However, the result does not give any indication of the behavior of the uctuation at nite temperature. To do this we take recourse to the mapping between the CSM and the exclusion statistics rst proposed by Haldane through a generalized Pauli principle [5,17].

A crucial property of exclusion statistical interactions is that they should cause shifts in single particle energies at all scales [14] (see next section). This property is realized by a large class of one dimensional models of interacting fermions where Fermi liquid theory breaks down [15,16]. In fact it has been shown exactly that quasiparticles with nontrivial exclusion statistics exist in a class of models that are solved by the Bethe ansatz [17]. Thus the result obtained below should be, in principle, valid in a large class of models with interactions. In particular, it is well known that the interacting particle of CSM may be regarded as ideal exclusion statistics particles or sim ply haldons [5]. The therm odynam ic properties of an ideal gas of exclusion statistics particles have been investigated widely [18{21}]. The distribution function has been computed and is given by

hn()i =
$$\frac{1}{(w() + g)}$$
; (12)

where ${\tt w}$ () is the solution of the equation

$$w ()^{g} (1 + w ())^{(1 g)} = e^{()}$$
(13)

At zero tem perature we have,

$$\operatorname{hni} = \frac{1}{g}; \quad \text{for} \quad k \tag{14}$$

and zero otherwise.

Note that the distribution function reduces to the usual Ferm i and Bose distribution functions for g = 1 and g = 0 respectively and in general g is regarded as the exclusion statistics parameter. Indeed, the statistical parameter g is precisely the interaction strength as given in Eq. (4) in one dimension. However, the distribution function as given above is valid in general and not necessarily restricted to one dimensionalm odels. The following discussion is therefore used for illustration in the case of one-dimensionalm odel but not restricted to this case alone.

The number uctuation at a given energy $_k$ is given by,

$$(n_k)^2 = T \frac{\theta n_k i}{\theta}$$
: (15)

Substituting for hn_k i from Eq. (12), we have for total number uctuation [18],

$$(N)^{2} = \int_{k=0}^{k} w_{k} (1 + w_{k}) hn_{k} i^{3}$$

$$= \int_{k=0}^{k} hn_{k} i (1 - ghn_{k} i) (1 + (1 - g)hn_{k} i):$$
(16)

The number uctuation vanishes at $T ! 0 \operatorname{since} n_k ! 1=g$ below the Ferm ienergy and zero otherwise. This result holds no matter how weak the interaction strength is. However, at g = 0, the bosonic limit, the number uctuation diverges as noted earlier. In this exactly solvable model, we have thus shown that interactions do remove the uctuation catastrophe encountered in the ideal B ose gas.

W hile these results have been derived in one dimension, extension to higher dimension is non-trivial since there is no suitable exactly solvable many body model. However, it has been shown that models with short range interactions in two dimension may be regarded as obeying exclusion statistics in the mean-eld picture [22]. We discuss the uctuation in these models in the next section.

B.Fluctuations in a two dim ensionalm odel

We consider a two-dimensional system of bosons interacting via a zero-range repulsive pseudo-potential. The quantum dynamics is then approximated by the following Ham iltonian

$$H = \sum_{i=1}^{N} \frac{p_i^2}{2m} + \frac{1}{2}m ! r_i^2 + \frac{2}{m} g_{i < j}^N (r_i - r_j); \qquad (17)$$

where the momenta and coordinates are planar vectors. The one-body potential generated by the above zero-range interaction (including exchange) is

$$U(n(r)) = \frac{2 h^2}{m} g n(r); \quad g = 2g; \qquad (18)$$

where n (r) is the local number density of the system. In two-dimensions, g 0 plays the role of the statistical parameter, with g = 0 for non-interacting bosons. At nite temperature, for $T > T_c$, the Thom as Fermi approximation yields [6,7]

$$n(\mathbf{r}) = \int_{-\infty}^{Z} \frac{d^2 \mathbf{p} = (2 \ h)^2}{\exp\left[\left(\frac{\mathbf{p}^2}{2m} + V(\mathbf{r})\right)\right] - 1} ;$$
(19)

where the Thom as Ferm im ean potential V (r) is given by

$$V(\mathbf{r}) = V_0(\mathbf{r}) + \frac{2 h^2}{m} g n(\mathbf{r})$$
 : (20)

Here V_0 (r) is the one-body harm onic trap. Note that Eq.(19) is valid only in the absence of a condensate. It has been shown [6,7]. however, that for a nonzero positive g, a self-consistent solution of this equation satisfying n (r)d²r = N m ay be obtained right down to T = 0. This solution has a lower free energy than the one with a condensate [8], so we may take $T_c = 0$ for g > 0.

The momentum integration may be done analytically:

$$n(r) = \frac{m}{2 h^2} \ln [l \exp [(V(r))]]$$
 (21)

The local number uctuation (between r and r + dr) in GCE is given by

$$(N)^{2} = T \frac{(0n)(r)}{(0)};$$
 (22)

where the coe cient of the tem perature T on the rhs is related to the com pressibility. Taking the derivative of the local density with respect to the chem ical potential we have

$$(N)^{2} = T \frac{m}{2 h^{2}} \frac{1}{\exp[(V(r))] + 1 + g};$$
 (23)

Note that is a function of temperature, and is determined by the condition that R n(r)d²r = N, and in the therm odynamic limit it approaches the lowest energy eigenstate as the temperature goes to zero. A few remarks on the therm odynamic limit are in order: The therm odynamic limit is reached when N ! 1 and ! ! 0. However, in the limit of no con nement the density of states becomes a constant and there is no critical temperature below which condensation takes place. Preserving the density of states as in a two dimensional oscillator, the condensation temperature of an ideal bose gas is given by

$$T_{c}^{(0)} = (6 = {}^{2})^{1=2} N^{1=2}!$$

Thus the limit N ! 1 and ! ! 0 is obtained keeping $T_c^{(0)}$ constant. In an ideal bose gas (g = 0), no self-consistent solution of Eq. (21) can be found for a xed N below this temperature $T_c^{(0)}$. However, when g > 0 no matter how small, the self-consistent solution of Eq. (21) may then be found for all T > 0.

In this lim it the uctuation is given by,

$$(N)^{2} = T \frac{m}{2 h^{2}} \frac{1}{\exp\left[\left(\frac{2 h^{2}}{m} gn\right)\right] + g};$$
 (24)

where n is the constant density given in the therm odynam ic lim it. (For the one-dimensional case, this was denoted by $_0$ earlier.)

In the absence of interaction, g = 0, the chemical potential goes to zero at $T = T_c$ and the above expression diverges as expected. However, as in the one dimensional case, the uctuation remains nite and approaches zero as T ! 0 when g is nite, however weak the interaction m ay be.

The same result m ay be obtained in the non-interacting exclusion statistics description in two-dimensions. The local density as a function of the radial coordinate is given by [6],

n (r) =
$$\frac{d^2 p = (2 h)^2}{[w + g]}$$
; (25)

where the local variable w (p;r) is de ned through W u's equation within TF approximation

$$w^{g}(1 + w)^{1 g} = \exp \left[\left(\frac{p^{2}}{2m} + V_{0}(r) \right) \right];$$
 (26)

and g is the exclusion statistics parameter which we will identify with the interaction strength in them ean-eld picture. We have retained the trap potential V_0 as in the interacting picture. O noe again the momentum integration may be done easily and we obtain,

$$n(r) = \frac{m}{2 h^2} \ln \frac{1 + w_0}{w_0} ; \qquad (27)$$

where the local variable w_0 (r) is determined through

$$w_0^g (1 + w_0)^{1 g} = \exp[(V_0(r))]$$
: (28)

The uctuation is then found by using Eq. (22) and we have

$$N^{2} = T \frac{(en (r))}{(e)};$$
 (29)

$$= T \frac{m}{2 h^2} \frac{1}{w_0 + g} :$$
 (30)

In the bosonic lim it taking g = 0 in Eq. (28), we have $w_0 = \exp[(V_0)]$ 1. Substituting this in the above expression for the uctuation in the therm odynam ic lim it it is easy to see that at T_c there is divergence. However, for positive de nite g there is no divergence. Furtherm one the equivalence between the non-interacting exclusion statistics picture and the mean-eld description is established using the following relationship

$$w_0(\mathbf{r}) = \exp[((\mathbf{V}(\mathbf{r})))] \quad 1;$$
 (31)

where V (r) is the self consistent mean-eld potential. Substituting this in Eq.(30), and taking the therm odynam ic lim it yields Eq.(24). Note that this equivalence allows one to calculate uctuations in either the mean eld picture or in the non-interacting exclusion statistics picture. Indeed this holds for the computation of other global therm odynam ic quantities as well.

III. GROUND STATE FLUCTUATION IN THE CANON ICAL ENSEMBLE

The total number uctuation de ned in the previous section in the GCE is obtained by summing over all the single particle states. In the canonical ensemble how ever, the number of particles is xed and therefore the uctuation in particle number has to be de ned with respect to a reference state. One way de ning the uctuation is to bok at the ground state occupancy as a function of temperature, which is present even when the total number of particles in the trap is xed. At T = 0 all the particles are in the ground state. At a nonzero tem perature (or excitation energy), there are m any ways of exciting the particles from the ground state, leading to a uctuation in the ground state population. This number uctuation for non-interacting bosons in a harm onic trap has been calculated by a number of authors [9] as a function of the excitation energy or tem perature. This has also been calculated for ferm ions in the CE by us in a previous publication [23]. Unlike the case of GCE for bosons, the ground state uctuation in the CE is nite at all tem peratures.

W e may extend the analysis of uctuations in CE to particles interacting via the inverse square pair-wise interaction in one dimension. It is more convenient to perform the calculation in the ferm ionic basis, although the form ulae given here are applicable for both interacting ferm ions, or interacting bosons. In this section, quantities like energy and ground state number uctuation of the interacting system with interaction strength g will be denoted by a bracketed superscript, e.g. E^(g), ((N_0)²)^(g). W e recall that the spectrum of the CSM H am iltonian given in Eq. (4) is exactly known. The states may be labeled by a set of ferm ionic occupation numbers fn_kg; k = 1; ...;1; n_k = 0;1. The energy E^(g) of the system in the ferm ionic basis is then given by,

$$E^{(g)}fn_{k}g = \sum_{k=1}^{N} n_{k} ! (1 g) \frac{N(N 1)}{2};$$
(32)

where $_{k} = (k \quad \frac{1}{2}h!)$ denotes the harm onic oscillator energy levels and $N = {}^{P} \frac{1}{k-1}n_{k}$. As can be seen from Eq. (32), the e ect of the interaction is that each particle shifts the energy of every other particle by a constant h! (g 1). The energy functional can also be written as

$$E^{(g)} fn_k g = \sum_{k=1}^{k} n_k + (1 - g) \sum_{k_1 < k_2 = 1}^{k} n_{k_1} n_{k_2} :$$
(33)

The exact spectrum of the model is thus reproduced by an elective H am iltonian of quasiparticles with constant density of states and constant Landau parameters. As mentioned before, this scale invariant energy shift is the basic reason for the occurrence of nontrivial exclusion statistics where g plays the role of exclusion statistics parameter with g = 0;1 for bosons and ferm ions.

The general canonical partition function in any basis is written in the occupation number representation as [5]

$$Z_{N}^{(g)} = \sum_{fn_{k}g}^{X} e^{-E^{(g)}fn_{k}g}$$
: (34)

U sing the energy spectrum in CSM given in Eq. (32), the N-particle partition function in this one dimensionalm odel is given by

$$Z_{N}^{(g)} = e^{\sim (1 - g)\frac{N - (N - 1)}{2}} Z_{N}^{F};$$
(35)

where $\sim = h!$ and Z_N^F is the N particle ferm ion partition function. Setting g = 0, the bosonic partition function is obtained,

$$Z_{N}^{B} = e^{-\frac{N(N-1)}{2}} Z_{N}^{F} :$$
 (36)

Combining Eqs. (35) and (36) we may write the partition function for CSM as,

$$Z_{N}^{(g)} = (Z_{N}^{F})^{g} (Z_{N}^{B})^{1 g};$$
(37)

The canonical partition function given above is exact in CE and may be used for calculating the therm odynam ic properties of the system in CSM within the canonical ensemble form alism. The moments of the occupation number are related to the partition function by

$$hn_{k}i^{(g)} = \frac{1}{Z_{N}^{(g)}} y_{k} \frac{\partial Z_{N}^{(g)}}{\partial y_{k}};$$
(38)

$$hn_{k}^{2}\dot{i}^{(g)} = \frac{1}{Z_{N}^{(g)}} Y_{k} \frac{\theta}{\theta Y_{k}} \overset{\circ}{\theta} Y_{k} \frac{\theta Z_{N}^{(g)}}{\theta Y_{k}} \overset{\circ}{A} ; \qquad (39)$$

where $y_k = \exp(k_k)$. Therefore, it follows that

h(
$$n_k$$
)² $i^{(g)} = y_k \frac{\theta n_k i^{(g)}}{\theta y_k}$: (40)

U sing Eqs. (37)-(39), $hn_k i^{(g)}$ can be expressed in terms of those of ferm ions and bosons via:

$$hn_k i^{(g)} = ghn_k i^F + (1 \quad g)hn_k i^B$$
(41)

Unlike $hn_k i^{(g)}$, $hn_k^2 i^{(g)}$ does not have a simple form as Eq. (41). However, the expression for the uctuation in the occupation number does:

$$((n_k)^2)^{(g)} = g((n_k)^2)^F + (1 g)((n_k)^2)^B$$
: (42)

Eq. (42) gives only the uctuation in the occupation of a given level k, while the quantity we are seeking is the ground state number uctuation. The latter is form ally de ned in any ensemble as:

$$(N_0)^2 = \sum_{k}^{X} (n_k)^2 = \sum_{k}^{X} (m_k^2 i m_k i^2)$$
 (43)

where the sum k runs over only the levels which are completely occupied at zero temperature. Thus, in an ab initio calculation, one would form ally sum over the quasiparticle levels which are occupied at T = 0 to get $((N_0)^2)^{(g)}$. Fig. 1 shows the level ow in CSM as a function of g obtained from Eq. (33) at T = 0. It can be seen that as g changes from the ferm ionic to the bosonic end, the number of levels contributing to the ground state remains constant, while the Ferm i energy decreases accordingly. This means that one may obtain $((N_0)^2)^{(g)}$ by simply substituting the ground state uctuations for ferm ions and bosons. ie:

$$((N_0)^2)^{(g)} = g((N_0)^2)^F + (1 g)((N_0)^2)^B$$
(44)

In the previous section we have discussed the ground state number uctuation in GCE and CE in an interacting system. We now compare the results obtained in these two ensembles. Due to numerical di culty at low temperatures, we are able to do the comparison only for N = 10 particles. Despite this limitation, some interesting points can be made. In Fig. 2 (a), we show the behavior of the relative ground state uctuation against temperature of interacting bosons for both ensembles for interacting strengths of g = 0; 1=2; 1. The g = 0 case corresponds to free bosons. Fig. 2 (b) shows the low temperature region of the curves in Fig. 2 (a). Note that as T ! 0 the GCE uctuation for free bosons diverges as expected, whereas those for interacting bosons remain nite and approach zero. Note also that in the ideal H aldane gas picture the g = 1=2 case corresponds to sem ions is found using Eqs. (12), (13) and (17), where is determined by the constraint that the average total number of particles is N.

Unlike in GCE, the CE ground state uctuation for free bosons remain nite as T ! 0. The ground state uctuation of free bosons, however, approaches zero exponentially (see Fig. 2(b)), contrary to previous results that nd a linear dependence with T all the way to T = 0 [9,25]. In Appendix A we give the low temperature expansions of the uctuation squared (N_0)² in powers of x, where x = e $\tilde{}$. Clearly, at very low temperatures (N_0)² is independent of the number of particles N (in both GCE and CE). Therefore, the exponential behaviour of the uctuation of free bosons at low temperatures should remain valid even in the large N limit. In the case of fermions the CE and GCE uctuations are similar except at very low temperatures. Clearly, the CE and GCE curves of fermions approach zero di erently as T ! 0. For high temperatures, on the other hand, GCE and CE give identical results, as expected.

It is to be noted from Fig. 2 (a) that the number uctuation for free bosons and free ferm ions cross at a certain temperature, with the ferm ion one getting bigger at higher temperatures. This is to be expected, since the number of possibilities of creating holes within the ferm isea, and distributing particles above, increases more rapidly than for bosons, whose ground state has only one level. Sem ions (g = 1=2), which may be regarded either as interacting bosons, or as interacting ferm ions, have uctuations interm ediate between those of bosons (g = 0) and ferm ions (g = 1).

In one dimension, we have calculated the number uctuation of bosons interacting via CSM using the ideal H aldane picture for both GCE and CE. The next challenge is to not the exact uctuation using combinatorics. This has been done for free ferm ions which correspond to interacting bosons with interacting strength of g = 1 [23]. W ork is in progress to calculate this exact uctuation for a general g.

ACKNOW LEDGM ENTS

Two of us (RKB and MNT) thank the Institute of M athem atical Sciences, where part of this work was done, for hospitality. This work was supported by Natural Sciences and Engineering Research Council (NSERC) of Canada.

APPENDIX:LOW TEMPERATURE EXPANSIONS IN GCE AND CE

The low temperature behaviour of them odynam ic quantities in GCE are well known for bosonic systems. However, a comparison between GCE and CE calculations at low temperatures is not usually discussed for either Bose or Ferm i systems. Further, making use of some asymptotic expansions, the CE uctuation for bosons was earlier found to be linear right down to T = 0. However, we give here the expansion of the uctuation squared at low temperature in power of x, where $x = e^{-}$, and show that the CE uctuation of bosons is in fact exponential at very low T. In GCE only expansion for ferm ions is possible, since the uctuation tends to in nity at low temperature for bosons. Both expansions are possible in CE.

a. Grand Canonical Ensemble: In GCE the (ferm ionic) occupation number is:

$$hn_{k}i_{GCE} = \frac{1}{x^{(-k)} + 1} = \frac{1}{x^{(-k-1-2)} + 1}$$
(45)

for a one dimensional system. The ground state number uctuation squared is given by:

$$h(N_0)^2 i_{GCE} = \frac{k_F}{k=0} \frac{x^{(k-1-2)}}{[x^{(mu-k-1-2)} + 1]^2}$$
(46)

where k_F is the Ferm i level. At low tem peratures, for the one-dimensional harm onic oscillator, $_0 = N$. Therefore,

h(N₀)²
$$i_{GCE} = \frac{p}{x} 2x + 4x^{3=2} 4x^2 + 6x^{5=2} 8x^3 + 8x^{7=2} 8x^4 \dots$$
; N 4: (47)

Note that the rst few terms are independent of N.

b. CanonicalEnsemble: In CE the rst and second moments of the occupation number are known [24]:

$$hn_{k}i = \frac{1}{Z_{N}} \int_{j}^{k} ()^{j+1} x^{j} Z_{N} j; \qquad (48)$$

$$hn_{k}^{2}i = \frac{1}{Z_{N}} \sum_{j}^{N} (j)^{j+1} [j (j 1)] x^{j} Z_{N} j;$$
(49)

where the upper and lower signs refer to bosons and ferm ions respectively. Sum m ing over the ground states up the Ferm i level gives the ferm ionic ground state num ber:

$$hN_{0}i_{CE} = \frac{1}{Z_{N}} \frac{x^{N}}{y} (1)^{j+1}x^{j=2} \frac{1}{1} \frac{x^{jN}}{x^{j}} Z_{N-j};$$
(50)

where we let $_{k} = k$ 1=2, with $k = 1;2; \dots$ Therefore,

$$h(N_0)^2 i_{CE} = \frac{x^N}{m_k i} \frac{x^N}{m_k i^2} m_k i^2$$

= x + 2x⁴ + ::: ; N 4; (51)

where we have used $hn_k i$ = $hn_k^2 i$ and again the $\mbox{ rst few term s are independent of the system size N .$

For bosons the ground state consists of one single lowest level, the low temperature expansion of the number uctuation is given by

Again as in the ferm ionic case the rst few terms in the low temperature expansions are independent of the system size. Indeed it is interesting to note that in CE, the uctuations in both the systems approach zero as $T \, ! \, 0$ in exactly identical fashion.

REFERENCES

- [1] Landau L D and Lifshitz E M 1958 Statistical Physics (Addison-Wesley, Reading) p. 354
- [2] Calogero F 1969 J. M ath. Phys. 10 2191 2197
- [3] Sutherland B 1971 J. M ath. Phys. 12 246; 1992 Phys. Rev. B 45 907
- [4] Haldane F D M 1991 Phys. Rev. Lett. 67 937
 W u Yong Shi1994 Phys. Rev. Lett. 73 922
- [5] M urthy M V N and Shankar R 1994 Phys. Rev, Lett. 73 3331; 1995 Phys. Rev, Lett. 75 353
 - HaZNC 1994 Phys. Rev, Lett. 73 1574
- [6] Bhaduri R K, Reim ann S M, Viefers S, Ghose Choudhury A, and Srivastava M K 2000 J. Phys. B: At. M ol. Opt. Phys. 33 3895
- [7] Van ZylB P, Bhaduri R K, and Sigetich J 2002 J. Phys. B: At. Mol. Opt. Phys. 35 1251
- [8] Fernandez J P and M ullin W J cond-m at/0203175
- [9] G rossm an S and Holthaus M 1996 Phys. Rev. E 54 3495; 1997 Phys. Rev. Lett. 79 3557
 - Politzer H D 1996 Phys. Rev. A 54 5048
 - Gajda M and Rzazewski K 1997 Phys. Rev. Lett. 78 2886
 - Wilkens M and Weiss C 1997 J. Mod. Opt. 44 1801
 - Navez P, Bitouk D, Gajda M, Idziaszek Z, and Rzazewski K 1997 Phys. Rev. Lett. 79 1879
- [10] Illum inatiF, Navez P and W ilkens M 1999 J. Phys. B 32 L461
- [11] Viefers S, Hansson T H, and Leinaas JM 2001 J. Phys. B: At. Mol. Opt. Phys. 34 4609
- [12] Mehta M L 1967 Random matrices and statistical theory of energy levels (A cademic Press, New York) p.199
- [13] Sen D and Bhaduri R K 1999 Can. J. Phys. 77 327
- [14] M urthy M V N and Shankar R 1994 Phys. Rev, Lett. 72 3629
- [15] Anderson P W 1992 Prog. Theor. Phys. Suppl. 107 45
- [16] M urthy M V N and Shankar R 1995 Physica B212 315
- [17] Bernard D and W u Yong Shicond-m at/9404025 (unpublished)
- [18] C.Nayak and F.W ilczek, Phys. Rev. Lett. 73, 2740 (1994).
- [19] Isakov S B, Arovas D P, M yrheim J, and Polychronakos A P 1996 Phys. Lett. A 212 299
- [20] Yang H S, Lee B, and Park C 1997 J. Korean Phys. Soc. 30 14
- [21] A oyam a T cond-m at/0005336 (unpublished)
- [22] Bhaduri R K, Murthy M V N, and Srivastava M K 1996 Phys. Rev. Lett. 76 165
- [23] Tran M N, Murthy M V N, and Bhaduri R K 2001 Phys. Rev. E 63 031105
- [24] Parvan A S, Toneev V D, and Ploszajczak M, 2000 Nucl. Phys. A 676 409
- [25] Tran M N 2000 G round State Number Fluctuations In Finite Bose And Ferm i Systems M Sc. Thesis
- [26] M urthy M V N and Shankar R 1999 Phys. Rev. B 60 6517

FIGURES

FIG.1. The level ow of quasi-particle energy levels in CSM as a function of g.

FIG.2. (a) The ground state uctuation for the one-dimensional CSM system in GCE (Eq: (17) and CE (Eq: (44) as a function of temperature for N = 10. We show the results for fermions, bosons and also semions (g = 1=2).

FIG.3. (b) Same as in (a) but using the low temperature expansions given in the Appendix

•





