

PACS numbers: 36.40.Qv, 71.15.Nc, 71.15.Pd.

Metastable quasi-one-dimensional ensembles of nitrogen clusters N_8

N. N. Degtyarenko, V. F. Elesin, L. A. Openov, A. I. Podlivaev

Moscow Engineering Physics Institute (State University),

Kashirskoe sh. 31, Moscow 115409, Russia

By means of *ab initio* and tight-binding calculations it is shown that metastable nitrogen clusters N_8 (boats) can form quasi-one-dimensional ensembles in which the nearest clusters N_8 are bound to each other by covalent bonds. Those ensembles are characterized by rather high energy barriers (~ 0.3 eV) that prevent the fission of the ensembles into isolated N_8 clusters and/or N_2 molecules.

A nitrogen molecule N_2 is known to be the most stable form of nitrogen at standard conditions. Contrary to the overwhelming majority of other chemical elements, the cohesive energy per atom, $E_{\text{coh}}(n) = E_{\text{tot}}(n) - E_1$ (where $E_{\text{tot}}(n)$ is the total energy of an n -atom cluster N_n , E_1 is the energy of an isolated nitrogen atom) decreases with n , so it is energetically favorable for clusters N_n with $n > 2$ to decompose into separated N_2 molecules and thus to lower their energy. However, recently it was shown theoretically [1, 2, 3] that several nitrogen clusters, e.g., N_4 and N_8 , can exist in metastable states which correspond to the local minima of the potential energy as a function of atomic coordinates. Although metastable clusters N_n with $n \geq 4$ have not been synthesized yet, they attract much theoretical attention as potential high energy density materials (HEDM). The energy $E_{\text{acc}}(n) = E_{\text{coh}}(2) - E_{\text{coh}}(n)$ accumulated in the N_n ($n \geq 4$) systems approaches the value of $E_{\text{acc}} \sim 3$ eV/atom

[1, 2, 3].

In order to make use of N_n ($n \geq 4$) clusters as new efficient and environmentally safe energy sources, one should be able to create bulk metastable nitrogen structures composed of N_n ($n \geq 4$) clusters. On the one hand, in order such structures could exist, the nitrogen clusters should be connected with each other by covalent bonds. On the other hand, those bonds should not be "too strong" in order the clusters preserved their individuality and the energy accumulated in such structures was not substantially lower than the energy accumulated in an isolated cluster, i.e., the clusters should form ensembles. Besides, there should be a sufficiently high energy barrier preventing the fission of ensembles into isolated nitrogen clusters and/or nitrogen molecules.

An extensive theoretical study of Mailhot et al. [4] suggests that nitrogen can form bulk polymeric metastable structures at high pressures. However, the polymeric structures cannot be considered as composed of nitrogen clusters, i.e., as ensembles of clusters. In this paper, we present the preliminary results of numerical simulations of bonding between nitrogen clusters and show that boats N_8 can form metastable quasi-one-dimensional ensembles in which the nearest clusters N_8 are bound to each other by covalent bonds.

The version of the self-consistent-field theory with 3-21G, 6-31G, and 6-311G basis sets in HyperChem was used in the present work. First, we studied the geometry and energetics of isolated clusters N_4 (tetrahedron) and N_8 (cubane and boat). Those clusters are shown in Figs. 1-3. The accumulated energies are $E_{acc}(4) = (1.98 - 2.21)$ eV/atom for the cluster N_4 , $E_{acc}(8) = (2.44 - 2.59)$ eV/atom for the cubane N_8 , and $E_{acc}(8) = (1.53 - 1.66)$ eV/atom for the boat N_8 , depending on the basis set used. Our results agree with previous works [1, 2, 3].

We have carefully analyzed a lot of different mutual arrangements of nitrogen clusters. In both cases of N_4 and cubane N_8 clusters, we have found no metastable configurations. However, in the case of boats N_8 , we have revealed that two boats can form a "cluster molecule" $(N_8)_2$. It is shown in Fig. 4. The energy accumulated in this molecule, $E_{acc}(16) = 1.79$ eV/atom (3-21G basis set), exceeds slightly the energy accumulated in an isolated boat N_8 . We have demonstrated by direct calculations that the molecule $(N_8)_2$ is mechanically stable, i.e., all its oscillation frequencies are real, giving evidence that the configuration shown in Fig. 4 corresponds to a local minimum in the total-energy surface as a function of atomic coordinates.

In order to study the formation of large ensembles of boats N_8 , we employed a version of TBM D (tight binding molecular dynamics) method with transferable interatomic potential [5, 6] used by us earlier for carbon clusters [7, 8, 9, 10, 11] and modified for calculations on nitrogen systems. Although the TBM D method is not as rigorous as ab initio approaches, it allows one to study relatively large systems as well as to calculate directly the lifetime of the metastable state and the height of the energy barrier. For N_4 and N_8 clusters and $(N_8)_2$ "boat dimers", the results of TBM D simulations agree qualitatively with ab initio calculations. We have also shown that rather long quasi-one-dimensional ensembles of boats can be formed through attaching of new boats N_8 to the $(N_8)_2$ "molecule". The ensembles $(N_8)_m$ with m up to 6 have been shown to be metastable with accumulated energy about 90 percents of the energy $E_{acc}(8)$ accumulated in an isolated boat. The heights of the energy barriers preventing the fission of metastable configurations exceed 0.3 eV.

Finally, we note a recent success in recovering the non-molecular phase of nitrogen at ambient pressure at temperatures below 100 K [12]. Structure and physical

properties of a new phase are not understood yet. Its characterization remains an important issue. Presumably, it is polymeric. This can be considered as a first step towards synthesis of bulk metastable nitrogen structures, including "cluster phases" studied in this paper.

Acknowledgments

The work was supported by the CRDF (project B-P/252), the Contract DSW A 01-98-C-0001, and the Russian Federal Program "Integration" (Project AO 133).

References

- [1] W. J. Lauderdale, J. F. Stanton, and R. J. Bartlett, *J. Phys. Chem.*, **96** (1992) 1173.
- [2] M. L. Leininger M. L., C. D. Sherril, and H. F. Schaefer, *J. Phys. Chem.*, **99** (1995) 2324.
- [3] C. Chen and K.-C. Sun, *Int. J. Quant. Chem.*, **30** (1996) 497.
- [4] C. Mailhot, L. H. Yang, and A. K. McMahan, *Phys. Rev. B*, **46** (1992) 14419.
- [5] C. H. Xu, C. Z. Wang, C. T. Chan, and K. M. Ho, *Phys. Rev. B* **47** (1993) 9878.
- [6] C. H. Xu, C. Z. Wang, C. T. Chan, and K. M. Ho, *J. Phys.: Condens. Matter* **4** (1992) 6047.
- [7] L. A. Openov and V. F. Elesin, *Pis'ma Zh. Eksp. Teor. Fiz.*, **68** (1998) 685 [*JETP Lett.*, **68** (1998) 726].
- [8] V. F. Elesin, A. I. Podlivaev, and L. A. Openov, *Phys. Low-Dim. Struct.*, **11/12** (2000) 91.
- [9] L. A. Openov and V. F. Elesin, *Mol. Materials*, **13** (2000) 391.

- [10] N N Degtyarenko, V F Elesin, N E L'vov, L A Openov, and A I Podlivaev, Abstracts of International Workshop "Fullerenes and Atomic Clusters" IW - FAC '2001, July 2-6, 2001, St. Petersburg. p.311.
- [11] N N Degtyarenko, V F Elesin, N E L'vov, L A Openov, and A I Podlivaev, Abstracts of International Conference "Physics of Low-Dimensional Structures" PLDS-3, October 15-20, 2001, Chernogolovka, p.16.
- [12] M I Eremets, R J Hemley, H K Mao, and E Gregoryanz, Nature, 411 (2001) 170.

Figure captions

Fig.1. Tetrahedron N_4 .

Fig.2. Cubane N_8 .

Fig.3. Boat N_8 .

Fig.4. "Cluster molecule" $(N_8)_2$ composed of two boats N_8 .









