

Counterflow Superfluidity of Two-Species Ultracold Atoms in a Commensurate Optical Lattice

A. B. Kuklov

Department of Engineering Science and Physics, CUNY, Staten Island, NY 10314

B.V. Svistunov

*Department of Physics, University of Massachusetts, Amherst, MA 01003, USA and
Russian Research Center “Kurchatov Institute,” 123182 Moscow, Russia*

In the Mott-insulator regime, two species of ultracold atoms in optical lattice can exhibit the low energy counterflow motion. We construct effective Hamiltonians for the three classes of the two-species insulators—fermion-fermion, boson-boson, and boson-fermion type—and reveal the conditions when the resulting groundstate supports super-counter-fluidity (SCF), with the alternative being phase-segregation. We emphasize a crucial role of breaking the isotopic symmetry between the species for realizing the SCF phase.

PACS numbers: PACS 03.75.Fi, 05.30.Jp

Recent pioneering experiment by Greiner *et al.* [1] (proposed in ref. [2]) on the superfluid–Mott-insulator transition in the optical lattice of ultracold atoms has opened up a new era of strongly correlated atomic systems. Now it is possible to experimentally study a Mott phase of ultracold atoms. As shown in ref.[3], the spinor ($S=1$) bosons in the lattice can exhibit very rich and exotic phase diagram.

In the Mott phase of *single* specie of atoms, any low-energy transport is suppressed [4]. The situation may change dramatically, if the Mott phase includes at least *two* species. While the net number-of-atoms transport is still suppressed, the counterflow (when the currents of the two species are equal in absolute values and are in opposite directions), generally speaking, survives, and, at certain conditions, it can be non-dissipative (super-counterflow).

In this Letter, we discuss typical cases of counterflow dynamics in two-species Mott phase. We consider three classes of the commensurate two-species systems: fermion-fermion (FF), boson-boson (BB), and boson-fermion (BF) mixtures (assuming that each of the species does not possess internal degrees of freedom). We confine our analysis to the strong coupling limit, when the effective counterflow Hamiltonian is readily obtained in the second order perturbation theory in the parameter $t/U \ll 1$, where t is the hopping amplitude and U is the on-site repulsion. For the same-statistics (BB or FF) mixtures, the effective Hamiltonian can be written in terms of $s = \nu/2$ iso-spin operators (the integer ν is the filling factor). In the FF case, we have a $s = 1/2$ anti-ferromagnetic asymmetric effective Hamiltonian. In the BB case, the counterflow term is represented by asymmetric ferromagnetic interaction of iso-spins.

Depending on particular means of breaking the isotopic $SU(2)$ symmetry, either the effective Hamiltonians are of the easy-axis type, which means phase segregation, or they are of the easy-plane type resulting in the SCF

groundstate. In the BF case, it is possible to eliminate bosonic variables and obtain a spinless fermionic model with nearest-neighbor interaction. The sign of the interaction depends on the parameters of the original Hamiltonian, and how they deviate from the exact fermion-boson symmetry. In particular, the p -wave Cooper pairing is possible which results in the SCF.

Our lowest-order effective Hamiltonians will involve only on-site and nearest-neighbor interactions. Hence, to derive them it is sufficient to consider a system consisting of two strong traps, 1 and 2, between which weak tunneling is allowed. Only the lowest one-particle state φ_σ for the species $\sigma = \uparrow, \downarrow$ (we use pseudo-spin notation) in each trap is taken into account. We introduce the creation and annihilation fermionic/bosonic operators $a_{b\sigma}^\dagger, a_{b\sigma}$, respectively, where $b = 1, 2$ labels the trap (site). Our original two-site Hamiltonian then reads

$$H_{12} = H_{12}^{(0)} + V_{12}, \quad (1)$$

$$H_{12}^{(0)} = \frac{1}{2} \sum_{b\sigma\sigma'} U_{\sigma\sigma'} : n_{b\sigma} n_{b\sigma'} :, \quad (2)$$

$$V_{12} = - \sum_{\sigma} (t_{\sigma} a_{1\sigma}^\dagger a_{2\sigma} + H.c.), \quad (3)$$

where $n_{b\sigma} = a_{b\sigma}^\dagger a_{b\sigma}$, $U_{\sigma\sigma'} = g_{\sigma\sigma'} \int d\mathbf{x} |\varphi_\sigma|^2 |\varphi_{\sigma'}|^2$, $g_{\sigma\sigma'}$ is the interaction constant between species σ and σ' , and $:(\dots):$ denotes normal form of the product of the creation-annihilation operators. We consider the tunneling matrix element t_σ as real and positive and depending on the component index.

At commensurate total filling $n_b = \sum_{\sigma} n_{b\sigma} = \nu$ on each site, and in the limit $|t_\sigma| \ll U_{\sigma\sigma'}$, single particle jumps change the total on-site populations, and, therefore, require high energy. In contrast, exchanging two different particles does not require such an energy. In order to describe these processes the single particle jumps should be eliminated in the second order with respect

to the ratio t/U . Thus, we choose $H_{12}^{(0)}$ in eq. (2) as zeroth order part, while V_{12} is the perturbation. The basis in our effective Hilbert space is given by the states $|\text{eff}\rangle = |n_{1\uparrow}, n_{1\downarrow}, n_{2\uparrow}, n_{2\downarrow}\rangle$, where $n_1 = n_2 = \nu$; the virtual excited states $|\text{ex}\rangle$ (where $n_1 = n_2 \pm 1$) are excluded in the second order. The effective Hamiltonian V'_{12} can be represented in terms of the matrix elements of the original Hamiltonian as

$$(V'_{12})_{\alpha\beta} = - \sum_{\gamma} \frac{V_{\alpha\gamma} V_{\gamma\beta}}{E_{\gamma} - (E_{\alpha} + E_{\beta})/2}, \quad (4)$$

where α, β label $|\text{eff}\rangle$, and γ denotes $|\text{ex}\rangle$; E 's are the eigenenergies of $H^{(0)}$ in eq. (2).

FF case. This case is the easiest for calculation. The only possible filling factor is $\nu = 1$ and the only possible intermediate virtual state γ is the state with two different fermions on one site. The only relevant interaction vertex is $U_{\uparrow\downarrow} = U$. The Hamiltonian of the system is the standard Hubbard model [5] widely used for electrons in metal, the only specific (and significant !) feature being the dependence of the hopping amplitude on the “spin” index. The effective Hamiltonian is readily obtained from eq. (4), in terms of iso-spin operators

$$\mathbf{S}_b = (1/2) \sum_{\sigma\sigma'} a_{b\sigma}^{\dagger}(\sigma)_{\sigma\sigma'} a_{b\sigma}, \quad (5)$$

where $(\sigma)_{\sigma\sigma'}$ are the Pauli matrices. The operators $\mathbf{S}_b = (S_{bx}, S_{by}, S_{bz})$, where the second index refers to the iso-spin components, obey standard spin commutation relation at same site and commute with each other on different sites. The resulting effective two-site Hamiltonian $H'_{12} = 2J\mathbf{S}_1\mathbf{S}_2 + 2J'S_{1z}S_{2z}$ extended to a lattice is

$$H_{FF} = \sum_{\langle ij \rangle} [J\mathbf{S}_i\mathbf{S}_j + J'S_{iz}S_{jz}] - \sum_i B S_{iz}, \quad (6)$$

$$J = 2t_{\uparrow}t_{\downarrow}/U, \quad J' = (t_{\uparrow} - t_{\downarrow})^2/U. \quad (7)$$

where $\langle \dots \rangle$ stands for the pairs of nearest neighbors; $B = \mu_{\uparrow} - \mu_{\downarrow}$, and μ_{σ} is the chemical potential of the σ -th component. If the two species are, e.g., just different isotopes of one and the same element, then the condition $t_{\uparrow} \neq t_{\downarrow}$ naturally occurs from the difference of the masses.

The groundstate of this system (see, e.g., [6]) at $B = 0$ is the spin-1/2 easy-axis anti-ferromagnet, given $J > 0$, $J' > 0$ in eq. (7). This corresponds to the checkerboard insulator (Néel state). If $J' = 0$ and for infinitesimally small B , this state is unstable with respect to forming the *canted* phase, that is, the easy-plane state in which the staggered magnetization resides in the (x, y) -plane, with the ferromagnetic component $\sim B$ oriented along the z -axis. This state maps into the superfluid bosonic system (see in, e.g., [7] and also [8]), which is equivalent to the SCF. For $J' \neq 0$, the *canted* state

occurs by the first order transition at $|B| > B_c$, where $B_c \approx \sqrt{J'(2J + J')}$. In the *canonical* ensemble, when the numbers of atoms of both components, N_{\uparrow} and N_{\downarrow} , are fixed, the SCF phase corresponds to a *general* case, while the checkerboard insulator occurs only at $N_{\uparrow} = N_{\downarrow}$. As shown in ref.[8], in canonical ensemble, the checkerboard insulator state can coexist with the superfluid, with the amount of the superfluid determined by $N_{\uparrow} - N_{\downarrow} \neq 0$.

It is important to stress, that in our situation—in contrast to the case of real easy-plane anti-ferromagnet—the staggered magnetization vector fundamentally cannot be attached to any direction in the easy plane due to the exact conservation of the atoms of each component. This guarantees the exact $U(1)$ symmetry. As discussed later, the easy-plane $U(1)$ symmetry can be broken only *spontaneously* by forming the superfluid counter-flow vacuum.

BB case. In this case we also introduce the site iso-spin operators according to eq. (5) with bosonic creation-annihilation operators. The expression for the effective Hamiltonian in a general case of non-equal interactions is rather cumbersome. It becomes much simpler in two important particular cases: (i) when $U_{\sigma\sigma'} = U + \delta U_{\sigma\sigma'}$, with $|\delta U_{\sigma\sigma'}| \ll U$, and (ii) when $\nu = 1$. We start with the former case in the limit $|\delta U_{\sigma\sigma'}|/U \rightarrow 0$. Then, the denominators in eq. (4) are the same and equal to U . Hence, the intermediate virtual states γ do not involve any projection operators, and the answer for the two-sites can be written as a bilinear form of $s = \nu/2$ iso-spin operators as $H'_{12} = -J\mathbf{S}_1\mathbf{S}_2 - J'S_{1z}S_{2z} - B_{12}(S_{1z} + S_{2z})$, $B_{12} = (\nu + 1)(t_{\uparrow}^2 - t_{\downarrow}^2)/U$, where the expressions for J and J' are given by eqs. (7). When $|\delta U_{\sigma\sigma'}| \ll U$, the parameters J , J' , and B_{12} remain, to the first order in $|\delta U_{\sigma\sigma'}|/U$, the same. The main correction to the Hamiltonian is associated with the term $H_{12}^{(0)}$ (1). Then, extending the two-sites Hamiltonian $H'_{12} + H_{12}^{(0)}$ to the lattice and omitting trivial constant terms, we find

$$H_{BB} = - \sum_{\langle ij \rangle} [J\mathbf{S}_i\mathbf{S}_j + J'S_{iz}S_{jz}] + \sum_i [(D/2)(S_{iz})^2 - B S_{iz}], \quad (8)$$

$$B = \mu_{\uparrow} - \mu_{\downarrow} - \frac{1}{2}(\nu - 1)(U_{\uparrow\uparrow} - U_{\downarrow\downarrow}) + p B_{12}, \quad (9)$$

$$D = U_{\uparrow\uparrow} + U_{\downarrow\downarrow} - 2U_{\uparrow\downarrow}, \quad (10)$$

with p being the number of close neighbors on the lattice. Note that in contrast to the FF case, the minus sign now stands in front of J and J' . That is, if the D -term in eq.(8) is ignored, we arrive at the easy-axis ferromagnetic model, that implies a phase segregation. However, for large enough D in eqs.(8,10), the easy-plane ground state can be realized. The mean field condition for this (in the case $B = 0$) is $D \gtrsim pJ'$ [9].

The effective Hamiltonian for the BB situation at $\nu = 1$ can readily be found for arbitrary $U_{\sigma\sigma'}$. The summation

over γ in eq.(4) in this case is very simple, because for any different states α and β in eq. (4), there is no more than one state γ for which matrix elements differ from zero. In the diagonal terms $\alpha = \beta$, if two sites are both occupied by the same iso-spin σ bosons, the energy change due to jumping of either boson is $U_{\sigma\sigma}$; when the bosons have opposite spins, the energy change becomes $U_{\uparrow\downarrow}$. The final result acquires the form eq. (8), the parameters J , J' , and B being

$$J = 2t_{\uparrow}t_{\downarrow}/U_{\uparrow\downarrow}, \quad (11)$$

$$J' = -(t_{\uparrow} + t_{\downarrow})^2/U_{\uparrow\downarrow} + 2t_{\uparrow}^2/U_{\uparrow\uparrow} + 2t_{\downarrow}^2/U_{\downarrow\downarrow}, \quad (12)$$

$$B_{12} = 2(t_{\uparrow}^2/U_{\uparrow\uparrow} - t_{\downarrow}^2/U_{\downarrow\downarrow}). \quad (13)$$

At certain conditions, one obtains the easy-plane situation $J' < 0$, that is the SCF. A solution for this can, in principle, be found exactly. We will, however, analyze it in a simplified situation when the system is almost $SU(2)$ symmetric (like ^{87}Rb [10]), which means $U_{\uparrow\uparrow} \approx U_{\downarrow\downarrow} \approx U_{\uparrow\downarrow}$, $t_{\downarrow} \approx t_{\uparrow}$, and will choose chemical potentials to have $B = 0$. We represent $U_{\uparrow\uparrow} = U + U' + D/2$, $U_{\downarrow\downarrow} = U - U' + D/2$, $U_{\uparrow\downarrow} = U$, $t_{\uparrow} = t + t'$, $t_{\downarrow} = t - t'$, with $|U'|/U \ll 1$, $|D|/U \ll 1$, $|t'|/t \ll 1$. Then, $J' < 0$ yields

$$|t'|/t < \sqrt{D/2U}, \quad D > 0. \quad (14)$$

The easy-plane state is equivalent to the SCF.

Now let us focus on why the easy-plane ground state of the models (6, 8) can support the superfluid counterflow of the components. First, we note that, according to Holstein and Primakoff [11], spin lattice is equivalent to the lattice bosons, with the spin commutation relation being essentially equivalent to the Bose commutation relation. Then, the site operators $\hat{S}_n^+ = \hat{S}_{nx} + i\hat{S}_{ny}$ are proportional to the effective boson annihilation operators \hat{b}_n [11], and the operator $\hat{S}_{nz} = (\nu/2) - \hat{b}_n^\dagger \hat{b}_n$. Accordingly, the formation of the easy-plane order parameter $S_n^+ = \langle S_{nx} + iS_{ny} \rangle = |S'| \exp(i\varphi) \sim \langle \hat{b}_n \rangle$, where $|S'| \approx \text{const} \neq 0$, is equivalent to the formation of the Bose-Einstein condensate (BEC) $\langle \hat{b}_n \rangle \neq 0$ of the Holstein-Primakoff boson \hat{b}_n . As long as the global invariance with respect to the phase φ of the bosons holds, which is insured by the conservation of the original species, and the bosons are interacting, the system is (counter-) superfluid.

Let us obtain expression for the superfluid counter-current density \mathbf{I} , which is the difference of the currents of the each specie. To be specific, we will discuss the BB case and keep in mind that the result for the FF case is, qualitatively, the same. In the long wave limit, the conservation law of the components takes the form $\hat{S}_z + \nabla \mathbf{I} = 0$, where $S_z(\mathbf{x}) \approx S_{nz}/\Omega$ is the difference of the densities of the species, with Ω standing for the unit cell volume and \mathbf{x} being close to the site n . This continuity

equation follows from the Heisenberg equation of motion $i\dot{\hat{S}}_{nz} = [\hat{S}_{nz}, H_{BB}]$ after performing the commutations and taking the long wave limit. Finally, replacing the operators by c -numbers, we find

$$\mathbf{I} = n_{\text{scf}} \nabla \varphi, \quad n_{\text{scf}} = Jp d^2 |S'|^2 / (12\Omega), \quad (15)$$

where d denotes the nearest neighbor distance; n_{scf} has a meaning of the effective superfluid density, with $\nabla \varphi$ being the corresponding velocity. Thus, the easy-plane ground state of the models (6, 8) supports a super-counterflow of the two components. The general condition for this is that the intrinsic symmetry (i.e., $SU(2)$) between the components is broken down to the $U(1)$ group.

BF case. Now we turn to the Bose-Fermi mixture on the lattice. The Hamiltonian (1) of two sites becomes

$$H_{12}^{(0)} = \sum_{b=1,2} [(U_0/2) n_b(n_b - 1) + U_1 n_b m_b], \quad (16)$$

$$V_{12} = -(t_B a_2^\dagger a_1 + t_F c_2^\dagger c_1 + H.c.), \quad (17)$$

where a_b^\dagger , a_b and c_b^\dagger , c_b stand for the site creation and annihilation operators of the bosons and the fermions, respectively; $n_b = a_b^\dagger a_b$, $m_b = c_b^\dagger c_b$; $U_0 > 0$, $U_1 > 0$. Below we will see that the SCF is possible, if (formal) symmetry between bosons and fermions (when $U_0 = U_1$, $t_B = t_F$) is broken.

The effective two-site Hamiltonian can be written in terms of the fermionic operators only. The elimination of the bosonic operators is possible because the truncated Hilbert space is exhausted by different fermionic occupations, the bosonic occupations being unambiguously defined by the constraint $n_b + m_b = \nu$. Formally, one introduces new operators $\tilde{c}_i = a_i^\dagger c_i / \sqrt{\nu}$ and observes that, given our constraint, these satisfy standard fermionic algebra. [Note, that the vacuum for the new fermions corresponds to the Bose Mott insulator state with the filling factor ν .] The effective Hilbert space and the Hamiltonian are then automatically defined:

$$H_{BF} = \sum_{\langle ij \rangle} [-\kappa c_i^\dagger c_j + (\lambda/2) m_i m_j] - \mu_F \sum_i m_i, \quad (18)$$

(we have omitted the tilde in the fermionic operators). Here $\mu_F = \mu + p\mu_{12}$ denotes the resulting fermionic chemical potential, with μ being the difference of the bare fermion and boson chemical potentials (adding one fermion to the lattice implies simultaneous removal of one boson); the other constants are:

$$\kappa = 2\nu t_B t_F / U_1, \quad (19)$$

$$\lambda = 2t_B^2 \left(\frac{\nu^2}{U_1} + \frac{\nu^2 - 1}{2U_0 - U_1} - \frac{2\nu^2}{U_0} \right) + \frac{2t_F^2}{U_1}, \quad (20)$$

$$\mu_{12} = \frac{t_F^2 + t_B^2 \nu^2}{U_1} + \frac{t_B^2(\nu^2 - 1)}{2U_0 - U_1} - \frac{2t_B^2 \nu(\nu + 1)}{U_0}, \quad (21)$$

and it is assumed that $2U_0 - U_1 > 0$ [12].

Note that in the case of the exact boson-fermion symmetry (in the meaning defined above), $\lambda = 0$ and SCF is not possible, because the problem maps on the ideal one-component fermion gas. Thus, the symmetry must be broken. From eqs. (20), it is seen that there is a region where $\lambda < 0$. At, e.g., $U_1 = U_0$, we find $\lambda = 2(t_F^2 - t_B^2)/U_0$, and $\lambda < 0$ when $t_B > t_F$. Thus, we obtain one-component lattice fermions with the nearest neighbor attraction. This leads to the Cooper pairing (see, e.g., [13]), which in our case implies the SCF. As our fermions are effectively spinless (spin-polarized), the pairing takes place in the p -channel, that is the superconducting order parameter is characterized by broken inversion symmetry, and can exhibit broken time-reversal symmetry (see in [14]). The particular structure depends on the lattice.

Experimental realization and detection of the SCF. The essential condition for the SCF is that the system is in the MI regime with respect to the net-atomic transport. It is exactly the same condition formulated for the one component case [2]. In the FF case at $T = 0$ and $|B| > B_c$, the system will be in the SCF state. In the BB case, the three interaction constants $U_{\sigma\sigma}$ must be tuned in order to avoid the phase separation. This, however, can be achieved by creating slightly displaced lattice potentials [2] for the species in order to reduce the overlap of the states \uparrow, \downarrow on each site, so that the value $U_{\uparrow\downarrow}$ is reduced and the parameter D (10) becomes positive and large enough to insure the condition (14) for, e.g., ^{87}Rb .

It is important to note that in the cases FF and BB, once $T \neq 0$ the phase transition from the ordered (SCF) phase into the “paramagnetic” (i.e., normal with respect to the counter-flow) state may occur. A typical critical (Néel or Curie) temperature T_c is given by the effective exchange constant as $T_c \approx pJ$. Thus, the SCF requires low enough temperatures $T \leq pt^2/U$. For typical parameters employed in ref.[2], this reads as $T \leq 10^{-9}\text{K}$. In the BF case, the SCF-normal counter-fluid transition temperature is given by the same energy scale.

The SCF could be detected by observing non-dissipative exchange of the components through the optical lattice. One of the possibilities consists of separating the components initially. Then, the species will be exchanging their position in the oscillating manner. Raising T above T_c will result in the abrupt increase of the damping of these oscillations.

The existence of the phase φ (15) implies that the counter-vortex can be supported in the SCF phase. Then, the winding of the phase in the SCF, which has a boundary with the regular BEC phase of the BB components, will imprint this winding on the BEC phases,

resulting in regular BEC vortices. Detecting these vortices will, then, signal a presence of the SCF-vortex.

The order associated with the SCF is revealed in the *two*-particle density matrix. Its detection can be done by atomic scattering of fast atoms off the lattice [15]. Specifically, the cross-section of the process, when incoming fast particle of, e.g., sort \uparrow strikes the atoms in the lattice and “transforms” into the outgoing particle \downarrow , is exactly given by the (iso) spin-spin correlators of the atoms in the lattice. This issue will be considered in greater detail elsewhere.

Finally, let us discuss a formation of the SCF vortex in a rotating lattice. To be specific, we will discuss the BB case. The lattice rotation can induce such vortices, if the components have different masses M_σ . Indeed, the total mass current can be expressed as $\mathbf{I}^{(M)} = (M_\uparrow - M_\downarrow)n_{\text{scf}}\nabla\varphi$, where n_{scf} , φ define the counter-current (15). External rotation of the lattice at some angular velocity ω changes the energy of the counter-vortex $\sim n_{\text{scf}}L\ln(L/d)$, where L stands for a typical system size, by $\sim \pm(M_\uparrow - M_\downarrow)n_{\text{scf}}L^3\omega$. The formation of the vortex becomes energetically favorable at [13]

$$|\omega| \geq \omega_c \approx L^{-2} \ln(L/d)/|M_\uparrow - M_\downarrow|. \quad (22)$$

Note that differences in masses can result in the phase separation induced by gravity/rotation. However, if the atomic magnetons of different species are different, the corresponding linear and oscillator fields can be compensated by adding non-uniform magnetic field.

Summarizing, we have demonstrated that there is a large variety of strongly correlated groundstates in the two-component system of ultracold atoms in an optical lattice, even if the system is in the Mott-insulator regime with respect to the net number-of-atoms transport. The most dramatic effect that may occur is the so-called super-counter-fluidity, when the system supports a non-dissipative counterflow of the two components. In the strong-coupling limit, the effective Hamiltonian for the super-counterfluids corresponds to: (i) easy-plane (*canted*) anti-ferromagnet (in the case of fermion-fermion mixture), (ii) easy-plane ferromagnet (in the case of boson-boson mixture), and (iii) p -wave superconductor (in the case of boson-fermion mixture). The experimental conditions and means of detecting the super-counterfluidity are outlined.

Authors acknowledge useful discussions of the results with David Schmeltzer, Nikolay Prokof'ev and Manfred Sigrist. ABK was supported by CUNY grant PSC-63499-0032. BVS acknowledges a support from Russian Foundation for Basic Research under Grant 01-02-16508, from the Netherlands Organization for Scientific Research (NWO), and from the European Community under Grant INTAS-2001-2344.

-
- [1] M. Greiner *et al.*, Nature, **415**, 39 (2002).
- [2] D. Jaksch *et al.*, Phys. Rev. Lett. **81**, 3108 (1998).
- [3] E. Demler and F. Zhou, Phys. Rev. Lett. **88**, 163001 (2002).
- [4] M. Ma *et al.*, Phys. Rev. B **34**, 3136(1986); M.P.A. Fisher *et al.*, Phys.Rev. B **40**, 546 (1989).
- [5] J. Hubbard, Proc. Roy. Soc., **A276**, 238 (1963); **A277**, 237 (1964); **A281**, 401 (1964).
- [6] C. Kittel, *Introduction to Solid State Physics*, John Wiley&Sons, 1986.
- [7] S. Sachdev, *Quantum Phase Transitions*, Ch.13.4.3, Cambridge University Press, 1999.
- [8] G. Schmid, et al, cond-mat/0110024; F. Hébert et al., Phys. Rev. B **65**, 014513 (2002).
- [9] The only reservation should be made for the special case of commensurate filling of the lattice by *each* of the species, when strong enough D leads to a Mott-insulator transition in the iso-spin system. Indeed, exchange of two bosons \uparrow and \downarrow between two sites costs, then, energy D . Accordingly, the condition $|t_{\uparrow}| \sim |t_{\downarrow}| \gtrsim D$ should hold.
- [10] D.S. Hall *et al.*, Phys. Rev. Lett. **81**, 1543 (1998); D. S. Hall *et al.*, Phys. Rev. Lett. **81**, 1539 (1998).
- [11] T. Holstein and H. Primakoff, Phys.Rev. **58**, 1098 (1940).
- [12] Otherwise, for $\nu > 1$, the ground state of (16) will have $\nu - 1$ particles on a site, containing fermion, and $\nu + 1$ bosons on the other site with no fermion. In this situation, no transport of the fermion is possible in the second order in t_B , t_F .
- [13] E.M. Lifshitz and L.P. Pitaevskii, *Statistical Physics. Part 2*, Pergamon, Oxford, 1980.
- [14] M. Sigrist and K. Ueda, Rev. Mod. Phys. **63**, 239 (1991); Y. Maeno, T.M. Rice, and M. Sigrist, Physics Today **54**, 42 (2001).
- [15] A.B. Kuklov and B.V. Svistunov, Phys. Rev. A **60**, R769 (1999).