

## Aging and Energy Landscapes: Application to Liquids and Glasses

Stefano Mossa,<sup>1,2</sup> Emilia La Nave,<sup>1</sup> Francesco Sciortino,<sup>1</sup> and Piero Tartaglia<sup>1</sup><sup>1</sup>Dipartimento di Fisica and INFN UdR and Center for Statistical Mechanics and Complexity,  
Università di Roma "La Sapienza" Piazzale Aldo Moro 2, I-00185, Roma, Italy<sup>2</sup>Center for Polymer Studies and Department of Physics,  
Boston University, Boston, Massachusetts 02215

(Dated: April 14, 2024)

The equation of state for a liquid in equilibrium, written in the potential energy landscape formalism, is generalized to describe out-of-equilibrium conditions. The hypothesis that during aging the system explores basins associated to equilibrium configurations is the key ingredient in the derivation. Theoretical predictions are successfully compared with data from molecular dynamics simulations of different aging processes, such as temperature and pressure jumps.

Glasses are out-of-equilibrium (OOE) systems, characterized by slow dynamical evolution (aging) and history dependent properties. Their thermodynamic and dynamic properties depend on the preparation method (cooling or compression schedules) as well as the time spent in the glass phase [1]. The extremely slow aging dynamics has often been considered an indication that a thermodynamic description of the glass state can be achieved by adding one or more history-dependent parameters to the equation of state (EOS) [2, 3, 4, 5, 6, 7, 8, 9, 10, 11]. Recent theoretical work [mostly based on mean field models of structural glasses, where analytic solutions of the OOE dynamics can be explicitly worked out [12]] supports such possibility.

In this Letter we derive an EOS for OOE conditions, based on a generalization of the potential energy landscape (PEL) thermodynamic approach [13, 14, 15]. The EOS is derived under the hypothesis that during aging the system explores basins associated with equilibrium configurations, a condition which, as shown later, is simple to implement in the PEL context. The proposed EOS for OOE conditions depends on one additional parameter, which for example can be chosen to be the average depth of the explored local minima of the PEL (the so-called inherent structures (IS) [13, 14]). The EOS allows for the first time a detailed comparison between predictions and "exact results" from out-of-equilibrium molecular dynamics simulations. We report such a comparison for a realistic model of the fragile liquid orthoterphenyl, one of the most studied glass forming liquids [16].

The PEL formalism [13, 14] allows for a clear separation of vibrational and configurational contributions in thermodynamic quantities. Indeed, the equilibrium liquid free energy  $F(T; V)$  at temperature  $T$  and volume  $V$  is expressed as a sum of a vibrational contribution  $f_{\text{vib}}$ , representing the intra-basin free energy [13], and a configurational contribution ( $TS_{\text{conf}} + e_{\text{IS}}$ ). Here the configurational entropy  $S_{\text{conf}}$  accounts for the number of

explored basins of the PEL of average energy depth  $e_{\text{IS}}$ .  $f_{\text{vib}}$ , for a system of  $N$  atoms, depends on the curvature of the potential energy surface, i.e., on the  $3N$  eigenvalues  $\lambda_i^2$  of the Hessian matrix evaluated at the IS. Recently, calculating the  $V$  derivative of  $F(T; V)$ , an analytic EOS has been derived, based only on the statistical properties of the landscape [17, 18].

The separation in vibrational and configurational parts can be carried out for several properties. In our case, it is particularly useful to separate the instantaneous pressure as sum of two contributions: the pressure felt by the system in the inherent structure configuration ( $P_{\text{IS}}$ ), and an additional contribution related to the finite temperature, which we refer to as the vibrational contribution ( $P_{\text{vib}}$ ) [19]

$$P(T; V; e_{\text{IS}}) = P_{\text{IS}}(e_{\text{IS}}; V) + P_{\text{vib}}(T; V; e_{\text{IS}}): \quad (1)$$

In equilibrium  $e_{\text{IS}}$  is a well defined function of  $T$  and  $V$  and, as well known,  $P$  depends only on  $T$  and  $V$ . Note that  $P_{\text{IS}}$ , being the pressure experienced by the liquid in the IS, depends only on the basin's depth and on  $V$ . Instead,  $P_{\text{vib}}$  depends on the basin's depth as well as on  $T$  and  $V$  [20].

For models for which the statistical properties of the PEL, i.e., the distribution of shapes and numbers of IS [17, 18] are known,  $P_{\text{IS}}$  and  $P_{\text{vib}}$  can be calculated theoretically. In other cases,  $P_{\text{IS}}$  and  $P_{\text{vib}}$  can be calculated numerically. In all cases, an landscape-based EOS based on Eq. (1) can be derived.

The separation of the pressure in two contributions is crucial for the possibility of extending the equilibrium EOS to OOE conditions, since it provides a direct way to rebuild the pressure once  $T$ ,  $V$  and  $e_{\text{IS}}$  are known. In the PEL formalism, the hypothesis that the system explores basins associated with typical equilibrium configurations, i.e., configurations sampled in equilibrium conditions, is equivalent to assuming that the relations calculated in equilibrium linking  $P_{\text{IS}}$  and  $P_{\text{vib}}$  to  $e_{\text{IS}}$  and  $V$  are valid also in OOE conditions. If this is the case, the knowledge of  $e_{\text{IS}}$ ,  $T$  and  $V$  is sufficient to calculate both  $P_{\text{IS}}$ ,  $P_{\text{vib}}$  and their sum  $P$  according to Eq. (1). Similarly, the values of  $P$ ,  $T$  and  $e_{\text{IS}}$  are sufficient to predict  $V$ , since both  $P_{\text{IS}}(e_{\text{IS}}; V)$  and  $P_{\text{vib}}(e_{\text{IS}}; T; V)$  can be estimated

Present address: Laboratoire de Physique Théorique des Liquides, Université Pierre et Marie Curie, 4 place Jussieu, Paris 75005, France




FIG. 1: (a) Equilibrium relation between basin pressure  $P_{IS}$  and basin depth  $e_{IS}$ . Symbols are simulation results at the five volumes per molecule  $0.337 \text{ nm}^3$  (down triangles),  $0.345 \text{ nm}^3$  (circles),  $0.353 \text{ nm}^3$  (diamonds),  $0.361 \text{ nm}^3$  (up triangles), and  $0.369 \text{ nm}^3$  (squares) from Ref. [21]. Dashed lines are calculated using the parameterization discussed in Ref. [17]. Results at four intermediate volumes are also shown. The local minima are calculated using standard minimization algorithms [21]. (b)  $P_{vib}$  as a function of  $T$  for the same five volumes. Each curve has been shifted by  $n = 20 \text{ MPa}$  to avoid overlaps. Dashed lines show a suitable parameterization of the  $e_{IS}$ ,  $V$  and  $T$  dependence of  $P_{vib}$ , as discussed in Ref. [17].

as a function of  $V$ . The predicted  $V$  is the value for which  $P_{IS}(e_{IS}; V) + P_{vib}(e_{IS}; T; V)$  matches the external pressure.

Next we apply the ideas discussed above to a system of  $N = 343$  molecules interacting via the Lewis and Wanstrom (LW) model for the fragile molecular liquid orthoterphenyl (OTP) [21, 22]. The LW model is a rigid three-site model, with intermolecular site-site interactions modeled by the Lennard-Jones (LJ) potential [22]. We refer to Ref. [21] for all numerical details. An equilibrium EOS, based on the assumption of a Gaussian landscape, has been recently presented for this model [17], and successfully compared with the "exact" EOS calculated from molecular dynamics simulations.

Fig. 1 shows the relation between  $P_{IS}$  and  $e_{IS}$  in equilibrium along different isochores and  $P_{vib}$  as a function of  $T$  for the LW model. Both quantities can be parameter-




FIG. 2: Paths in the  $P$  -  $T$  plane of the four OOE simulations discussed in the text. Arrows indicate: (i) a  $T$ -jump at constant  $V$  between liquid states; (ii) a  $P$ -jump at constant  $T$  between liquid states; (iii) a  $P$ -jump at constant  $T$  starting from a glass; (iv) a constant  $P$  heating of a glass.  $P$  and  $T$ -jumps have been simulated by changing the thermostat and barostat value at  $t = 0$ . The time constant of the thermostat and barostat has been fixed to 20 ps. The full lines indicate the equilibrium  $P(T)$  at various  $V$  values, from Ref. [21]. The dashed lines connect the equilibrium state points and the glass states obtained by fast constant- $V$  cooling.

ized in a quite accurate way. Indeed, for this model,  $P_{vib}$  is weakly dependent on  $V$  and  $e_{IS}$  and it is well represented by a linear  $T$ -dependence. The parameterization of  $P_{IS}$  and  $P_{vib}$  offers the possibility of estimating the system pressure once  $T$ ,  $V$  and  $e_{IS}$  are known.

We study four different OOE protocols via computer simulation. The imposed external conditions are illustrated in Fig. 2. We study i) a  $T$ -jump at constant  $V$ ; ii) a  $P$ -jump at constant  $T$ ; iii) a  $P$ -jump at constant  $T$  in the glass phase and iv) an isobaric heating of a glass. For the cases i) and ii), conditions are chosen in such a way that within the simulation time window (50 ns) the system reaches the equilibrium state. In all studied cases, averages over 50 independent simulations have been performed.

Case (i). | Fig. 3 shows the comparison between the theoretical predictions and the numerical results for the constant- $V$   $T$ -jump case. The  $e_{IS}$  values, calculated from the simulation (Fig. 3(a)), are used together with the values of  $T$  and  $V$  as input to predict the evolution of  $P$  (Fig. 3(b)). Figs. 3(c) and (d) also show the comparison for  $P_{IS}$  and  $W$ .  $W(e_{IS})$ , defined as  $\frac{3N}{i=1} \ln(|\epsilon_i(e_{IS})|)$ , provides an indication of the similarity in basin shape (at the level of the harmonic approximation) between basins explored in equilibrium and in OOE conditions. The quality of the predictions indeed supports the validity of the EOS and the hypothesis that the basins sampled during aging have the same relationship  $W(e_{IS})$  as in equilibrium.

Case (ii). | Fig. 4 shows the evolution of the system after a constant- $T$  pressure jump. In this case, the  $e_{IS}$




FIG. 3: Dynamics after a T-jump, at volume per molecule  $V = 0.345 \text{ nm}^3$ , from 480 K to 340 K. (a)  $e_{IS}$ ; (b)  $P$ ; (c)  $P_{IS}$ ; (d)  $W$  ( $l_o = 1 \text{ cm}^{-1}$ ). Open symbols are simulation results, closed circles are theoretical predictions based on the OOE PEL-EOS and using as input the data shown in panel (a). Dashed lines indicate the known equilibrium values at the initial and final state. The error in the prediction at short and long times, when equilibrium conditions are met, provides a measure of the quality of the equilibrium EOS [17], which agrees with the simulation data within 5 MPa.

values calculated from the simulation are used together with the values of  $T$  and  $P$  as input to successfully predict the evolution of  $V$ . For times shorter than 20 ps, the barostat time constant, the fast increasing external pressure forces the system to change the volume with a solid-like response. In this time window, the initial basins in conformation space are continuously deformed by the rapid volume change, in agreement with the findings of Ref. [23]. For time longer than 20 ps, when the external pressure has reached the equilibrium value, the system starts to explore basins different from the (deformed) original ones, and the time evolution becomes controlled by the aging kinetics. To support this interpretation we show in Fig. 4(e) the actual path in the  $e_{IS}$ - $P_{IS}$  plane, where the change around 20 ps is clearly evidenced.

Cases (iii) and (iv). | The last two cases are respectively a constant  $T$   $P$ -jump and a heating at constant  $P$  (with an heating rate of  $1 \text{ K/ps}$ ), both starting from glass conformations. These initial conformations are generated by rapid constant- $V$  quenches of equilibrium conformations, as shown schematically in Fig. 2. Fig. 5 shows the comparison between the theoretical predictions and the numerical calculations for  $V$ , using  $e_{IS}$  as input. In the case of a  $P$ -jump (Fig. 5-Top), again two different dynamical behaviors are observed: a fast dynamics pro-




FIG. 4: Dynamics after a  $P$ -jump at  $T = 320 \text{ K}$  from 13.4 MPa to 60.7 MPa. (a)  $e_{IS}$ ; (b)  $V$ ; (c)  $P_{IS}$ ; (d)  $W$ . Open symbols are simulation results, closed circles are theoretical predictions based on the OOE PEL-EOS, using as input the data shown in panel (a). Horizontal dashed lines indicate the known equilibrium values at the initial and final state. The vertical dashed lines indicate the time at which the external pressure reaches the final value. Panel (e) shows the actual path of the aging process in the  $P_{IS}$ - $e_{IS}$  plane.

cess describing the mechanical response of the glass to the external pressure change, followed by an extremely slow aging dynamics, during which basin changes are taking place. In both cases and for all times, the out-of-equilibrium EOS is able to predict quantitatively the  $V$  changes.

In summary, we have derived an extension of the PEL equation of state to model the thermodynamic properties of liquids under out-of-equilibrium conditions. Such an




FIG. 5: Top: Simulation of a P-jump from 270.1 MPa to 115.4 MPa, starting from a glass conformation, obtained by quickly cooling to 70 K equilibrium conformations at volume per molecule  $V = 0.369 \text{ nm}^3$  and  $T = 170 \text{ K}$ . Bottom: Iso-baric heating, starting from the IS at  $V = 0.337 \text{ nm}^3$  per molecule and  $T = 320 \text{ K}$ . The heating rate is  $1 \text{ K / ps}$ . Circles are the  $e_{\text{IS}}$  values used as input to predict the evolution of  $V$  (lines). Squares are the  $V$  calculated directly from the simulations.

extension | based on the hypothesis that an aging system evolves through regions of conformation space which are typically sampled in equilibrium | requires an additional thermodynamic parameter which, for convenience, we have associated with the IS energy of the explored basins.

The results reported in this Letter (Figs.3-5) show that the proposed generalization of the PEL equation of state is successful in predicting out-of-equilibrium thermodynamics, at least under the conditions and time scales probed by state-of-the-art numerical simulations. Larger  $T$  or  $P$  jumps and/or longer aging time could require a more detailed thermodynamic description, with more than one additional parameter. Despite such a possibility, the energy landscape approach developed in this Letter is a promising starting point for looking into more complex problems in the physics of OOE liquids and glasses.

We thank A. Angell, P. Debenedetti, T. Keyes, L. Leuzzi, G. McKenna, G. Ruocco, S. Sastry, A. Scala, and M. Yamada for discussions and comments. We acknowledge support from MUR-COFIN 2000, INFN-PRA and INFN Initiative Parallel Computing.

- 
- [1] P. G. Debenedetti, *Metastable liquids* (Princeton Univ. Press, Princeton, 1996).
  - [2] A. Q. Tool, *J. Am. Ceram. Soc.* 29, 240 (1946).
  - [3] R. O. Davies and G. O. Jones, *Adv. in Physics* 2, 370 (1953).
  - [4] A. J. Kovacs, *Fortschr. Hochpolym.-Forsch.* 3, 394 (1964).
  - [5] G. B. McKenna, in *Comprehensive Polymer Science*, Vol. 2, ed. by C. Booth and C. Price, Pergamon, Oxford, 311 (1989).
  - [6] R. J. Speedy, *J. Chem. Phys.* 100, 6684 (1994).
  - [7] L. F. Cugliandolo, et al., *Phys. Rev. E* 55, 3898 (1997).
  - [8] Th. M. Nieuwenhuizen, *Phys. Rev. Lett.* 80, 5580 (1998).
  - [9] A. C. Angell, et al., *J. Appl. Phys.* 88, 3113 (2000).
  - [10] S. Franz and M. A. Virasoro, *J. Phys. A* 33, 891 (2000).
  - [11] L. Leuzzi and Th. M. Nieuwenhuizen, *J. Phys.: Condens. Matter* 14, 1637 (2002).
  - [12] L. F. Cugliandolo, et al., *Phys. Rev. Lett.* 76, 2390 (1996).
  - [13] F. H. Stillinger and T. A. Weber, *Phys. Rev. A* 25, 978 (1982).
  - [14] P. G. Debenedetti and F. H. Stillinger, *Nature* 410, 259 (2001).
  - [15] S. Sastry, *Nature* 409, 164 (2001).
  - [16] A. Tolle, *Rep. Prog. Phys.* 64, 1473 (2001).
  - [17] E. La Nave, et al., *Phys. Rev. Lett.* 88, 225701 (2002).
  - [18] P. G. Debenedetti, et al., *Adv. in Chem. Eng.* 28, 21 (2001).
  - [19] In equilibrium,  $P$  can be unambiguously separated in a conformational  $@_V [T S_{\text{conf}}(T; V) - e_{\text{IS}}(T; V)]$  and a vibrational  $@_V f_{\text{vib}}$  contribution. In the IS conformation, the vibrational part is strictly zero. Still, the conformational component may not coincide with  $P_{\text{IS}}$ . Different proposals for  $P_{\text{IS}}$  arise depending on the meaning of  $T$  in the  $T S_{\text{conf}}$  term above. See Refs. [17, 24]. Such ambiguity affects the theoretical expression for  $P_{\text{IS}}$  based on the statistical properties of the landscape. Luckily, this ambiguity does not hamper the possibility of developing an OOE-EOS when  $P_{\text{IS}}$  is calculated directly from the simulation data.
  - [20] For example, in an harmonic solid,  $P_{\text{vib}}$  is related to the  $V$  derivative of the eigenfrequencies as  $P_{\text{vib}}(T; V; e_{\text{IS}}) = k_B T @_V \sum_{i=1}^{3N-3} \ln(\sim!_i(e_{\text{IS}}))$ .
  - [21] S. Mossa, et al., *Phys. Rev. E* 65, 041205 (2002).
  - [22] L. J. Lewis and G. Wahnstrom, *Phys. Rev. E* 50, 3865 (1994).
  - [23] D. L. Malandro and D. J. Lacks, *Phys. Rev. Lett.* 81, 5576 (1998).
  - [24] F. Sciortino and P. Tartaglia, *Phys. Rev. Lett.* 86, 107 (2001).