

Energy Bounds of Linked Vortex States

A. P. Protogenov, V. A. Verbus

Institute of Applied Physics of the RAS, 603950 Nizhny Novgorod
Institute for Physics of Microstructures of the RAS, 603950 Nizhny Novgorod

22nd March 2024

Abstract

Energy bounds of knotted and linked vortex states in a charged two-component system are considered. It is shown that a set of local minima of free energy contains new classes of universality. When the mutual linking number of vector order parameter vortex lines is less than the Hopf invariant, these states have lower-lying energies.

PACS: 74.20.De, 74.20.Mn, 11.15.Tk, 11.10.Lm, 11.27.+d

A tangle of vortex elements is a system which attracts attention due to several reasons. Along with the coherent state, which is the background of this vortex field distributions, the element system also contains a disorder combination due to free motion of its fragments and a topological order because of the effects of knotting and linking [1, 2] of its separate parts.

The study of soft condensed matter, whose universal behavior is determined by topological characteristics, is recognized as one of the most challenging problems of modern condensed matter physics [3]. The aim of this paper is to find the energy bounds for vortex states with a set of numbers determining the knotting and linking degree of the fields that take part in the description of the coherent state. We will use the Ginzburg-Landau model

$$F = \int d^3x \left[\frac{1}{2m} (\nabla_k \psi + i \frac{2e}{c} A_k \psi)^2 + \frac{1}{2} (b_j \psi^2 + \frac{c}{2} j^2 \psi^4 + \frac{B^2}{8}) \right] \quad (1)$$

with a two-component order parameter

$$\psi = \frac{1}{\sqrt{2m}} \sum_j \psi_j e^{i\phi_j} ; \quad \mathbf{j} = \sum_j \mathbf{j}_j e^{i\phi_j} ; \quad (2)$$

satisfying the CP^1 condition, $j_1 \mathbf{j}_1 + j_2 \mathbf{j}_2 = 1$. This model is used in the context of the two-gap superconductivity [4] and in the non-Abelian field theory [5, 6].

It has been shown in paper [4] that there exists an exact mapping of the model (1), (2) into the following version of n-field model:

$$F = \int d^3x \left[\frac{1}{4} (\partial_k n)^2 + (\partial_k \phi)^2 + \frac{1}{16} c^2 + (F_{ik} - H_{ik})^2 + V(n, \phi) \right] ; \quad (3)$$

e-mail: aprot@applsci-nnov.ru

Setting in Eq.(3) $c = 0$ we get Faddeev-Niemi model [7]. The numerical study of the knotted configurations of n -eld in this model has been done in [8, 9, 10]. The lower energy bound in this case

$$F > 32 \quad 2 \quad 0 \quad 3=4 \quad (4)$$

is determined ([11, 12, 13]) by the Hopf invariant,

$$Q = \frac{1}{16^2} \int d^3x \, {}^{(3)}\mathcal{L}_{\text{ADM}}(a_i, \partial_k a_i) : \quad (5)$$

At compactification $R^3 \rightarrow S^3$ and $n \geq 2$ S^2 , the integer $Q = \pi_3(S^2) = \mathbb{Z}$ shows the degree of linking or knotting of lamental manifolds, where the vector field $n(x; y; z)$ is defined. In particular, for two linked rings (Hopf linking) $Q = 1$, for the trefoil knot $Q = 6$ and etc. Significant point is the following: $\pi_3(CP^M) = 0$ at $M > 1$ and $\pi_3(CP^1) = \pi_3(S^2) = \mathbb{Z}$ [14]. In the latter case the order parameter (2) is two-component one [4] and linked or knotted soliton configurations are labeled by the Hopf invariant (5). In the $(3+0)D$ case of the free energy (3), Hopf invariant (5) is analogous to the Chern-Simons action ($k=4$) $\int d^4x \epsilon^{\mu\nu\rho\sigma} a_\mu \partial_\nu a_\rho \partial_\sigma a_\sigma$ determining strong correlations of $(2+1)D$ modes [15, 16] at semion value $k' = 2$.

Let us assume that β can be found from the minimal value of the potential $V(\beta)$, but the velocity c does not equal zero. Equation (3) in this case has the following form:

$$\mathbf{F} = \mathbf{F}_n + \mathbf{F}_c \quad \mathbf{F}_{int} = \int_{\Omega} d^3x \quad (\partial_k n)^2 + H_{ik}^2 + \frac{1}{4}C^2 + F_{ik}^2 - 2F_{ik}H_{ik} \quad : \quad (6)$$

It is seen from Eq.(6) that a superconducting state with $c \neq 0$ has the energy which is less than the minimum in Eq.(4) due to renormalization of the coefficient ($= 1$) in the second term of the functional F_n . To find the lower free energy bound in the superconducting state with $c \neq 0$ we will use the following inequality:

$$F_n^{5=6} F_c^{1=2} > (32^{-2})^{4=3} \text{LJ}; \quad (7)$$

where

$$L = \frac{1}{16^2} \int d^3x \, {}^{(3)}C_{ikl} C_l @_k a_1 \quad (8)$$

is the degree of mutual linking of the velocity c lines and of the magnetic field $H = [r, a]$ lines. It is also an integral of motion [17].

The proof of the inequality (7) is based on employ of the Hölder inequality chain $\|f\|_6 \|g\|_6 \leq \|f\|_{\frac{6}{5}} \|g\|_6$, $\|g\|_6^2 = \|g\|_{\frac{6}{2}}^2$ and on the Ladyzhenskaya [8] inequality $\|g\|_6 \leq C \|g\|_{\frac{6}{5}}^{1-\epsilon} \|g\|_{\frac{6}{5}}^\epsilon$, as well as on the results of the paper [13]. Here $\|f\|_p = (\int_{\Omega} |f|^p dx)^{1/p}$. It follows also from the Schwartz-Cauchy-Bunyakovsky inequality that

$$2F_n^{1=2} F_c^{1=2} > F_{int} : \quad (9)$$

Setting the boundary value F_{int} from Eq.(9) into Eq.(6), we get $F_{min} = (F_n^{1=2} - F_c^{1=2})^2$. Using the minimum value $32^{-2} D_j^{3=4}$ of the function F_n and the Eq.(7) we have finally

$$F > 32^{-2} D_j^{3=4} (1 - j D_j^2) : \quad (10)$$

The inequality (10) is the main result of the paper. The trivial case $Q = 0$ should be considered after the limit $L = 0$. Let us pay also an attention to the self-dual relation $F_n = F_c$ which follows from F_{min} .

It follows from Eq.(10) that for all numbers $L < Q$ the energy of the ground state is less than that in the model described in [7], for which the inequality (4) is valid. The origin of the energy decrease can be easily understood. Even under the conditions of the existence of the paramagnetic part j of the current J , the diamagnetic interaction in the superconducting state consumes its own current energy and a part of the energy relating to the n -eld dynamics for all state classes with $L < Q$.

The case $\epsilon \neq \text{const}$ both for $c = 0$ [19] and $c \neq 0$ arouses certain interest. It is more complicated due to some reasons and will be considered in a separate paper. We only mention that in a soft case $\epsilon \neq \text{const}$ we need to have the proof of the compatibility of the peculiar value of the coefficient in r.h.s of Eq.(7) with stability condition [13] with respect to dilatation transformations. The equality in Eq.(12) under this remark should be understood as an ideal limit depending on topological characteristics of knots and links only.

In conclusion, we have found the energy bounds of the superconducting states using CP^1 version of Ginzburg-Landau model under the conditions of the existence of linking and knotting phenomena of the n - and c -elds being the gauge invariant order parameters of the considered system. We have shown that the energy space of the local minimum contains new state classes with $L < Q$.

We would like to thank A.G. Abanov, L.D. Faddeev, E.A. Kuznetsov and G.E. Volovik for advices, V.F. Gantmakher for the crucial remark, and G.M. Fraiman, A.G. Litvak, V.A. Mironov for useful discussions. This work was supported in part by the RFBR under the grant # 01-02-17225.

References

- [1] V.M.H. Ruutu, U. Parts, J.H. Koivumäki, M. Krusius, E.V. Thuneberg, and G.E. Volovik, JETP Lett. 60, 671 (1994).
- [2] Yu.G. Makhlin, T.Sh. Misirpashaev, JETP Lett. 61, 49 (1995).
- [3] V. Katrich, J. Bednar, D. Michoud, R.G. Scharein, J. Dubochet, and A. Stasiak, Nature 384 142 (1996).
- [4] E. Babaev, L.D. Faddeev, A.J. Niemi, Phys. Rev. B 65, 100512 (2002); cond-mat/0009438.
- [5] L.D. Faddeev, A.J. Niemi, Phys. Lett. B 525, 195 (2002).
- [6] Y.M. Cho, Phys. Rev. Lett. 87, 252001 (2001); hep-th/0110076.
- [7] L.D. Faddeev, A.J. Niemi, Nature 387, 58 (1997).

- [8] J.Gladikowski, M.Helmut, Phys.Rev.D 56, 5194 (1997).
- [9] R.A.Battye, P.M.Sutcliffe, Phys.Rev.Lett. 81, 4798 (1998).
- [10] J.Hietarinta, P.Salo, Phys.Lett.B 451, 60 (1999).
- [11] A.F.Vakulenko, L.V.Kapitansky, Sov.Phys.Dokl. 24, 433 (1979).
- [12] A.Kundu, Yu.P.Rubakov, J.Phys.A 15, 269 (1982).
- [13] R.S.Ward, Nonlinearity 12, 1 (1999), hep-th/9811176.
- [14] A.G.Abanov, P.W.Wiegmann, Geometrical phases and quantum numbers of solitons in nonlinear sigma-models, hep-th/0105213.
- [15] L.A.Abramyan, A.P.Protogenov, V.A.Verbus, JETP Lett. 69, 887 (1999).
- [16] A.P.Protogenov, JETP Lett. 73, 255 (2001).
- [17] V.E.Zakharov, E.A.Kuznetsov, Phys.Usp. 40, 1087 (1997).
- [18] O.A.Ladyzhenskaya. The mathematical theory of viscous incompressible flow, Gordon and Breach, 1969.
- [19] M.Lubcke, S.M.Nasir, A.Niemai, and K.Torokko. Twisted vortices in a gauge theory, hep-th/0106102.