## C harge D ensity B ounds in Superconducting States of Strongly C orrelated System s

A lexander P. Protogenov

Institute of Applied Physics of the RAS, Nizhny Novgorod 603950

Charge density bounds of knotted and linked vortex states in two-component G inzburg-Landau m odel are considered. W hen the m utual linking number of vector order parameter vortex lines is less than the H opf invariant, these states have the lower-lying energies. It is shown that a set of local m inim a of free energy contains new classes of universality which exist in a hole density range limited at both nite ends.

PACS numbers: 7420De, 7420Mn, 11.15.Tk, 11.10Lm, 11.27.+d

Introduction. { A tangle of vortex lam ents is a system which attracts attention due to several reasons. A long with the coherent state, which is the background of this vortex eld distributions, the lam ent system also contains a disorder combination due to free motion of its fragm ents and a topological order because of the e ects of knotting and linking of its separate parts. The degree of this order can be characterized by the linking num ber of vortices. Persistent linked eld con gurations along with a large value of interaction energy present the basic content of what we mean when talking about the origin of strong correlations.

The study of soft condensed m atter, whose universal behavior is determined by topological characteristics, is an active area of research in physics [1, 2, 3, 4] and beyond [5]. Indeed, the dynam ics of entangled vortex states is recognized as one of the most challenging problem s of m odern condensed m atter physics. The emergence of novel analytical m ethods [2, 3, 4] to treat nonlinear eld equations create now an unique opportunity to understand the dynam ics of entangled vortex states.

The vortex dynam ics can be understood in detail in the fram ework of basic models, such as the G inzburg-Landau functionalorm ore complicated multi-component system s. M ain theoretical approaches for analyzing universalbehavior in these models are based on the methods of the topological eld theory [6] and nonlinear dynam ics [7]. The universality classes are de ned both by sym metry and topological characteristics of the background nonlinear elds, while a system with in nite number of degrees of freedom contains in the ground state only the nite num ber of sym metrically invariant physical states.

The aim of this paper is to elucidate the physical mechanism leading to form ation of the charge density range bounds for superconducting states with a set of numbers determ ining the knotting and linking degree of the elds that take part in the description of the coherent state. The use of the G inzburg-Landau functional and the methods of the topological gauge eld theory gives us the hope that the obtained answers are universal. A combination of novel methods will make it possible to advance in description of the processes which are determined by the contribution of topological excitations. Model. { Wewilluse the Ginzburg-Landau free energy

$$F = \frac{Z}{d^{3}x} \frac{X}{2m} \frac{1}{2m} \sim \theta_{k} + i\frac{2e}{c}A_{k}$$
$$+ \frac{X}{bj} \frac{1}{j} + \frac{c}{2}j \frac{4}{j} + \frac{B^{2}}{8}$$
(1)

with a two-component order parameter

$$= \frac{p}{2m}$$
;  $= j j e^{i'}$ ; (2)

satisfying the CP<sup>1</sup> condition,  $j_1 f + j_2 f = 1$ . This model is used in the context of the two-gap superconductivity [8] and in the non-Abelian eld theory [9, 10].

It has been shown in paper [8] that there exists an exact mapping of the model (1), (2) into the following version of n - eld m odel:

$$F = \int_{-\infty}^{2} d^{3}x \frac{1}{4} \left[ \left( \left( \theta_{k} n \right)^{2} + \left( \left( \theta_{k} \right)^{2} + \frac{1}{16} \right)^{2} c^{2} + \left( F_{ik} - H_{ik} \right)^{2} + V(; n_{3}) \right] ; (3)$$

To write down Eq. (3), dimensionless units and gauge invariant order parameter elds of the unit vector n =

, and the velocity  $c = J = {}^{2}$  have been used. Here = ( $_{1}$ ;  $_{2}$ ) and -Paulim atrices. The elective coupling constant  $512e^{2} = \sim c$  in this case [8] has the order of unity. The full current  $J = 2 {}^{2}$  (j 4A) has a param agnetic (j = i[ $_{1}r_{1}$  cx:+ (1 ! 2)]) and a diam agnetic (4A) parts. Besides, in Eq. (3)  $F_{ik} = 0$   $Q_{k}c_{i}$ ,  $H_{ik} = n$  (A)  $Q_{k}n$ ] =  $0a_{k}$   $Q_{k}a_{i}$ .

Setting in Eq. (3) c = 0 we come back to the model of [11]. The numerical study of the knotted congurations of n - eld in this model has been done in [12, 13, 14]. The lower energy bound in this case

$$F > 32^{2} j_{2} j^{3=4}$$
 (4)

is determ ined [15, 16, 17] by the Hopf invariant,

$$Q = \frac{1}{16^{-2}} d^{3}x "_{ikl}a_{i}\theta_{k}a_{l}:$$
 (5)

2

At compacti cation R<sup>3</sup>! S<sup>3</sup> and n 2 S<sup>2</sup>, the integer Q 2  $_{3}$  (S<sup>2</sup>) = Z shows the degree of linking or knotting of lam entalmanifolds M 2 S<sup>3</sup>, where the vector eld n (x;y;z) is de ned. In particular, for two linked rings (H opf linking) Q = 1, for the trefoil knot Q = 6 and etc. Signi cant point [18] is as follows:  $_{3}$  (C P<sup>M</sup>) = 0 at M > 1 and  $_{3}$  (C P<sup>1</sup>) =  $_{3}$  (S<sup>2</sup>) = Z. In the latter case the order parameter (2) is two-component one [8] and linked or knotted soliton con gurations are labeled by the H opf invariant (5).

Energy bounds. { Let us assume that  $= _0$  can be nd from the minimal value V ( $_0$ ) of the potential V (), but the velocity c does not equal zero. Equation (3) in this case has the following form :

$$F = F_{n} + F_{c} \quad F_{int} = d^{3}x \quad (\theta_{k}n)^{2} + H_{ik}^{2} + \frac{1}{4}c^{2} + F_{ik}^{2} \quad 2F_{ik}H_{ik} : (6)$$

It is seen from Eq. (6) that a superconducting state with jcj 1 has the energy which is less than the minimum in Eq. (4). To nd the lower free energy bound in the superconducting state with  $c \in 0$ , it has been proved [19] that the following inequality for free energy parts takes place

$$F_n^{5=6} F_c^{1=2} > (32^2)^{4=3} j_j$$
; (7)

where

$$L = \frac{1}{16^{-2}} Z^{2} d^{3}x "_{ikl}c_{i}\theta_{k}a_{l}$$
(8)

is the degree of mutual linking [21] of the velocity c lines and of the magnetic eld  $H = [r \ a]$  lines. Quantities like (5), (8) arise as rst integrals in the theory of an ideal or barotropic uid [21]. In general, the integrals in Eqs. (5), (8) could be de ned by the asymptotic linking numbers [22].

It follows from the Schwartz-Cauchy-Bunyakovsky inequality that  $2F_n^{1=2}$   $F_{rat}^{1=2} > F_{int}$ . Setting the boundary value  $F_{int}$  into Eq.(6), we get  $F_{m in} = (F_n^{1=2} - F_c^{1=2})^2$ . The mimimum value 32  $^2Q^{3=4}$  of the function  $F_n$  and Eq. (7) lead [19] to

$$F > 32^{2} p j^{3=4} (1 j_{1} j_{2} j_{2})^{2} :$$
 (9)

The trivial case Q = 0 should be considered after the lim it L = 0. Let us pay also an attention to the self-dual relation  $F_n = F_c$  which follows from  $F_{m \text{ in}}$ .

In general, the Hopf-Chern-Sim onsm atrix,

$$K = \frac{1}{16^{-2}}^{Z} d^{3}x "_{ikl}a_{i} @_{k}a_{l} = \frac{Q L^{0}}{L Q^{0}} : (10)$$

can be de ned in our case by linking numbers Q 2  $\mathbb{Z}$  of spin and spin-charge (fL;Q<sup>0</sup>g  $\neq$   $\mathbb{Z}$  at all) degrees of

freedom. In Eq. (10), K is a sym m etrical matrix ( $L^0 = L$ ) [22] and  $a_i^1$   $a_i; a_i^2$  g. Following the approach of [19] which is based on employ of the Holder inequality chain as well as on the Ladyzhenskaya [20] inequality, one can nd that

$$F_n^{1=2} F_c^{5=6} > (16^{-2})^{4=3} j_c^0 j_i$$
 (11)

The distiction of the coe cient in Eq. (11) from Eq. (7) results from the charge 2e of pairs (due to the coe cient 1=4 in the rt term of the free energy part  $F_c$ ).

It follows from Eq. (9) that for all numbers L < Q the energy of the ground state is less than that in the model described in [11], for which the inequality (4) is valid. The origin of the energy decrease can be easily understood. Even under the conditions of the existence of the param agnetic part jof the current J, the diam agnetic interaction in the superconducting state consumes its own energy of the current and a part of the energy relating to the n-eld dynam ics for all state classes with L 6 Q.

{ In the context of the high-Charge density. tem perature superconductivity problem s [23, 24, 25, 26], Eq. (2) has the sense of the condition of the correlated or linked factorization of charge and spin degrees of freedom (from the gauge invariant order param eter point of view (3)). In fact, the constant value of the charge density plays the role of a tuning parameter of the system. In this case, the order parameter n describes the distribution of spin degrees of freedom, while the order param eter c contains the contribution both of spin degrees of freedom in the param agnetic part of the current and of the U(1) charge degrees of freedom in the diam agnetic part of the current J. Due to such mixing, the gauge invariant order parameters n and c dier principally and topologically. If the mutual linking index is de ned by the current J, the measure in Eq. (8) depends on the charge density. The vector  $c = J = 2^{2}$  is normalized to the charge density and, as distinct from n, belongs to the noncompact manifold. This leads to the fact that the Hopfnum bers for it are not integer valued ((L;Q<sup>0</sup>)  $\neq$  Z), and the linking numbers are included into the factor of Eq. (9) in the form of the ratio L=Q.

Let the parameter  $_0$  change in some range. Since all terms in Eq. (3) are of the same order, the velocity c and therefore the number L decrease as  $_0$  increases. In this case, if L is rather small, the smallest superconducting gap goes down with the growth of Q at the background of large value 32  $^2$ Q  $^{3=4}$  of the spin gap.

As  $_0$  decreases, the following e ect takes place. The radius R of the compactication R<sup>3</sup> ! S<sup>3</sup> being proportional to  $_0^{-1}$ , increases till it exceeds a certain critical value, R<sub>cr</sub>. At R > R<sub>cr</sub> the Hopf map [27] is unstable relative to in nitesimal perturbation of vortex linked eld distribuions. As a result, the U (2) symmetry associated with identical Hopf map is spontaneously broken.

This means that the topological solitons, instead of being spread out over the whole of S<sup>3</sup>, are then localized around a particular point (the base point of the stereographic projection) and collapse into localized structures [17]. This picture corresponds to the existence of an optim alvalue of the relation L=Q. Under such a key condition of the restoration of com pactness of the base manifold, inhom ogeneous superconducting state appears. As the parameter  $_0$  decreases further, the free energy F due to the inhom ogeneity term ( $(e_k)^2$  increases again and suppresses the superconducting gap.

Up to now the vector A characterized the internal charge U (1) gauge symmetry. If we apply an external electrom agnetic eld the vector potential A equals the sum of the internal and the external gauge potentials. As a result, due to the diam agnetism of the superconducting state, the velocity c decreases. Like in the case of increasing  $_0$ , this leads to suppression of the superconducting gap. The answer to the question on the existence of full or partial M eisner screening in these states depends on the result of the com petition of contributions from neutral j and charged 4A parts to the full current J.

The soft case  $\Leftrightarrow$  const both for c = 0 [29] and  $c \Leftrightarrow 0$ arouses certain interest. It is more complicated due to some reasons and will be considered in a separate paper. The point  $F_c=F_n = 1=4$  where the elds of the inequalities (7), (11) intersect, deserve an additional careful study as well. The equality in Eq. (9) under this remarks should be understood as an ideal lim it depending on topologial characteristics of knots and links only.

D iscussion. { In the (3+0)D case of the free energy (3), H opf invariant (5) is analogous to the C hem-Sim ons action (k=4) dtd<sup>2</sup>x " a @ a determ ining strong correlations of (2 + 1)D modes [30, 31] at k ' 2. In planar system s, this coe cient has the sense of braiding of the excitation world lines. In particular, for the sem ion k = 2. K exping in m ind the relation of spatial dim ensionality of the system s in their quantum and statistical descriptions, we note that the (2 + 1)D dynam ical case k = 2 of the open world line ends of excitations is equivalent to the compact (3 + 0)D statistical exam ple of the H opf linking Q = 1 in Eq. (3) (see Fig. 1).

Similarly to the description of the states in the fractional quantum Halle ect with the lling factor = p=q and p;q 2 Z, the Eq. (9) depends on L=Q. The Hopf invariant Q 2 Z enum erating gauge vakua [28], is equivalent to the degree q of the ground state degeneracy and L plays the role of the lling degree p of the incom pressible charged uid state. We have stressed already that the number L is, generally, not integer valued. From this point of view the multiplier (1 L=Q) in Eq. (9) is equivalent to the Hall lling factor 1 for holes. (Let us rem ark that the mentioned parameter  $\frac{2}{0}$  has the sense of the particle density). The distinction of our system from the fractional quantum Halle ect state is that due



FIG.1: Examples of n- and c- eld line con gurations in the form of Borrom ean rings with vanishing pairwise linking numbers K (on the lh.s.) and in the form of the closed chain of correlated charge-spin sem ion loops with Q = 1 (on the rh.s.).

to compressibility of the superconducting state, where the charge gauge U (1) symmetry is broken, the elective number L of the charge degrees of freedom is continuous. It plays the role of an interpolation parameter [32] which connects sectors with L  $\geqq$  Z and L 2 Z in one and the same universality class of P fa an states. Apparently it is true for all P fa an-like states with Q > 1 and numbers L < Q.

In conclusion, we have studied the charge density bounds of the superconducting states using the C P<sup>1</sup> version of G inzburg-Landau m odel under the conditions of the existence of linking and knotting phenom ena of the n-and c-elds being the gauge invariant order parameters of the considered system . We have shown that the superconducting states exist in the nite range  $\frac{2}{1} < \frac{2}{0} < \frac{2}{2}$  of the charge density values. In this range, the eld con g-urations with eld lines characterized by sem ion values of the linking num bers are preferable.

I would like to thank A G. Abanov, E A. Kuznetsov and G E. Volovik for advices, V F. G antmakher for the crucial remark, and L D. Faddeev, G M. Fraim an, A G. Litvak, V A. Verbus for useful discussions. This work was supported in part by the RFBR under the grant No. 01-02-17225.

- [1] H.K.Mo att, Nature 347, 367 (1990).
- [2] V M H.Ruutu, U.Parts, JH.Koivuniem i, M.Knusius, E.V.Thuneberg, and G E.Volovik, JETP Lett. 60, 671 (1994).
- [3] Yu G. Makhlin, T.Sh. M isinpashaev, JETP Lett. 61, 49 (1995).
- [4] M J. M onastyrsky, P.V. Sasorov, Sov. Phys. JETP 66, 683 (1987).
- [5] V. Katritch, J. Bednar, D. M ichoud, R.G. Scharein, J. Dubochet, and A. Stasiak, Nature 384 142 (1996).
- [6] E.W itten, Commun.M ath.Phys.121, 351 (1989).
- [7] I.Aranson, L.Kramer, Rev.Mod.Phys.74, 99 (2002).

- [8] E.Babaev, LD.Faddeev, A.J.Niem i, Phys.Rev.B 65, 100512 (2002); cond-m at/0009438.
- [9] L.D.Faddeev, A.J.Niemi, Phys.Lett.B 525, 195 (2002).
- [10] Y M. Cho, Phys. Rev. Lett. 87, 252001 (2001); hepth/0110076.
- [11] L.D. Faddeev, A.J. Niemi, Nature 387, 58 (1997).
- [12] J. G ladikowski, M. Hellmund, Phys. Rev. D 56, 5194 (1997).
- [13] R A. Battye, P M. Sutcli e, Phys. Rev. Lett. 81, 4798 (1998).
- [14] J.Hietarinta, P.Salo, Phys. Lett. B 451, 60 (1999).
- [15] A F. Vakulenko, L.V. Kapitansky, Sov. Phys. Dokl. 24, 433 (1979).
- [16] A.Kundu, YuP.Rubakov, J.Phys.A 15, 269 (1982).
- [17] R.S.W ard, Nonlinearity 12, 1 (1999), hep-th/9811176.
- [18] A G. Abanov, P W. W iegm ann. G eom etrical phases and quantum numbers of solitons in nonlinear sigm a-m odels, hep-th/0105213.
- [19] A P. Protogenov, V A. Verbus, JETP Lett. (to be published).
- [20] O A. Ladyzhenskaya. The m athem atical theory of viscous incom pressible ow, G ordon and B reach, 1969.
- [21] V E. Zakharov, E A. Kuznetsov, Phys. Usp. 40, 1087

(1997).

- [22] V J. A mold, B A. K hesin. Topological M ethods in Hydrodynam ics. Appl. M ath. Sci. 125, Chapt.3.
- [23] M. Sigrist, D.B. Bailey, and R.B. Laughlin, Phys. Rev. Lett. 74 3249 (1995).
- [24] P A. Lee, N. N agaosa, T K. Ng, and X G. W en, Phys. Rev.B 57, 6003 (1998).
- [25] P A. Lee and X G. W en. Vortex structure in underdoped cuprates, cond-m at/0008419.
- [26] S.Chakravarty, R B.Laughlin, D.Morr, C.Nayak.Hidden order in the cuprates, cond-m at/0005443.
- [27] E A. Kuznetsov, A.V. M ikhailov, Phys. Lett. A 77 37 (1980).
- [28] P. van Baal, A. W ipf. Classical gauge vacua as knots, hep-th/0105141.
- [29] M. Lubcke, S.M. Nasir, A. Niemi, and K. Toroko, Phys. Lett. B 534 195 (2002); hep-th/0106102.
- [30] LA.Abram yan, A P. Protogenov, VA. Verbus, JETP Lett. 69, 887 (1999).
- [31] A P. Protogenov, JETP Lett. 73, 255 (2001).
- [32] N. Read, and Dm itry Green, Phys. Rev. B 61, 10267 (2000).