

Slow light and the phase of a Bose-Einstein condensate

P. Ohberg

School of Physics and Astronomy, University of St Andrews,
North Haugh, St Andrews, Fife, KY16 9SS, Scotland

We investigate the propagation of light with ultra low group velocity in a Bose-Einstein condensate where the phase is not uniform. The light is shown to couple strongly to the phase gradient of the condensate. The interaction between the light and the condensate enables us to perform a phase imprinting where the phase of the condensate is imprinted on the light. We illustrate the effect by showing how one can measure the fluctuating phase in an elongated quasicondensate.

03.75.Fi, 05.30.Jp

Superfluids such as Bose-Einstein condensates of alkali atoms [1] are systems which show a conceptual simplicity yet provide all the intriguing qualities of an interacting many-body system. Many superfluid phenomena are strongly connected to the phase of the fluid. For instance vortices [2] which have been created in a stirring manner similar to the rotating bucket experiment [3], and solitons [4,5], are typical examples of a condensate with a nonuniform phase. Also quantum shock waves causing topological defects in the superfluid, have been created using ultra-compressed pulses of slow light [6]. Here we show how the properties of slow light can be used to detect the phase of the macroscopic wavefunction of the condensate nondestructively, where the slow light receives the phase of the condensate in a way similar to phase imprinting [4,5].

The recent experimental demonstrations with slow light has revealed a whole new playground with interesting applications [6-8]. In this paper we show the advantages with slow light when the phase of the condensate is of interest. Light, and especially the phase contrast imaging method, has been successfully used to probe the density of the condensate in situ [9]. In order to measure the phase, on the other hand, destructive methods have been used where the interference pattern has revealed the phase of the condensate [10]. In this paper we show using slow light that the phase of the condensate can be imprinted on the light in a nondestructive way, making it possible to follow the phase evolution of the condensate in situ.

The envisioned experiment is based on Electromagnetically Induced Transparency (EIT) [11-13]. Suppose that the condensate is trapped in an elongated trap and illuminated by a uniform control beam. The beam controls the group velocity v_g of a second beam, the probe beam. The probe shall be a pulse with a frequency in the laboratory frame that matches exactly the atomic transition frequency between the ground state and one of the excited states coupled by the control beam. The condensate density, which can be monitored in-situ by phase-contrast microscopy [9], is supposed to be station-

ary, but the phase can in certain cases fluctuate [14,15]. For low group velocities the probe beam couples strongly to the condensate phase gradient which results in a phase imprinting on the light. The condensate phase can consequently be read out by analyzing the interference pattern between the initial and final probe light.

Let us start by deriving the equations of motion for the slow light and the Bose-Einstein condensate. The probe light is described by a real scalar field ψ where we ignore the polarization, and ψ represents the electric field in units of the vacuum noise, $E = (\hbar \omega_0)^{1/2} \psi$. Here ω_0 denotes the resonance frequency of EIT. In a medium at rest, slow light is subject to the principle of least action with the Lagrangian density [12]

$$L_L = \frac{\hbar}{2} (1 + \chi) (\partial_t \psi)^2 - c^2 (\nabla \psi)^2 - \frac{1}{2} \omega_0^2 \psi^2 \quad (1)$$

where the group index χ corresponds to a group velocity of

$$v_g = \frac{c}{1 + \chi} \quad (2)$$

and χ is proportional to the density of the condensate, n , and inversely proportional to the intensity of the control beam I_c [16,12],

$$(1 + \chi) \frac{\omega_0^2 \psi^2}{\hbar \omega_0} = \frac{1}{2} \frac{I_p}{I_c} \quad ; \quad (3)$$

Here the intensities of the probe and control fields are calibrated in terms of the Rabi frequencies Ω_p and Ω_c , respectively. The control beam will dominate and in practice Ω_p^2 / Ω_c^2 does not exceed 10^{-1} [7]. This means the less intense the control beam is the slower the light is. The corresponding momentum density of the light which is needed when describing the coupling between the light and the condensate, is obtained from the symmetric energy-momentum tensor [17,12]

$$P = \hbar (\partial_t \psi) \nabla \psi \quad ; \quad (4)$$

Consider for the time being the general situation with a moving condensate. The condensate with the slow u

and the probe light must now be taken as a combined dynamical system with the total Lagrangian density

$$L = L_L + L_M \quad (5)$$

where the condensate is described by the Gross-Pitaevskii Lagrangian density

$$L_M = \int d^3r \left[\frac{\hbar^2}{2} |\nabla \psi|^2 + \frac{\hbar^2}{2m} (\nabla \psi^-)^2 + \frac{g}{2} \psi^\dagger \psi + V \psi^\dagger \psi \right] \quad (6)$$

Here m is the atomic mass, g characterizes the atom-atom collisions, V denotes the external potential and the condensate density. The coupling between light and matter is given by the relation [18]

$$m \dot{u} = \hbar r S + \rho_0 P; \quad \dot{\phi} = 0 \quad (7)$$

where ρ_0 is a constant. This coupling gives the correct equation of motion

$$\partial_t u + r(u) = 0 \quad (8)$$

from the Euler-Lagrange equation.

Using the Lagrangian in Eq. (5) we derive the equation of motion for the light field. The resulting wave equation is

$$(1 + \epsilon) \partial_t^2 u - c^2 r^2 + \epsilon_0^2 + \partial_t u - r(u) = 0 \quad (9)$$

In order to simplify things we assume the light field can be expressed as a propagating pulse with a single frequency ω_0 and a slowly varying amplitude. The resulting equation of motion is then of the Schrodinger type

$$i \partial_t u(r) = \frac{c^2}{2(\epsilon + 1)\epsilon_0} r^2 u(r) + \frac{i}{2(\epsilon + 1)} (2u - r' u + (r - u)' u) \quad (10)$$

where the coupling between light and matter is primarily described by the phase gradient of the condensate.

Let us illustrate the consequences of the coupling between the light and the phase of the condensate by using an elongated quasicondensate where the phase can fluctuate [14,15]. Fluctuations of the density and the phase of a condensate are related to the elementary excitations. The density fluctuations are dominated by the excitations of the order of the chemical potential. If the condensate is very elongated but still of a 3D character, the wavelength of the density excitations are much smaller than the radial size of the condensate. The fluctuations are therefore of an ordinary 3D form and are small. Consequently the total field operator can be written as

$$\hat{\psi}(r) = \int \frac{d^3k}{(2\pi)^3} \hat{a}(k) e^{i\mathbf{k}\cdot\mathbf{r}} \quad (11)$$

where $\rho_0(r)$ is the stationary density and $\hat{\psi}(r)$ is the operator describing the phase of the condensate. This operator is given by [19]

$$\hat{\psi}(r) = \int \frac{d^3k}{(2\pi)^3} \hat{a}(k) e^{i\mathbf{k}\cdot\mathbf{r}} \quad (12)$$

where \hat{a} is the quasiparticle annihilation operator with quantum number \mathbf{k} and energy ϵ_k . The mode functions $f^+ = u + v$ are the sum of the two functions u and v which are the solutions to the Bogoliubov-deGennes equations.

In an elongated condensate the excitations are mainly of two kinds: axial excitations with $\epsilon_k < \hbar\omega_0$ and radial excitations $\epsilon_k > \hbar\omega_0$. The latter has a 3D character since the wavelength is typically less than the radial size of the cloud and consequently the fluctuations are small. The axial excitations on the other hand have wavelengths larger than the radial size of the cloud and have a 1D behavior. Therefore these excitations will be most important for the axial fluctuations of the phase. In order to actually calculate the phase we note that for a harmonic external potential the low energy axial modes [20] are described by the energy $\epsilon_k = \frac{1}{2} \hbar \omega_0 \sqrt{k^2 + 3}$ and the functions

$$f^+ = \frac{S}{4(\epsilon + 1)R^2 L} P_{(1,1)} \left(\frac{z}{L} \right) \quad (13)$$

where $P_{(1,1)}$ are the Jacobi polynomials, R the radial size and L the axial length of the cloud. The quasiparticle annihilation operators are now replaced by complex amplitudes u and v . To reproduce the quantum statistical properties of the phase the amplitudes are sampled as random variables with a zero mean value, $\langle u \rangle = \langle v \rangle = 0$, and bosonic number density

$$\langle |u|^2 \rangle = \frac{1}{e^{\beta \epsilon_k} + 1} \quad (14)$$

where β is the inverse temperature [14].

If the coupling is dominated by the phase gradient, in other words we neglect the nonlinear term for the light in Eq. (10), it is clear that we can use the light as a weak probe which is affected by the condensate phase gradient. The condensate will necessarily be very elongated. We can therefore solve the dynamics in 1D since the low energy axial excitations will acquire a 1D character. From Eq. (10) it is immediately clear that the light will pick up a phase

$$\phi(r; z) = \phi_0(r; z) e^{i\phi(z)} \quad (15)$$

where the phase is given by the integral

$$\phi(z) = \frac{k_0}{v_g} \int_{-z}^z dx u(x) = \frac{\hbar k_0}{m v_g} S(z) \quad (16)$$

with $k_0 = \frac{1}{c}$ and $S(z)$ the phase of the condensate. This scenario requires that the interaction between the probe light and the condensate is turned on suddenly. This can indeed be achieved by tuning the control beam such that the probe beam is rapidly slowed down and allowed to propagate in the condensate. This results in a pulse delay typically of the order of a few micro seconds [7].




FIG. 1. a) The experimental setup with the control beam and the probe beam. The probe beam reveals the phase properties of the condensate. b) The probe beam is split where one part is let to interact with the condensate. Detection of the condensate phase is made with an interference experiment. Note the transversal interference pattern.

To illustrate the phase measuring process we solve Eq. (10) for a light pulse moving with the group velocity 1 m/s during 1.5 ms in a condensate with $N = 3.3 \cdot 10^4$ Na atoms, $\hbar\omega_z = 2 \cdot 10 \text{ Hz}$ and $\hbar\omega_x = 2 \cdot 400 \text{ Hz}$. We choose a low group velocity and a relatively long propagation time in order to show any effects on the light intensity from the condensate phase. In Fig. (2) the condensate together with one possible phase representation is shown at a temperature well below the condensate critical temperature, $T = 0.75 T_{\text{BEC}}$. From Fig. (2a) it is clear that only the low excitations are significant, as expected. The phase fluctuations are clearly present and are typically of the order of ≈ 3 in the center of the con-

densate. The shape of the light pulse, in Fig. (2b), is only slightly altered due to the acquired condensate phase. The pulse shape changes are typically small compared to the overall pulse width and are therefore not necessarily very well suited for detecting any acquired phase from the condensate. In Fig. (2c) we show the resulting intensity when the light which has acquired the phase of the condensate, and the initial probe beam interferes. Note here that the interference pattern will be in the transverse direction compared to the pulse propagation. For sufficiently low group velocities the acquired condensate phase is amplified by the factor $\hbar k_0 = m v_g$, which makes it possible to study also small phase fluctuations not detectable by a time of flight measurement [14]. As in any measurement process, there is inevitably going to be a back-action on the condensate from the light. This back-action can, however, be made arbitrarily small by choosing an appropriate ratio j_p/j_c . The group velocity, which is a function of the control beam, is not affected by this ratio.

The 1D character of the condensate-light phase imprinting has here been used for simplicity. Similar phase fluctuations in 2D condensates can also exist [21] and consequently the same situation emerges concerning the properties of the light. One important factor, however, is the inherent time dependence of the phase fluctuations in the condensate. As was already seen in Fig. (2) the phase fluctuations are dominated by the lowest modes. This means also that the dynamics is governed by the corresponding frequencies $\omega = \hbar^{-1} \epsilon$. For a typical external harmonic potential with $\hbar\omega_z = 2 \cdot 10 \text{ Hz}$ we obtain a timescale for the phase fluctuations of the order of 20 ms , corresponding to the ten lowest modes, which is long enough to treat the phase as stationary in the imprinting process. The typical timescale to apply and slow down the light pulse is of the order of micro seconds [7] which allows us to assume an instant phase imprinting.

In summary we have shown that light with extremely low group velocity can be used to probe the phase of a condensate. The method was illustrated by imprinting the fluctuating phase of a quasicondensate onto the slow light propagating through the condensate. The phase imprinting technique also allows for studying other forms of phase gradients such as vortices [22] and solitons [23]. Especially solitons in two-component condensates with repulsive interaction could be detected without opening the trap [4]. Two-component solitons are significantly more difficult to observe because the total density will always stay constant, compared to the single component condensate soliton which shows a density notch at the position of the soliton. The probe light in this case would simply pick up the phase of the soliton solution. If the light propagates during a long time in a medium with a nonuniform phase it is also possible to study scattering of the light from the condensate phase [18] which in principle can also be used to measure the phase gradi-

ent, although this method will not necessarily reveal the condensate phase as clearly as a direct interference experiment.

This work was supported by EPSRC. The author wishes to thank T. Kiss and U. Leonhardt for fruitful discussions. The Department of Physics at Abo Akademi University is acknowledged for their hospitality.




FIG. 2. (a) One possible phase at the temperature $T = 0.75T_{\text{BEC}}$. Only the lowest modes are important in the phase fluctuations. (b) The shape of the light pulse (solid black line) is only slightly altered due to the acquired phase. The grey curve shows the initial pulse shape (arb. units). The dashed line shows the density of the condensate (arb. units). (c) The resulting intensity of the light when interfering with the initial probe light (arb. units).

- 2498 (1999); K. W. Madison, F. Chevy, W. Wohlleben, and J. Dalibard, *ibid.* 84, 806 (2000); J. R. Abo-Shaeer, C. Raman, J. M. Vogels, and W. Ketterle, *Science* 292, 476 (2001).
- [3] D. R. Tilley and J. Tilley, *Superfluidity and Superconductivity*, (Adam Hilger, Bristol, 1990).
- [4] S. Burger, K. Bongs, S. Dettmer, W. Ertmer, K. Sengstock, A. Sanpera, G. V. Shlyapnikov, and M. Lewenstein, *Phys. Rev. Lett.* 83, 5198 (1999).
- [5] J. Denschlag, J. E. Sarsian, D. L. Feder, Charles W. Clark, L. A. Collins, J. Cubizolles, L. Deng, E. W. Hagley, K. Helmerson, W. P. Reinhardt, S. L. Rolston, B. I. Schneider, and W. D. Phillips, *Science* 83, 97 (2000).
- [6] Z. Dutton, M. Budde, Ch. Slowe, and L. V. Hau, *Science* 293, 663 (2001).
- [7] L. V. Hau, S. E. Harris, Z. Dutton, and C. H. Behroozi, *Nature* 397, 594 (1999); Ch. Liu, Z. Dutton, C. H. Behroozi, and L. V. Hau, *ibid.* 409, 490 (2001).
- [8] U. Leonhardt and P. P. Winkler, *Phys. Rev. Lett.* 84, 822 (2000).
- [9] M. R. Andrews, M. O. Mewes, N. J. van Duten, D. S. Durfee, D. M. Kum, and W. Ketterle, *Science* 273, 84 (1996).
- [10] S. Inouye, S. Gupta, T. Rosenband, A. P. Chikkatur, A. Gorlitz, T. L. Gustavson, A. E. Leanhardt, D. E. Pritchard, and W. Ketterle, *Phys. Rev. Lett.* 87, 080402 (2001).
- [11] M. O. Scully and M. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).
- [12] U. Leonhardt, *arXiv:gr-qc/0108085*.
- [13] I. Canusotto, M. Artoni and G. C. La Rocca, *JETP Letters* 72, 289 (2000).
- [14] S. Dettmer, D. Hellweg, P. Ryyty, J. J. Arlt, W. Ertmer, K. Sengstock, D. S. Petrov, G. V. Shlyapnikov, H. Kreutzmann, L. Santos, and M. Lewenstein, *Phys. Rev. Lett.* 87, 160406 (2001).
- [15] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, *Phys. Rev. Lett.* 87, 050404 (2001).
- [16] M. Fleischhauer and M. D. Lukin, *Phys. Rev. Lett.* 84, 5094 (2000).
- [17] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Pergamon, Oxford, 1975).
- [18] U. Leonhardt and P. Ohberg, *arXiv:cond-mat/0110514*.
- [19] D. V. Fil and S. I. Shevchenko, *Phys. Rev. A*, 64 (2001) 013607.
- [20] S. Stringari, *Phys. Rev. A* 58, 2385 (1998).
- [21] D. S. Petrov, M. Holzmann, and G. V. Shlyapnikov, *Phys. Rev. Lett.* 84, 2551 (2001).
- [22] M. Artoni and I. Canusotto, *arXiv:cond-mat/0206104*.
- [23] P. Ohberg and L. Santos, *Phys. Rev. Lett.* 86, 2918 (2001).

[1] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* 71, 463 (1999).

[2] M. R. Matthews, B. P. Anderson, P. C. Haljan, D. S. Hall, C. E. Wieman, and E. A. Cornell, *Phys. Rev. Lett.* 83,