

Substrate effects in magneto-optical second-harmonic generation from first principles: Fe/Cu(001)

Torsten Andersen^{1,*} and W. Hübner^{1,2,†}

¹*Max-Planck-Institut für Mikrostrukturphysik, Weinberg 2, D-06120 Halle/Saale, Germany*

²*Institut für Theoretische Physik, Karl-Franzens-Universität Graz, Universitätsplatz 5, A-8010 Graz, Austria*

We compute the nonlinear optical response of an Fe monolayer placed on top of 1 to 4 monolayers of Cu(001). Our calculation is based on *ab initio* eigenstates of the slab, which are obtained within the full-potential linearized augmented plane-wave method. The ground-state spin-polarized electronic structure is converged self-consistently to an accuracy better than 0.1 mRy. Subsequently, we take the spin-orbit interaction into account within a second variational treatment. The new set of eigenstates allows us to calculate the magneto-optical transition matrix elements. The second-harmonic response is determined in the reflection geometry with magnetization perpendicular to the surface (the so-called polar configuration) using the surface-sheet model. Adding layers of a noble metal (Cu) to the Fe monolayer gives a new degree of freedom for the inclusion of nonmagnetic Cu d bands to the nonlinear magneto-optical response of the slab, and the energy bands show that such an addition converges essentially to an addition of d-states and a small broadening of the d-band with growing number of Cu layers. The screened nonlinear optical susceptibility is calculated and converges quite well with a growing number of Cu layers. Our first-principles results confirm that the magnetic tensor elements of the nonlinear optical response tensor are roughly of the same order of magnitude as the nonmagnetic ones (in contrast to linear optics, where the magnetic response is only a minor correction).

PACS numbers: 78.20.Ls, 73.20.At, 78.66.Bz

* Electronic address: thor@mpi-halle.mpg.de

† Electronic address: huebner@mpi-halle.mpg.de