

On the ground state properties of antiferromagnetic half-integer spin chains with long-range interactions

Tigran Hakobyan*

Theory Division, Yerevan Physics Institute, 375036, Alikhanyan 2, Yerevan, Armenia

(Dated: March 22, 2022)

The Lieb-Shultz-Mattis theorem is extended to Heisenberg chains with long-range interactions. We prove that the half-integer spin chain has no gap, if it possesses *unique* ground state and the exchange decays faster than the *inverse-square* of distance between spins. The results can be extended to a wide class of one-dimensional models.

PACS numbers: 75.10.Jm 75.50.Ee 71.10.-w 68.35.Rh 05.50.+q 02.30.Ik

The zero-temperature ground state properties of one-dimensional antiferromagnets have been studied intensively over a long period. There are various numerical and field theoretical approaches to the problem. At the same time there are just few exact non-perturbative methods. The spin $S=1/2$ antiferromagnetic (AF) Heisenberg spin chain with nearest-neighbor (n.n.) interactions was solved exactly by Bethe [1]. Bethe's ansatz was applied later to peculiar higher spin chains with polynomial spin exchange [2].

Lieb, Schultz and Mattis [3] offered another non-perturbative approach. They proved that $S=1/2$ AF n.n. Heisenberg chain has *gapless* excitations above the *unique* ground state. Later Haldane [4], using σ -model description, argued, that the AF n.n. *integer* spin Heisenberg chains have a *gap* between the ground state and the first excited state, whereas the *half-integer* spin chains have *no gap*. Haldane's conjecture was proven exactly for the *half-integer* spin chains [5, 6] by means of extension of the Lieb-Schultz-Mattis (LSM) theorem. The LSM theorem was extended also to the case with an applied magnetic field [7]. It was generalized to Heisenberg chains with $SU(n)$ [6] and other [8] symmetries, to fermionic chains [9], to spin [10] and fermionic [11] models on ladders, etc. Very recently, its extension to higher dimensions has been discussed [12, 13, 14].

This letter is devoted to the extension of LSM theorem to AF Heisenberg chains with *long-range* interactions. The $S = 1/2$ chain with the spin exchange decaying as $1/r^2$, where r is the distance between spins, has been extensively investigated since it was solved exactly by Haldane and Shastri [17]. The integrability is a consequence of an infinite symmetry of this model [18]. Despite of presence of long-range interactions, its low-energy properties are similar to those of the chain with n.n. interactions: both models are gapless and belong to the same universality class. Here we discuss the zero-temperature ground state properties of the *translationally* invariant AF *half-integer* spin chain with *arbitrary* exchange coupling. We will prove that, in the thermodynamic limit, it either has *gapless* excitations or *degenerate* ground states, provided that the coupling decays faster than $1/r^2$.

The Hamiltonian of one-dimensional Heisenberg chain with long-range interactions is

$$\mathcal{H}_L = \sum_{r=1}^{L'} \sum_{i=0}^{L-1} J(r) \vec{S}_i \cdot \vec{S}_{i+r}, \quad L' = (L-1)/2, \quad (1)$$

where the chain length L is chosen to be *even*. Note, that the spin exchange coupling $J(r)$ depends on the distance between the spins only. Thus, \mathcal{H}_L is translationally invariant. The periodic boundary conditions are assumed and the sum over r is taken up to the half-size L' of the chain.

Let us consider the nonlocal unitary operator

$$U = e^{\mathcal{A}}, \quad \mathcal{A} = \frac{2\pi i}{L} \sum_{k=1}^L k S_k^z. \quad (2)$$

It rotates the spins around the z -axis with the relative angle $2\pi/L$ between the neighboring spins. We are interested in formal algebraic properties of U . However, as it was shown recently, it has a clear physical meaning too. In spin chains and ladders, the ground state expectation value of U can be treated as an order parameter, which characterizes various valence-bond-solid ground states [15]. In sine-Gordon theory, it is related to expectation values of vertex operators [16].

It is well known that the action of (2) on the ground state of the Heisenberg chains with *short-range* interactions gives rise to the state with an energy, which approaches the energy of the ground state in the thermodynamic limit $L \rightarrow \infty$ [3, 6]. Below we will discuss the possibility to extend this property to the Hamiltonian (1) with *long-range* interactions.

In this letter, for the sake of simplicity, we assume the *uniqueness* of the ground state $|\Omega\rangle$ of finite chain. Note, that in this case $|\Omega\rangle$ is a *spin-singlet*. Our assumption is fulfilled for wide class of models on even length chains. In case of alternating couplings, i.e. $J(2r-1) > 0$, $J(2r) \leq 0$, it can be proven *exactly* using Perron-Frobenius type arguments [6].

It is convenient to rewrite \mathcal{H}_L in the following form:

$$\mathcal{H}_L = \sum_{r=1}^{L'} J(r) \mathcal{H}_L^r, \quad \text{where} \quad \mathcal{H}_L^r = \sum_{i=0}^{L-1} \vec{S}_i \cdot \vec{S}_{i+r}.$$

Then the energy gap between the ground state $|\Omega\rangle$ and the shifted state $U|\Omega\rangle$ is

$$\begin{aligned}\Delta E_L &= \sum_{r=1}^{L'} \Delta E_L^r, \quad \text{where} \\ \Delta E_L^r &= \langle \Omega | U^+ \mathcal{H}_L^r U | \Omega \rangle - \langle \Omega | \mathcal{H}_L^r | \Omega \rangle.\end{aligned}\quad (3)$$

The straightforward calculations show that

$$\begin{aligned}U^+ \mathcal{H}_L^r U - \mathcal{H}_L^r &= \sum_{i=0}^{L-1} \{ \mathcal{K}(r/L) S_i^+ S_{i+r}^- + \text{h. c.} \} \\ &+ [\mathcal{H}_L^r, \mathcal{A}], \quad \text{where} \\ \mathcal{K}(\phi) &= \frac{1}{2} \{ \exp(2\pi i \phi) - (1 + 2\pi i \phi) \}\end{aligned}\quad (4)$$

and $S_i^\pm = S_i^x \pm i S_i^y$. Note, that the coefficients in front of spin operators don't depend on the absolute positions of the spins, but depend on their distance. Taking the sum of (4) over r , we obtain:

$$\begin{aligned}U^+ \mathcal{H}_L U - \mathcal{H}_L &= \sum_{r=1}^{L'} J(r) \{ \mathcal{K}(r/L) \mathcal{H}_L^r + \text{h. c.} \} \\ &+ [\mathcal{H}_L, \mathcal{A}], \quad \text{where} \quad \mathcal{H}_L^r \pm = \sum_{i=0}^{L-1} S_i^\pm S_{i+r}^\mp.\end{aligned}\quad (5)$$

The mean value of the commutator $[\mathcal{H}_L, \mathcal{A}]$ vanishes on $|\Omega\rangle$, because $|\Omega\rangle$ is an eigenstate of \mathcal{H}_L . The mean value of \mathcal{H}_L^\pm is of the order of L :

$$|\langle \Omega | \mathcal{H}_L^\pm | \Omega \rangle| \leq 1/4L. \quad (6)$$

Now, using (3), (5) and (6), we can estimate the upper bound of the gap:

$$\begin{aligned}|\Delta E_L| &\leq \frac{L}{2} \sum_{r=1}^{L'} J(r) |\mathcal{K}(r/L)| \\ &= \frac{1}{L} \sum_{r=1}^{L'} r^2 J(r) \frac{|\mathcal{K}(r/L)|}{2(r/L)^2}.\end{aligned}\quad (7)$$

The function $|\mathcal{K}(\phi)|/(2\phi^2)$ is continuous on the unit interval $0 \leq \phi \leq 1$ (see the definition in (4)). So, it can be replaced in (7) by its maximal value C and we obtain:

$$|\Delta E_L| \leq \frac{C}{L} \sum_{r=1}^{L'} r^2 J(r). \quad (8)$$

We conclude, that the energy gap between the ground state and the shifted state $U|\Omega\rangle$ vanishes in the thermodynamic limit, provided that

$$\lim_{L \rightarrow \infty} \frac{1}{L} \sum_{r=1}^{L'} r^2 J(r) = 0.$$

Any function, obeying

$$\lim_{L \rightarrow \infty} r^2 J(r) = 0, \quad (9)$$

satisfies this condition. Note, that up to now we haven't distinguished between the integer and the half-integer spin chains.

In order to prove that $U|\Omega\rangle$ is a real excitation, i.e. it doesn't coincide with the ground state $|\Omega\rangle$ and doesn't approach it in the thermodynamic limit, it is sufficient to show that both states are orthogonal [3, 6]. We use the translation T by one lattice site to demonstrate this. The ground state is an eigenstate of T , because T commutes with the Hamiltonian (1): $T|\Omega\rangle = e^{ip}|\Omega\rangle$, where $p = 0, \pi$ due to the reflection symmetry. The shift operator (2) transforms as: $T^{-1}UT = U \exp(2\pi i S_z^z) \exp(-2\pi i S^z/L)$, where $S^z = \sum_{i=0}^{L-1} S_i^z$ is the z -component of the total spin \vec{S} . We have $S^z|\Omega\rangle = 0$, because $|\Omega\rangle$ is a singlet. Using the equations above, we obtain: $TU|\Omega\rangle = e^{-2\pi i S} e^{ip} U|\Omega\rangle$. We see, that the eigenvalues of T on $|\Omega\rangle$ and $U|\Omega\rangle$ differ for *half-integer* S only. So, in this case the two states are orthogonal. Then, in the $L \rightarrow \infty$ limit, either i) $U|\Omega\rangle$ generates another ground state(s) or ii) $|\Omega\rangle$ remains unique ground state with gapless excitations, created by $U|\Omega\rangle$. In the second case, of course, the ground state should be a *spin-singlet*.

We come to the conclusion that:

For the translationally invariant AF Heisenberg half-integer spin chain with long-range interactions the low-energy state, approaching the ground state in the thermodynamic limit, exists, if the exchange coupling of interacting spins decays faster than the inverse-square of their distance. If, in addition, the ground state remains unique, gapless excitations appear above it.

This statement excludes the possibility to have a gap and unique ground state simultaneously under the aforementioned conditions. This fact remains true even if we omit the ground state uniqueness requirement for finite L . The examples of $J(r)$, which satisfy the conditions above, are:

- a) $1/(r^2 \log(r))$; b) $1/r^\alpha$, $\alpha > 2$;
- c) $r^n \exp(-r/a)$, $a > 0$.

The last case corresponds to an effective short-range interaction.

The statement above gives only the *sufficient* condition for the chain to be gapless or to have degenerate ground states in the thermodynamic limit. The integer spin chains or chains with the exchange coupling decay slower than $1/r^2$ can have gap and unique ground state simultaneously. They can also be gapless. For example, the $S = 1$ chain with the alternating exchange $J(r) = (-1)^r/r^\alpha$, $1 < \alpha < 3$ exhibits gapless behavior [20]. However, the same chain with $J(r) = 1/r^2$ has a gap, as it was shown numerically in [19]. Its ground state

properties are similar to those of Haldane chain with n.n. interactions [4].

As we have mentioned before, the $S = 1/2$ chain with $J(r) = 1/r^2$ corresponds to the Haldane-Shastry integrable model. In case of the *periodic* boundary conditions the coupling is slightly modified [21]: $J_L^{\text{HSh}}(r) = ((L/\pi) \sin(\pi r/L))^{-2}$. For $J(r) = J_L^{\text{HSh}}(r)$ the upper bound of the gap in (7) is nonzero. This fact doesn't guarantee the existence of a low-energy state [22]. Nevertheless, the exact solution shows that this model has gapless excitations.

Very recently, the LSM theorem is considered in higher dimensions [12, 13, 14], where the gap estimation method, used in this letter, can't be applied. Nevertheless, the reasonable arguments, used by Laughlin [23] in quantum Hall effect, can be presented in support of LSM theorem in this case too. These arguments are based on the assumption that during the certain adiabatic process the gap remains robust. This process consists in the insertion and the subsequent increment of an fictitious magnetic flux, which penetrates the ring of the fermionic chain. In the spin language, the magnetic flux is equivalent to a twist of the boundary conditions. It would be interesting to clarify, if the same argument can be applied in the presence of long-range interactions. This would be a way to extend LSM theorem to the chains with the exchange couplings, which decay slower than $1/r^2$.

The results in this letter can be generalized to a wide class of one-dimensional chains with long-range interactions, including fermionic chains, spin chains with higher symmetries, with an applied external field, etc.

The author is grateful to Prof. R. Flume for hospitality during his visit at Physikalisches Institut, Universität Bonn. He thanks R. Flume, R. Poghosian and D. Karakhanyan for stimulating discussions.

This work is supported by Volkswagen Foundation of Germany, grants INTAS #00-561, INTAS #99-01459 and Swiss SCOPE grant.

- [3] E. H. Lieb, T. D. Schultz, and D. C. Mattis, Ann. Phys. (N.Y.) **16**, 407 (1961).
- [4] F.D.M. Haldane, Phys. Lett. A **93**, 464 (1983); Phys. Rev. Lett. **50**, 1153 (1983).
- [5] M. Kolb, Phys. Rev. B **31**, 7494 (1985).
- [6] I. Affleck and E. Lieb, Lett. Math. Phys. **12**, 57 (1986); I. Affleck, Phys. Rev. B **37**, 5186 (1988).
- [7] M. Oshikawa, M. Yamanaka and I. Affleck, Phys. Rev. Lett. **78**, 1984 (1997).
- [8] Y.Q. Li, Phys. Rev. Lett. **87**, 127208 (2001).
- [9] M. Yamanaka, M. Oshikawa and I. Affleck, Phys. Rev. Lett. **79**, 1110 (1997).
- [10] T. Fukui, M. Sigrist and N. Kawakami, Phys. Rev. B **56**, 2530 (1997).
- [11] P. Gagliardini, S. Haas and T.M. Rice, Phys. Rev. B **58**, 9603 (1998).
- [12] M. Oshikawa, Phys. Rev. Lett. **84**, 1535 (2000).
- [13] M. Oshikawa, Phys. Rev. Lett. **84**, 3370 (2000).
- [14] G. Misguich *et al.*, Eur. Phys. J. B **26**, 167 (2002); G. Misguich and C. Lhuillier, cond-mat/0002170 (2000).
- [15] M. Nakamura and S. Todo, cond-mat/0112377 (2001); cond-mat/0201204, in Proceedings for Yukawa-Institute-for-Theoretical-Physics International Workshop "Order, Disorder, and Dynamics in Quantum Spin Systems" (November, 2001); see also N.E. Bonesteel, Phys. Rev. B **40**, 8954 (1989).
- [16] M. Nakamura and J. Voit, Phys. Rev. B **65**, 153110 (2002).
- [17] F.D.M. Haldane, Phys. Rev. Lett. **60**, 635 (1988); B.S. Shastri, Phys. Rev. Lett. **60**, 639 (1988).
- [18] F.D.M. Haldane *et al.*, Phys. Rev. Lett. **69**, 2021 (1992); D. Bernard *et al.*, J. Phys. A: Math. Gen. **26**, 5219 (1993).
- [19] P.D. Sacramento and V.R. Vieira, Z. Phys. B: Cond. Mat. **101**, 441 (1996); Z. Phys. B: Cond. Mat. **103**, 231 (1997).
- [20] P.D. Sacramento and V.R. Vieira, J. Phys.: Cond. Mat. **9**, 10687 (1997).
- [21] Of course, we have: $\lim_{L \rightarrow \infty} J_L^{\text{HSh}}(r) = 1/r^2$. Note, that the dependence on the chain length doesn't change anything in the final result (7), where we can impose $J(r) = J_L(r)$.
- [22] For $J(r) = J_L^{\text{HSh}}(r)$ the right hand side of (7) is equal to the nonzero finite value of the integral

$$\frac{\pi^2}{2} \int_0^{1/2} \frac{|\mathcal{K}(\phi)|}{\sin^2(\pi\phi)} d\phi.$$

* Electronic address: hakob@lx2.yerphi.am

- [1] H. Bethe Z. Phys., **71**, 205 (1931).
- [2] L. Takhtajan, Phys. Lett. A **87**, 479 (1982); H. Babujan, Nucl. Phys. B **215**, 317 (1983).

- [23] R.B. Laughlin, Phys. Rev. B **23**, 5632 (1981).