

# Reply to the comment by E. Menendez-Proupin on "Multiphonon Raman scattering in semiconductor nanocrystals: Importance of non-adiabatic transitions"

E. P. Pokatilov

Department of Theoretical Physics, State University of Moldova,  
str. A. Mateevici 60, MD-2009 Kishinev, Republic of Moldova

S. N. Klimin, V. M. Fomin and J. T. Devreese

Theoretische Fysica van de Vaste Stof, Universiteit Antwerpen (U.I.A.), B-2610 Antwerpen, Belgium

F. W. Wise

Department of Applied Physics, Cornell University, Ithaca, New York 14853, USA

(Dated: May 23, 2002)

In our recent paper [Phys. Rev. B 65, 075316 (2002)], the selection rules for the Raman scattering in spherical nanocrystals are analyzed. Here we show, that a physically expedient choice of the coordinate system, related to the polarization vector of the incoming or scattered light, simplifies and clarifies the analysis.

In the paper<sup>1</sup>, multiphonon Raman scattering in semiconductor nanocrystals is investigated. In that work, we have considered two groups of selection rules: (i) for active exciton states and (ii) for participating phonons. Within the dipole approximation, the physically expedient choice implies that the axis  $Oz$  is parallel to one of the polarization vectors, either  $e^I$  of the linearly polarized incoming light or  $e^S$  of the linearly polarized scattered light. The selection rules on p. 5 of the paper<sup>1</sup> correspond just to this choice of the axis  $Oz$ . We keep one and the same coordinate system when describing both exciton and phonon states throughout the paper<sup>1</sup>.

The selection rules for participating phonons can be expressed in terms of the vector addition of angular momenta. Let us denote  $L_{I(S)} = 1$  the angular momentum of the incoming (scattered) light, and  $l$  the angular momentum of a phonon. The corresponding  $z$ -projections (for an arbitrary coordinate system) are  $M_{I(S)}$  and  $m$ ; respectively. The rules for the angular momentum addition are

$$\begin{aligned} J &= 1 \pm 1; \\ M_I &= M_S + m; \end{aligned} \quad (1)$$

Herefrom, we obtain the selection rules for the phonon angular momentum and for its  $z$ -projection

$$\begin{aligned} l &= 0, 2; \\ m &= M_I - M_S; \end{aligned} \quad (2)$$

Eq. (2) means, in particular, that only  $s$ -,  $p$ - and  $d$ -phonons may participate in the one-phonon Raman scattering. The selection rule (3) is just the conservation law for the  $z$ -projection of the angular momentum.

Let us consider the case when  $Oz \parallel e^I$ ; so that  $M_I = 0$ : For the Raman scattering in the parallel polarizations  $e^S \parallel Oz$ ;  $M_S = 0$ ; and, according to Eq. (3), only the scattering with the participation of phonons with  $m = 0$  is allowed. Otherwise, when one considers the same Raman scattering in the parallel polarizations choosing  $Ox \parallel e^I \parallel e^S$ ; the projections of the angular momenta for both incoming and scattered photons on the axis  $Oz$  do not take definite values, because photon states in this case are superpositions of the states with  $M_{I(S)} = 1$  and  $M_{I(S)} = -1$ : For this coordinate system, following Eq. (3), phonons with  $m = 0; \pm 2$  can participate in the one-phonon Raman scattering. This result is in agreement with the statement of Ref.<sup>2</sup>, that in parallel polarizations, phonons with  $m \neq 0$  can participate in the Raman scattering when the axis  $Oz$  does not coincide with the light polarization vectors. However, in this case, the choice of the axis  $Oz$  is not physically expedient.

For the scattering in the crossed polarizations, let us consider one of the aforesaid physically expedient coordinate systems with  $Oz \parallel e^I; Ox \parallel e^S$ . In this system,  $M_I = 0$ ; while the scattered light is a superposition of states with  $M_S = \pm 1$ : Consequently, the selection rule (3) takes the form  $m = \pm 1$ : It is worth noting, that for any coordinate system with  $Oz \parallel e^I$  or  $Oz \parallel e^S$ , we obtain the same selection rule  $m = \pm 1$  for the crossed polarizations.

For the other coordinate system  $Ox \parallel e^I; Oy \parallel e^S$ , described in the comment<sup>2</sup>, the photon angular momentum projections are:  $M_I = \pm 1; M_S = \pm 1$ . As a result,  $m$  can take the values  $m = 0; \pm 2$ : It is worth mentioning, that when one chooses  $Ox \parallel e^I \parallel e^S$  for the parallel polarizations and  $Ox \parallel e^I; Oy \parallel e^S$  for the crossed polarizations, the selection rule (3) leads to the same allowed values of  $m$  for the parallel and crossed polarizations, what makes a distinction between these two scattering configurations less clear.

In summary, in different coordinate systems the selection rule (3) for the participating phonons (as well as the selection rules for the active exciton states) can formally distinguish just because a rotation of the axis  $Oz$  can lead

to superpositions of states with different values of  $m$  in a new coordinate system. As shown above, the choice of the coordinate system in the paper<sup>1</sup> is physically opportune. When a coordinate system is chosen in another way, the content of scattering can be masked, though it describes the same physical processes.

---

\* Permanent address: Department of Theoretical Physics, State University of Moldova, str. A. M. Mateevici 60, MD-2009 Kishinev, Republic of Moldova

\*\* Also at: Universiteit Antwerpen (RUCA), Groenenborgerlaan 171, B-2020 Antwerpen, Belgium and Technische Universiteit Eindhoven, P.B. 513, 5600 MB Eindhoven, The Netherlands

<sup>1</sup> E. P. Pokatilov, S. N. Klimin, V. M. Fomin, J. T. Devreese, and F. W. Wise, Phys. Rev. B 65, 075316 (2002).

<sup>2</sup> E. Menendez-Proupin, cond-mat/0204502 (unpublished).