

# INTERACTING LINEAR POLYMERS ON THREE-DIMENSIONAL SIERPINSKI FRACTALS

Jelena Maricic<sup>1</sup> and Suncica Elezovic-Hadzic<sup>2</sup>

<sup>1</sup> Physics Department, University of Hawaii at Manoa

<sup>2</sup> Faculty of Physics, University of Belgrade

## Abstract

Using self-avoiding walk model on three-dimensional Sierpinski fractals (3d SF) we have studied critical properties of self-interacting linear polymers in porous environment, via exact real-space renormalization group (RG) method. We have found that RG equations for 3d SF with base  $b = 4$  are much more complicated than for the previously studied  $b = 2$  and  $b = 3$  3d SFs. Numerical analysis of these equations shows that for all considered cases there are three fixed points, corresponding to the high-temperature extended polymer state, collapse transition, and the low-temperature state, which is compact or semi-compact, depending on the value of the fractal base  $b$ . We discuss the reasons for such different low-temperature behavior, as well as the possibility of establishing the RG equations beyond  $b = 4$ .

## Introduction

At low temperatures and in poor solvents linear polymers are in compact globule state, while at high temperatures and in good solvents they have extended coil conformation [1]. Self-interacting self-avoiding walk (SISAW) on a lattice is a good model for linear polymer in solvent. If  $\epsilon < 0$  is the attractive energy per pair of nearest-neighbors monomers (not directly connected) and  $w = \exp(\epsilon/k_B T)$  the corresponding Boltzmann factor, then for  $N$ -step SISAW and  $N \gg 1$  the following behavior of the mean-squared end-to-end distance  $\langle R_N^2 \rangle$  is expected:

$$\langle R_N^2 \rangle \sim N^2 : \quad (1)$$

There is a critical value of interaction  $w = w_c$ , such that for any  $w < w_c$  critical exponent  $\nu = \nu_{SAW}$ , which corresponds to extended coil phase, whereas for any  $w > w_c$  exponent  $\nu$  has smaller value, equal to  $1/d$  for  $d$ -dimensional homogeneous lattices. The value  $w = w_c$  corresponds to the

collapse transition, with  $\beta_c = \beta_c$ , between the low  $\beta$  and high  $\beta$  temperature values:  $1 = d < \beta_c < \beta_{SAW}$ .

On nonhomogeneous lattices collapse transition also occurs, with the different values of the exponent  $\beta$ . Three-dimensional Sierpinski fractals (3d SFs) are nonhomogeneous lattices where exact real space renormalization group (RG) method can be applied for studying asymptotic properties of SISAW [2,3,4].

Renormalization group scheme

Each member of the 3d SF family is labelled by  $b = 2; 3; 4;$  and can be constructed recursively starting with a tetrahedron of base  $b$ , containing  $b(b+1)(b+2)/6$  unit tetrahedrons (so called generator  $G_1(b)$ ) (see fig.1).




Figure 1: First two stages of the construction of 3d SF with  $b = 4$

The subsequent fractal stages are constructed self-similarly, by replacing each unit tetrahedron of the initial generator by a new generator. To obtain the  $l$ th stage fractal lattice  $G_l(b)$ , which we shall call the  $l$ th order generator, this process of construction has to be repeated  $l-1$  times, and the complete fractal is obtained in the limit  $l \rightarrow \infty$ . Fractal dimension  $d_f$  of 3d SF is equal to

$$d_f = \ln[(b+2)(b+1)b/6] / \ln b : \quad (2)$$

In this case RG parameters are restricted generating functions, which are for the  $G_l(b)$  equal to

$$A^{(l)}(x;w) = \sum_{N,P} A_{N,P} x^N w^P ; \quad B^{(l)}(x;w) = \sum_{N,P} B_{N,P} x^N w^P ; \quad (3)$$

where  $x$  is weight of any step of a SISAW (fugacity), and  $A_{N,P}$  ( $B_{N,P}$ ) is the number of  $N$  step SISAW with  $P$  nearest neighbor contacts, traversing the  $G_l(b)$  once (twice) (see fig.2). For elementary tetrahedrons  $G_0(b)$  one has

$$A^{(0)}(x;w) = x + 2x^2w + 2x^3w^3 ; \quad B^{(0)}(x;w) = x^2w^4 : \quad (4)$$

For any b RG equations have the form

$$\begin{aligned} A^{(l+1)} &= \sum_{n,m} A_{n,m} A^{(l)n} B^{(l)m}; \\ B^{(l+1)} &= \sum_{n,m} B_{n,m} A^{(l)n} B^{(l)m}; \end{aligned} \quad (5)$$

where  $A_{n,m}$  ( $B_{n,m}$ ) is the number of SISAW configurations consisting of  $n$  branches of "A" type and  $m$  branches of "B" type, traversing  $G_{l+1}$  once (twice).




Figure 2: Examples of A and B type of SISAW on  $G_1(2)$

Analysis of the  $b = 4$  case

As  $b$  increases the number of possible SISAW configurations quickly grows, which is the reason why we don't quote the corresponding extremely cumbersome RG equations [4] for the studied  $b = 4$  case here. Numerical analysis of these equations shows that there are three physically relevant fixed points.

(1) The high temperature fixed point ( $A = 0.289912; B = 0.0122393; \dots$ ) is reached for any  $w < w_c = 5.028; \dots$  and corresponding critical fugacity  $x = x(w)$ . Linearizing RG equations around this fixed point one gets only one relevant eigenvalue  $\lambda_1 = 8.6924; \dots$ , and the critical exponent is equal to  $\nu_{SAW} = \ln b / \ln \lambda_1 = 0.641076; \dots$ .

(2) The low temperature fixed point ( $A = 0; B = 0.356883; \dots = 22^{-1/3}$ ) is reached when starting with  $w > w_c$ . There is one relevant eigenvalue  $\lambda_1 = 16$ , and the critical exponent is  $\nu = 1/2$ .

(3) The tricritical fixed point ( $A = 0.19292; B = 0.3388; \dots$ ) corresponds to the collapse transition, when  $w = w_c$ . In this case there are two relevant eigenvalues:  $\lambda_1 = 15.4194; \dots$  and  $\lambda_2 = 5.52427; \dots$ . The critical exponent is equal to  $\nu_c = \ln b / \ln \lambda_1 = 0.506756; \dots$ .

#### Discussion

Values of  $\nu$  for SISAW on all so far studied 3d SF [2-4] are depicted on fig. 3. One can see that for any fractal  $\nu$  has its largest value in the high temperature phase, and that it decreases as  $b$  grows, approaching the

corresponding Euclidean value  $E_{SAW}^E = 0.588$  [5]. Respective values are smaller at the collapse transition, also decreasing with  $b$ , being close to the collapse Euclidean value  $E_c^E = 0.5$  [1]. Low temperature behavior is more interesting. In contrast to the  $b = 2$  case for which  $\nu = 1/d_f$  (meaning that polymer densely fills the underlying lattice, as on homogeneous lattices), critical exponent for  $b = 3$  and  $b = 4$  is larger than  $1/d_f$ , indicating that polymer is in semi-compact, rather than compact phase. This is due to the topological frustration, which is not the same for  $b = 3$  and  $b = 4$ , being also the reason for strange behavior of the low temperature RG fixed point:  $B$  is equal to  $22^{1/3}$  for both  $b = 2$  and  $b = 4$ , whereas for  $b = 3$  it is  $B = 1$  (of course,  $A = 0$  for any  $b$ ). It would be interesting to find out what happens beyond  $b = 4$ , but unfortunately the number of SISAW configurations grows extremely fast with  $b$ . Estimated computer time required for enumeration and sorting SISAWs for  $b = 5$  is not accessible, so one should search other methods for calculating  $\nu$ .




Figure 3: Critical exponent  $\nu$  for 3d SFs with  $b = 2, 3$ , and  $4$

#### References

- [1] P.G. de Gennes, "Scaling Concepts in Polymer Physics" (Itacha, NY : Cornell University Press, 1979)
- [2] D.Dhar and J.Vannimenus, J.Phys. A 20 (1987) 199
- [3] M.Knezevic and J.Vannimenus, J.Phys. A 20 (1987) L969
- [4] J.Maricic, "Random Walks on Three-dimensional Sierpinski Fractals" { diploma work (Faculty of Physics, University of Belgrade, 1998)
- [5] D.Lee et al, J.Stat.Phys. 80 (1995) 661; D.MacDonald et al, J.Phys. A 33 (2000) 5973