

Logarithmic corrections to finite size spectrum of $SU(N)$ symmetric quantum chains

Kingshuk Majumdar⁽¹⁾ and Manash Mukherjee⁽²⁾

⁽¹⁾ Department of Physics, Berea College, Berea, KY 40404,

⁽²⁾ Department of Physics, Miami University, Oxford, OH 45056.

We consider $SU(N)$ symmetric one dimensional quantum chains at finite temperature. For such systems the correlation lengths, ground state energy, and excited state energies are investigated in the framework of conformal field theory. The possibility of different types of excited states are discussed. Logarithmic corrections to the ground state energy and different types of excited states in the presence of a marginal operator, are calculated. Known results for $SU(2)$ and $SU(4)$ symmetric systems follow from our general formula.

PACS numbers: 75.10.Jm, 75.40.Gb

One of the fundamental models of solid state physics is the Heisenberg model of insulating magnets. In the one-dimensional case ("spin chains"), the spin-1/2 Heisenberg models have been studied extensively: most of our understanding of their quantum critical behavior is based on the Bethe-Ansatz solution for the ground state and excitation spectrum,^{1,2} mapping to the Sine-Gordon theory,³ non-abelian bosonization,⁴ and mapping to the sigma model.⁵ Although spin-1/2 Heisenberg chains are $SU(2)$ symmetric systems, fruitful generalizations have been accomplished in two different directions: (a) enlarging the representation of $SU(2)$ group to study quantum chains with higher spins, and (b) introducing higher symmetry groups such as $SU(N)$.

Here we consider generalizations of the type (b) and investigate how higher symmetry affects the ground state properties [Eqs. (8)-(10)] and finite size spectrum of quantum "spin" chains. Earlier studies of Aeck⁶ show that any one dimensional system with $SU(N)$ symmetry is critical, and at the very low energy scale these models are equivalent to $(N-1)$ free massless bosons. This system of free bosons – when viewed in the framework of two dimensional conformal field theory, are the primary fields of the $SU(N)_{k=1}$ WZNW model. Adopting this model we give an explicit derivation of logarithmic corrections to finite size spectrum of $SU(N)$ symmetric quantum chains. Logarithmic shifts in excited states energy levels have been theoretically observed for $SU(2)$ and $SU(4)$ symmetric systems away from the $T = 0$ quantum critical point.^{7,12} Known results for $N = 2$ and $N = 4$ follow from our general formulas [obtained in Eqs. (27)-(28) and the paragraphs below Eq. (30)].

An one-dimensional $SU(N)$ symmetric quantum chain of length L is described by the Hamiltonian¹³

$$H = \sum_{j=1}^{N-1} \sum_{A=1}^{N-1} S_j^A S_{j+1}^A \quad (1)$$

Here $n = L/a_0$ is the total number of discrete points

(a_0 being the lattice spacing) and S_j^A are the $(N^2 - 1)$ generators of the $SU(N)$ Lie algebra at each lattice site j . For convenience, the interaction strength and the lattice spacing a_0 have been set equal to one in Eq. (1). At each site j , the generators S_j^A can be represented by N "flavors" of fermions, a_j [$a = 1; \dots; N$]

$$S_j^A = \sum_{a,b=1}^{N-1} X_j^{ab} (T^A)_a^b \delta_{bj} \quad I=N; \quad (2)$$

where I is the identity operator, and T^A 's are a complete set of $(N^2 - 1)$ traceless normalized matrices so that $\text{Tr}[T^A T^B] = (1/2) \delta^{AB}$. Eq. (2) satisfies the constraint that at each site the total number of fermions is conserved, i.e. $\sum_{a=1}^N \sum_j^{N-1} a_j^a = 1$.

For such a fermionic system the theory can be bosonized using non-abelian bosonization at low temperatures. In the continuum limit, the bosonized Hamiltonian⁴ (H_e) can then be written in terms of the Kac-Moody currents,

$$H_e = \sum_{A=1}^{N-1} \int_{-\infty}^{\infty} dx J_L^A J_L^A + J_R^A J_R^A + 2J_L^A J_R^A; \quad (3)$$

where the normal ordered Kac-Moody currents for the left (with Fermi momentum $k_F < 0$) and the right moving (with $k_F > 0$) fermions are defined as,

$$J_L^A = : \sum_{a,b=1}^{N-1} (T^A)_a^b a_{-L}^a a_{-L}^b : ;$$

$$J_R^A = : \sum_{a,b=1}^{N-1} (T^A)_a^b a_R^a a_R^b : ; \quad (4)$$

At zero temperature ($T = 0$) the interaction term, $\sum_{A=1}^{N-1} \int_{-\infty}^{\infty} dx J_L^A J_R^A$, in Eq. (3) renormalizes to zero, and the sum of first two terms in the Hamiltonian that are quadratic in left and right moving currents, corresponds to the $SU(N)_{k=1}$ WZNW model. The fundamental unitary $N \times N$ matrix field g of the WZNW model is given by,

$$g_b^a = (\text{const}) : \frac{a_y}{L} R_b : : \quad (5)$$

The eld g transform s as the fundam ental representation of $SU(N)_L \times SU(N)_R$ which describes the exact sym m etry of the H am iltonian in Eq. (3) at zero tem perature. It is know n that this ferm ionic theory is equivalent to a theory of $(N-1)$ free massless bosons at the criticality with velocities v_s ; they correspond to $(N-1)$ excitation m odes of the $SU(N)$ sym m etric quantum chain that oscillates at different values of k_F .⁶ Furthermore, these oscillating m odes are prim ary elds of the $SU(N)_{k=1} \times WZNW$ m odel and their scaling dim ensions (Δ_p) can be obtained from¹⁴

$$\Delta_p = \frac{2C_p}{C_{adj} + 1} : \quad (6)$$

For $SU(N)$, C_p ($p = 1; 2; \dots; N-1$) is the eigenvalue of the Casim ir operator in the p -th fundam ental representation [having a Young tableau with p boxes in a single column], and $C_{adj} = N$ is the eigenvalue of the Casim ir operator in the adjoint representation [having a 2-column Young tableau with $(N-1)$ boxes in the first column and 1 box in the second column]. In a highest weight (λ) representation of $SU(N)$, the corresponding Casim ir eigenvalue is given by¹⁵

$$C = \frac{(\lambda, \lambda)}{2} = \frac{m}{N} + \sum_{i=1}^{r_0} \frac{X_i^2}{(b_i)^2} + \sum_{i=1}^{c_0} \frac{X_i^2}{(a_i)^2} ; \quad (7)$$

where λ is the highest weight corresponding to the adjoint representation and is norm alized to 1, m is the total number of boxes in the Young tableau with r_0 rows of length $b_1; b_2; \dots; b_{r_0}$ and c_0 columns of length $a_1; a_2; \dots; a_{c_0}$. Using this formula we find $C_p = p(N-p)(N+1)/2N$ and $C_{adj} = N$. Hence, the scaling dim ensions (Δ_p) of prim ary elds of the $SU(N)_{k=1} \times WZNW$ m odel are given by

$$\Delta_p = \frac{2C_p}{C_{adj} + 1} = \frac{p(N-p)}{N} : \quad (8)$$

For exam ple, in the case of $SU(4)$ the three oscillating components have scaling dim ensions $(3/4, 1, 3/4)$ [for $p = 1; 2; 3$]. For $SU(N)$, the m ode (dom inant) that oscillates at $k_F = 2\pi/N$ has a scaling dim ension $(1 - 1/N)$ [$p = 1$ or $(N-1)$ in this case].

Finite-size corrections to H eisenberg chain with $SU(2)$ and $SU(4)$ sym m etry has been studied using conform al eld theory.^{7,17} The relevance of studying finite size chains are twofold. One can not only compare the theoretical results with numerical simulations and experiments which are limited to finite size of the system but also can study the finite tem perature behavior of the system by identifying the finite size in the imaginary time

direction, that corresponds to finite tem perature. To obtain the finite-size corrections of a 1D chain of length L and with periodic boundary conditions we first introduce a conform al mapping from the finite plane (with coordinate z) to the cylinder (with coordinate w) via $w = (L/2\pi) \ln z$. Identifying the length to be inverse of the tem perature [$L = v_s/T$] the finite tem perature results¹⁷ of the ground state energy E_0 can be generalized to the $SU(N)$ sym m etric system ,

$$E_0(T) = E_0(0) - \frac{T(N-1)}{6v_s} : \quad (9)$$

Here $E_0(0)$ refers to the ground state energy at zero tem perature. The therm odynam ic quantities like specific heat and entropy can now be obtained by taking the appropriate derivatives with respect to the tem perature. Other quantities of interest are the finite tem perature corrections to the correlation lengths (ξ) of the different m odes. These inverse of the correlation lengths, ξ^{-1} are signature of energy gaps, $(E_n - E_0)$ between the ground state and the lowest lying excited states (E_n) that are created by finite tem perature of the system . Using the general formula for the scaling dim ension [Eq. (8)], we obtain ξ_p^{-1} of the p -th staggered m ode,

$$\xi_p^{-1} = \frac{E_n^p - E_0}{v_s} = \frac{2T}{v_s} \Delta_p = \frac{2T}{v_s} \frac{p(N-p)}{N} : \quad (10)$$

The tem perature dependence of the correlation lengths is in fact m odified by logarithmic corrections in the presence of marginal operators in the theory.¹⁷ The generic form of the H am iltonian at the critical point containing a marginal operator (ϕ ; t) is

$$H = H_0 + g_0 \int dx \phi(x, t) ; \quad (11)$$

where g_0 is the coupling constant and H_0 is the H am iltonian at the fixed point. In our case [Eq. (3)], the norm alized marginally irrelevant operator is

$$\phi = \sum_{A=1}^{N_X-1} D_A J_L^A J_R^A : \quad (12)$$

For such a marginally irrelevant operator, the H am iltonian at the critical point ($T = 0$) becomes equal to the fixed point H am iltonian. In Eq. (12), D is the norm alization constant to be determined from the two-point correlator of ϕ . The operator product expansion (OPE) of the $J_{L(R)}^A$ with any norm alized K ac-M oody prim ary eld is given by¹⁶

$$J_L^A(z)(z^0) = \frac{J_{0;L}^A}{2i(z-z^0)}(z^0) + \quad ; \quad (13)$$

$$J_R^A(z)(z^0) = \frac{J_{0;R}^A}{2i(z-z^0)}(z^0) + \quad ; \quad (14)$$

where the operators $J_{0;L}^A$ and $J_{0;R}^A$ are the generators of the global $SU(N)_L$ $SU(N)_R$ transformations, and satisfy the characteristic equations, $J_{0;L}^A j(z^0)i = P_{T_L^A} j(z^0)i$ and $J_{0;R}^A j(z^0)i = j(z^0)i T_R^A$. Note that $P_{T_L^A} (J_{0;L}^A)^2$ and $P_{T_R^A} (J_{0;R}^A)^2$ are the Casimir operators of $SU(N)_L$ and $SU(N)_R$ groups, respectively. The two-point correlators of the left currents (for $k=1$) are,

$$h J_L^A(z) J_L^B(z^0)i = \frac{\text{Tr}[T_L^A T_L^B]}{4^2 (z-z^0)^2} = \frac{A B}{8^2 (z-z^0)^2}; \quad (15)$$

$$h J_R^A(z) J_R^B(z^0)i = \frac{A B}{8^2 (z-z^0)^2}; \quad (16)$$

Using these results we explicitly calculate:

$$h(z;z)(z^0;z^0)i = \frac{D}{8^2} \frac{(N^2-1)}{(z-z^0)^2 (z-z^0)^2}; \quad (17)$$

and then compare it to the standard conformal field theory result i.e. $h(z;z)(z^0;z^0)i = \frac{1}{2} \frac{1}{(z-z^0)^2} \frac{1}{(z-z^0)^2}$ to obtain the value of the constant,

$$D = \frac{8^2}{N^2-1}; \quad (18)$$

Thus the normalized irrelevant marginal operator is given by,

$$(z;z) = \frac{8^2}{N^2-1} \sum_{A=1}^{N^2-1} J_L^A(z) J_R^A(z); \quad (19)$$

Perturbation to the normalized excited state (n) energies due to the marginal operator can now be calculated¹⁷ from

$$(E_n - E_0) = g_0 \int_0^Z dx h_n j_n i; \quad (20)$$

where j_n and j_n are Virasoro primary fields generated by applying Fourier modes of J_L^A and J_R^A on Kac-Moody primary fields. For large length (equivalently, small temperature), we may replace the coupling g_0 by its renormalization group improved value (upto the log-log term),¹⁸

$$g_0(T) = \frac{1}{b \ln(T_0=T)} \quad 1 - \frac{1}{2 \ln(T_0=T)} \ln[\ln(T_0=T)]; \quad (21)$$

Here T_0 is the model dependent parameter of the system and the coefficient b is defined via the following 3-point correlator,

$$h(z_1; z_1)(z_2; z_2)(z_3; z_3)i = \frac{1}{b} j_{12} j_{23} j_{13}; \quad (22)$$

Substituting this in Eq. (20) we obtain,

$$(E_n - E_0) = \frac{2T}{v_s \ln(T_0=T)} \quad \frac{2b_n}{b} \left(1 - \frac{1}{2 \ln(T_0=T)} \ln[\ln(T_0=T)] \right); \quad (23)$$

The coefficient b_n is again defined through the 3-point correlator,

$$h_n(z_1; z_1)(z_2; z_2)_n(z_3; z_3)i = \frac{1}{b_n} j_{12} j_{23} j_{13}^{x_n-2}; \quad (24)$$

Here x_n is the scaling dimension of the Virasoro primary field j_n . Substituting Eq. (19) in Eq. (24) and using the OPE's as in Eqs. (13) and (14) it follows that b_n is directly proportional to the sum of the product of the eigenvalues of the generators $J_{0;L}^A$ and $J_{0;R}^A$,

$$b_n = \frac{2}{N^2-1} \sum_{A=1}^{N^2-1} T_L^A T_R^A; \quad (25)$$

To evaluate $\sum_{A=1}^{N^2-1} T_L^A T_R^A$, we observe that the full symmetry, $SU(N)_L \times SU(N)_R$, of the quantum chain at the critical point is broken by the presence of the marginal operator, $(z;z)$. Only the diagonal $SU(N) \times SU(N)_L \times SU(N)_R$ is an exact symmetry of the quantum chain. Under this subgroup, the representation $V_L \times V_R$ of $SU(N)_L \times SU(N)_R$ decomposes into direct sum of various irreducible subrepresentations. If an excited state $(j_n i)$ belongs to a highest weight subrepresentation, $V_L \times V_R$ and C is the corresponding Casimir invariant of the diagonal $SU(N)$ in V , then we have¹⁹

$$\sum_{A=1}^{N^2-1} T_L^A T_R^A = \frac{1}{2} [C_L + C_R]; \quad (26)$$

where C_L and C_R are the Casimir invariants of $SU(N)_L$ and $SU(N)_R$ in the highest weight representations V_L and V_R , respectively. Therefore, using Eqs. (25) and (26) we find

$$b_n = \frac{1}{N^2-1} [C_L + C_R]; \quad (27)$$

The above formula may also be used to evaluate the renormalization group coefficient, b [Eq. (22)]: since

$(z; z)$ is a Virasoro primary field of conformal dimensions $(1, 1)$, we set $\chi_n(z; z) = (z; z)$ and $x_n = 2$ in Eq. (24), and hence, $b_h = b$. This can be seen as follows. The Virasoro primary fields J_L^A and J_R^A of conformal dimensions $(1; 0)$ and $(0; 1)$ transform as the adjoint representations, V_L^{adj} and V_R^{adj} of $SU(N)$, and since V_R^{adj} is conjugate to V_L^{adj} , the direct sum decomposition of $V_L^{\text{adj}} \oplus V_R^{\text{adj}}$ under the diagonal $SU(N) \rightarrow SU(N)_L \times SU(N)_R$ must contain a unique singlet. Hence, the Virasoro primary field $(z; z)$ in Eq. (19) transforms as this singlet representation and we have $C = 0; C_L = C_R = N$ in Eq. (27). This implies

$$b = \frac{2N}{p \frac{N^2 - 1}{N^2 - 1}} : \quad (28)$$

For example in the case of Heisenberg spin-chain with $SU(2)$ symmetry, $b = \frac{4}{p} = \frac{4}{3}$ and for the spin-orbital model with $SU(4)$, $b = \frac{8}{p} = \frac{8}{15}$. Our result for $SU(4)$ is new. Together with Eq. (21), the constant b also determines the correction of $O(g_0^3)$ in the ground state energy,

$$E_0(T) - E_0(0) = \frac{T}{6v_s} (N - 1) + 2^{-3} b g_0^3 : \quad (29)$$

To determine the logarithmic shifts in the excited states energy levels, we need the ratio $2b_h = b$ in Eq. (23). From Eqs. (27) and (28) we get,

$$\frac{2b_h}{b} = \frac{1}{N} [C - C_L - C_R] : \quad (30)$$

To evaluate b_h (and hence $2b_h = b$) we must know the excited states. In $SU(N)$ invariant quantum chains, the low lying excited states (j, n) correspond to $(N - 1)$ primary fields of the $SU(N)_{k=1}$ WZNW model with the scaling dimensions, Δ_p . These fields transform as $(q; q)$ representations of $SU(N)_L \times SU(N)_R$, where

$$q = \frac{N(N - 1)}{p!} (N - p + 1)$$

is the dimension of the p -th fundamental representation of $SU(N)$ for $p = 1; 2; \dots; N - 1$. For instance, the lowest excited states correspond to the fundamental primary field g with the scaling dimensions, $\Delta_1 = (1 - 1/N)$. This field transforms under the $(N; N)$ representation which decomposes under the diagonal $SU(N)$ into the adjoint and singlet representations. For the excited states, $\text{Tr}[gT^A]$, belonging to the adjoint representation, we have $C = N, C_L = C_R = (N^2 - 1)/2N$ which implies $b_h = \frac{1}{2} \frac{N^2 - 1}{N^2 - 1}$ and $2b_h = b = \frac{1}{2} N^2$. For example, in the case of $SU(2)$, this ratio $2b_h = b = \frac{1}{2} 4$; and for $SU(3)$ and $SU(4)$ they are $\frac{1}{2} 9$, and $\frac{1}{2} 16$ respectively.

For the excited state, $\text{Tr} g$, belonging to the singlet representation we have $C = 0; C_L = C_R = (N^2 - 1)/2N$. In this case, $b_h = \frac{1}{2} \frac{N^2 - 1}{N^2 - 1} = \frac{1}{2} N$ and the $2b_h = b = \frac{1}{2} 1 = \frac{1}{2} N^2$. This result is new. In case of $SU(2)$, the value $2b_h = b = \frac{3}{2} 4$ has been previously obtained⁷ but for $SU(3)$, $2b_h = b = \frac{8}{3} 9$ and for $SU(4)$, $2b_h = b = \frac{15}{4} 16$ are predictions from our general formula.

We consider one more application of the formula (30) of current interest { the $SU(4)$ symmetric quantum chain described by the $SU(4)_{k=1}$ WZNW model. In this case, to compute logarithmic corrections to the excited states energy we note that there are three primary fields with scaling dimensions $\Delta_p = 3/4; 1; 3/4$ for $p = 1; 2; 3$ respectively, as seen from Eq. (8). The case of $p = 1$ [and $p = 3$], as discussed above, is the fundamental field g [and its hermitian conjugate g^\dagger] which transforms under the $(4; 4)$ [and $(4; 4)$] representation of $SU(4)_L \times SU(4)_R$. From Eq. (10), the next lowest energy excited states correspond to the primary field operator (denoted by ψ) with $\Delta_2 = 1$. The field transforms under the $(6; 6)$ representation of $SU(4)_L \times SU(4)_R$. This $(6; 6)$ representation decomposes as direct sum of a singlet, an adjoint and a 20-dimensional representation (as in Fig. 1) under the diagonal $SU(4)$.

$\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \otimes \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} \oplus \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \oplus \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$

(6) (6) (1) (15) (20)

FIG. 1. Young tableau for the decomposition of $(6; 6)$ representation of $SU(4)$. The number in the parenthesis denotes the dimension of the corresponding representation.

We now compute the ratio $2b_h = b$ for the excited states corresponding to 20-dimensional representation which has a Young tableau with 2 rows and 2 columns. For this representation, the Casimir invariant C in Eq. (30) is obtained from the formula (7): we find $C = 6$, and $C_L = C_R = C_{p=2} = 5/2$. Thus, for the excited states corresponding to ψ in the 20-dimensional subrepresentation of $(6; 6)$, we have $2b_h = b = \frac{1}{2} 4$.

In summary, we have studied finite size spectrum for one dimensional $SU(N)$ symmetric quantum chains using both conformal field theory and representation theory of $SU(N)$. We have calculated in general the scaling dimensions of all the oscillating modes, and obtained the ground state energy as well as correlation lengths of the staggered modes for a finite size system with $SU(N)$ symmetry. Possibilities of different types of excited states

are also briefly discussed and a general formula to compute the logarithmic correction to the excited state energies has been derived. The existing results for $N = 2, 4$ agree with the predictions from our general formula.

Authors acknowledge discussions with F.C. Zhang and M.M. a.K.M. also acknowledge partial support by PRF No. 33611-AC6 and partial support from Berea College.

-
- ¹ H. Bethe, Z. Phys. 71, 205 (1931).
 - ² A. Luther and I. Peschel, Phys. Rev. B 12, 3908 (1975).
 - ³ C. N. Yang and C. P. Young, J. Math. Phys. 10, 1115 (1969).
 - ⁴ I. A. A. eck, Nucl. Phys. B 265, 409 (1986); I. A. eck, Phys. Rev. Lett. 55, 1355 (1985); Phys. Rev. Lett. 54, 966 (1985).
 - ⁵ F. D. M. Haldane, Phys. Rev. Lett. 45, 1358 (1980).
 - ⁶ I. A. eck, in Les Houches, session XLIX, Champs, Cordes et Phenomenes Critiques/Fields, strings and critical phenomena, Elsevier, New York, 1989.
 - ⁷ I. A. eck, D. Gepner, H. J. Schultz, and T. Ziman, J. Phys. A 22, 511 (1989).
 - ⁸ C. Itoi, S. Qin, and I. A. eck, Phys. Rev. B 61, 6747 (2000).
 - ⁹ I. A. eck, Phys. Rev. Lett. 55, 1355 (1985); I. A. eck, J. Phys. A 31, 4573 (1998).
 - ¹⁰ B. Frischmuth, F. Mila, and E. Troyer, Phys. Rev. Lett. 82, 835 (1999).
 - ¹¹ P. A. zaria, E. Boulat, and P. Lecheminant, Phys. Rev. B 61, 12112 (2000); P. A. zaria, A. O. Gogolin, P. Lecheminant, and A. A. Nersisyan, Phys. Rev. Lett. 83, 624 (2000).
 - ¹² Y. Yamashita, N. Shibata, and K. Ueda, Physica B 281, 542 (2000).
 - ¹³ B. Sutherland, Phys. Rev. B 12, 3795 (1975).
 - ¹⁴ V. G. Knizhnik and A. B. Zamolodchikov, Nucl. Phys. B 247, 83 (1984).
 - ¹⁵ J. Fuchs and C. Schweigert Symmetries, Lie Algebras and Representations (Cambridge University Press, 1997).
 - ¹⁶ P. Di Francesco, P. Mathieu, and D. Senechal Conformal Field Theory (Springer, 1996).
 - ¹⁷ J. L. Cardy, Nucl. Phys. B 240, 514 (1984); J. L. Cardy, J. Phys. A 19, L1093 (1986).
 - ¹⁸ K. Nomura and M. Yamada, Phys. Rev. B 43, 8217 (1991).
 - ¹⁹ P. I. Etingof, I. B. Frenkel and A. A. Kirillov Lectures on Representation Theory and Knizhnik-Zamolodchikov Equations (American Mathematical Society, 1998).