

## Glauber Critical Dynamics: Exact Solution of the Kinetic Gaussian Model

Jian-Yang Zhu<sup>a,b,c</sup> and Z.R. Yang<sup>a,b</sup><sup>a</sup>CCAST (World Laboratory), Box 8730, Beijing 100080, China<sup>b</sup>Department of Physics, Beijing Normal University, Beijing 100875, China<sup>y</sup><sup>c</sup>Department of Physics, Jiangxi Normal University, Nanchang 330027, China

In this paper, we have exactly solved Glauber critical dynamics of the Gaussian model on three dimensions. Of course, it is much easy to apply to low dimensional case. The key steps are that we generalize the spin change mechanism from Glauber's single-spin flipping to single-spin transition and give a normalized version of the transition probability. We have also investigated the dynamical critical exponent and found surprisingly that the dynamical critical exponent is highly universal which refer to that for one- two- and three-dimensions they have same value independent of spatial dimensionality in contrast to static (equilibrium) critical exponents.

PACS numbers: 64.60.Ht, 75.10.Hk

## I. INTRODUCTION

Irreversible dynamical systems exhibit complicated and interesting non-equilibrium phenomena near the critical point. The study of non-equilibrium statistical mechanics is much more difficult than equilibrium state due to the complexity. However, the interesting dynamical critical behaviors have been attracting a large number of researchers to work hard for many years.

Up to now, there is no general theory based on the first principle to describe the dynamical problems. However, a great progress has been made since the pioneering work completed by Glauber [1] and Kawasaki [2]. According to their theory, the time-evolving of the order parameters is described by Markov processes with Glauber single spin flipping mechanism or Kawasaki exchange mechanism between two spins. Since then people have been paid much attentions to the study of critical dynamics. The research so far have been extended from the kinetic Ising model to the kinetic Potts model and from integer to fraction dimension, in which many approximate methods such as Monte Carlo simulation, high-temperature series expansion,  $\epsilon$ -expansion, bond-moving renormalization-group method, etc. have been applied [3,19].

Now let us turn on master equation, a basic equation for treating critical dynamics. As we know, the key step for solving master equation is the determination of the transition probability. Usually the transition probability between different states, i.e., different spin configurations of the system, is only determined in terms of the detailed balance condition. Since such a choice is not unique, then leaves behind some arbitrariness. For removing the arbitrariness, at least in part, we suggest a normalized transition probability, this means in unit time interval the transition may occur or may not. We apply this point to continuous spin (Gaussian) system and obtain exact solutions of one-, two- and three dimensional kinetic Gaussian model.

This paper is organized as follows. In Sec.II, a single-spin transiting critical dynamics which suits arbitrary spin systems is presented. As an application, the Gaussian model is treated in Sec.III. We not only obtain the exact solutions of 1D, 2D and 3D kinetic Gaussian model, but also find that the dynamical critical exponent is highly universal. Finally, the conclusion is devoted to Sec.IV.

## II. FORMALIZATION

For an irreversible dynamical system subjected to the time-dependent perturbation, once the perturbation is removed, the system will very slowly approach to equilibrium state because of the large-scale fluctuation near the critical point, which is what we call the critical slowing down phenomenon. The critical dynamics will attempt to explain why the short-range interactions lead to long-time relaxation. Due to the complexity there have been no microscopic theory

---

E-mail: zhu.jy@bnu.edu.cn

<sup>y</sup>Mailing address

based on the first principle so far, thus a suitable theoretical model will be quite important. As already mentioned, both the Glauber dynamics and Kawasaki dynamics have proven to be successful in many dynamic systems.

In this section, we give a brief review of Glauber dynamics. Of course, we will give some improvement so that it can suits arbitrary-spin systems. For cleanness, we start from the one-dimensional case (1D), and then the formulation is easy to extend to 2D and 3D.

The 1D lattice-spin model we shall discuss is a stochastic one. The spins of  $N$  fixed particles are represented as stochastic functions of time  $\sigma_j(t)$ , ( $j = 1; \dots; N$ ), which can be taken discrete values (discrete-spin model) or continuous values (continuous-spin model), and make transitions among these values. The transition, according to the Glauber dynamics, can only change single spin value each time, such as  $\sigma_j(t) \rightarrow \sigma_j'(t)$  because of the interacting of the system with heat reservoir. The transition probability  $W_j(\sigma_j(t) \rightarrow \sigma_j'(t))$  from configuration  $(\sigma_1(t); \sigma_2(t); \dots; \sigma_j(t); \dots; \sigma_N(t))$  to configuration  $(\sigma_1(t); \sigma_2(t); \dots; \sigma_j'(t); \dots; \sigma_N(t))$ , in general, depends on the momentary values of the neighboring spins as well as on the influence of the heat bath. For this reason statistical correlations exist between different spins. Therefore, it is necessary to deal with the entire  $N$ -spin system as a unit. The evolution of spin functions describing system form a Markov process of  $N$  discrete or continuous random variables with a continuous time variable as argument.

We introduce a probability distribution function  $p(\sigma_1; \dots; \sigma_N; t)$ , which denotes the probability of spin system being in the state  $(\sigma_1; \dots; \sigma_N)$  at time  $t$ . Let  $W_j(\sigma_j \rightarrow \sigma_j')$  be the transition probability per unit time that the  $j$ th spin transits from one value  $\sigma_j$  to another possible value  $\sigma_j'$ , while the others remain fixed. Then, on the supposition of single-spin transition, we may write the time derivative of the function  $P(\sigma_1; \dots; \sigma_N; t)$  as

$$\frac{d}{dt}P(\sigma; t) = \sum_j \sum_{\sigma_j'} [W_j(\sigma_j \rightarrow \sigma_j')p(\sigma; t) + W_j(\sigma_j' \rightarrow \sigma_j)p(\sigma_j'; \sigma; t)] \quad (1)$$

This is a probability equation, in which the first term in the hand-right side of Eq.(1) denotes the decrease of the probability distribution function  $P(\sigma; t)$  per unit time due to the transition of the spin state from the initial value  $\sigma_j$ , ( $j = 1; 2; \dots; N$ ) to various possible final values and the second term denotes the increase of the probability distribution function  $P(\sigma; t)$  per unit time due to the transition of the spin state from the various possible initial values  $\sigma_j'$ , ( $j = 1; 2; \dots; N$ ) to final value  $\sigma_j$ . We shall refer to the Eq.(1) as the master equation since its solution would contain the most complete description of the system available.

It is the most crucial step, obviously, that the transition probability must be determined before the master equation can be solved. Then, how to determine the transition probability? For this problem, Glauber's theory leaves some leeway to select. However, inappropriate selection will probably make the problem difficult to solve. So, we hope to find out a more definite expression to apply the Glauber's theory to arbitrary-spin systems. Now we consider it from both mathematics and physics aspects. In mathematics, generally speaking, the probability must be ergodic and positive definite, and can be normalized; in physics, we often request that an equilibrium thermodynamic system satisfies the detailed balance condition. Based on these considerations, we can restrict the spin transition probability  $W_j(\sigma_j \rightarrow \sigma_j')$  to satisfy the following conditions so that ensure the system in thermodynamic equilibrium state finally:

For the set  $(S; \hat{S})$  composed of a subset  $S$  and its dual subset  $\hat{S}$  in phase space, existing  $\sigma_j$  belonging to  $S$  and  $\sigma_j'$  belonging to  $\hat{S}$ , we have

(1) ergodicity

$$\forall \sigma_j, \sigma_j' : W_j(\sigma_j \rightarrow \sigma_j') \neq 0; \quad (2)$$

(2) positivity

$$\forall \sigma_j, \sigma_j' : W_j(\sigma_j \rightarrow \sigma_j') \geq 0; \quad (3)$$

(3) normalization

$$\forall \sigma_j : \sum_{\sigma_j'} W_j(\sigma_j \rightarrow \sigma_j') = 1; \quad (4)$$

(4) detailed balance

$$\forall \sigma_j, \sigma_j' : \frac{W_j(\sigma_j \rightarrow \sigma_j')}{W_j(\sigma_j' \rightarrow \sigma_j)} = \frac{P_{eq}(\sigma_1; \dots; \sigma_j; \dots; \sigma_N)}{P_{eq}(\sigma_1; \dots; \sigma_j'; \dots; \sigma_N)}; \quad (5)$$

in which

$$P_{eq} = \frac{1}{Z} \exp[-H(f, g)]; \quad Z = \sum_{f, g} \exp[-H(f, g)];$$

where  $P_{eq}$  is the equilibrium Boltzmann distribution function,  $Z$  is the partition function,  $H(f, g)$  is the system Hamiltonian.

Although the spin transition probabilities are not determined uniquely by above restriction conditions, there are less room to choose them. Furthermore, considering the fact that the transition of the individual spin depends merely on the momentary values of the neighboring spins as well as on the influence of the heat bath, we can suppose that the transition probability from  $s_j$  to  $s_j'$  depend only on the heat Boltzmann factor of the neighboring spins, i.e.

$$W_i(s_i \rightarrow s_i') / \exp[-H_i(s_i)] = \sum_{\langle ij \rangle} \exp[-H_j(s_j')] A_j^5;$$

or

$$W_i(s_i \rightarrow s_i') = \frac{1}{Q_i} \exp[-H_i(s_i')] \sum_{\langle ij \rangle} \exp[-H_j(s_j')] A_j^5; \quad (6)$$

where  $\sum_{\langle ij \rangle}$  means that the summation for  $j$  is only related to the neighboring values of  $i$ . By means of the normalized condition (4), the normalized factor  $Q_i$  can be determined as

$$Q_i = \sum_{s_i} \exp[-H_i(s_i)] \sum_{\langle ij \rangle} \exp[-H_j(s_j')] A_j^5; \quad (7)$$

Obviously,  $Q_i$  is independent of  $s_i$  and is related to the temperature and neighboring spins.

Compared with Glauber's expression [1], Eq.(6) is a normalized version of transition probability. As we know, the constant in Glauber's expression is a free constant determined by the time scale. Actually, our expression is only a definite selection for constant by extra restriction conditions and physical considerations.

Usually, we are interested in local magnetization and the spin-pair correlation, they are defined as follows [1]

$$q_k(t) = \langle s_k(t) \rangle = \sum_{f, g} s_k P(f, g; t); \quad (8)$$

$$k_1(t) = \langle s_k(t) s_1(t) \rangle = \sum_{f, g} s_k s_1 P(f, g; t); \quad (9)$$

According to the definitions (8)–(9) and the master equation (1), and using the normalized condition (4), time-evolving equations of  $q_k(t)$  and  $k_1(t)$  can be derived (see Appendix A)

$$\frac{d}{dt} q_k(t) = -q_k(t) + \sum_{f, g} \sum_{s_k} s_k W_k(s_k \rightarrow s_k') P(f, g; t); \quad (10)$$

$$\begin{aligned} \frac{d}{dt} k_1(t) = & -2k_1(t) + \sum_{f, g} \sum_{s_k} s_k \sum_{s_1} s_1 W_1(s_1 \rightarrow s_1') \\ & + \sum_{f, g} \sum_{s_k} s_k W_k(s_k \rightarrow s_k') P(f, g; t); \end{aligned} \quad (11)$$

Similarly, time-evolving equation of equal-time multi-spin correlation function can be further derived

(12)

All of these formulas can readily be extended to spin systems on square lattice and cubic lattice. The only corrections are changing subscript  $i$  into  $ij$  and  $ijk$  respectively.

In this section, three examples of the application are given, including one-, two-, and three-dimensional kinetic Gaussian model.

The Gaussian model, proposed by T.H. Berlin and M.Kac [20] at first in order to make Ising model more tractable, is an continuous-spin model. Comparing with the Ising model, besides having the same Hamiltonian form (three-dimensional case)

(13)

$\mathbf{y}_{ijk}$  and  $\mathbf{y}_{ijk} + \mathbf{d}_{ijk}$  is assumed to be the Gaussian-type distribution

(14)

We now proceed to treat the isotropic kinetic Gaussian model on the cube lattice. The system Hamiltonian and the spin distribution probability are (13) and (14) respectively. In this case the spin transition probability can be expressed as

(15)

$$X \int_1^{\infty} \frac{Z_1}{f(\cdot)} d\cdot; \quad (16)$$
$$Q_{ijk} = \frac{d_{ijk}^8}{k^2} \exp \left( i \omega_{ijk} t + i \omega_{jk} z + i \omega_{jk+w} w \right) \quad (17)$$

and the another useful combination formula can also be obtained

$$\begin{aligned} \sum_{ijk} \hat{c}_{ijk} W_{ijk} (\hat{c}_{ijk} - \hat{c}_{ijk}) &= \sum_{ijk} \hat{c}_{ijk} W_{ijk} (\hat{c}_{ijk} - \hat{c}_{ijk}) f(\hat{c}_{ijk}) d\hat{c}_{ijk} \\ &= \frac{k}{b} \sum_w (c_{i+w,j;k} + c_{i;j+w,k} + c_{i;j,k+w}) : \end{aligned} \quad (18)$$

Substituting (18) into the following time-evolving equations of the local magnetization and the equal-time spin-pair correlation function

$$\frac{d}{dt} c_{ijk}(t) = c_{ijk}(t) + \sum_{fg} \sum_{ijk} \hat{c}_{ijk} W_{ijk} (\hat{c}_{ijk} - \hat{c}_{ijk}) A_P(fg;t); \quad (19)$$

$$\begin{aligned} \frac{d}{dt} c_{ijk;i^0j^0k^0}(t) &= 2 c_{ikj;i^0j^0k^0}(t) + \sum_{fg} \sum_{ijk} \sum_{i^0j^0k^0} \hat{c}_{i^0j^0k^0} W_{i^0j^0k^0} (\hat{c}_{i^0j^0k^0} - \hat{c}_{i^0j^0k^0}) A_{13} \\ &+ \sum_{i^0j^0k^0} \sum_{ijk} \hat{c}_{ijk} W_{ijk} (\hat{c}_{ijk} - \hat{c}_{ijk}) A_{5P}(fg;t); \end{aligned} \quad (20)$$

we get

$$\frac{d}{dt} c_{ijk}(t) = c_{ijk}(t) + \frac{k}{b} \sum_w (c_{i+w,j;k} + c_{i;j+w,k} + c_{i;j,k+w}); \quad (21)$$

$$\begin{aligned} \frac{d}{dt} c_{ijk;i^0j^0k^0}(t) &= 2 c_{ikj;i^0j^0k^0}(t) + \frac{k}{b} \sum_w (c_{ijk;i^0+j^0,k^0}(t) + c_{ijk;i^0,j^0+w,k^0}(t) \\ &+ c_{ijk;i^0,j^0;k^0+w}(t) + c_{i+w,j;k;i^0j^0k^0}(t) + c_{i;j+w,k;i^0j^0k^0}(t) + c_{i;j,k+w;i^0j^0k^0}(t)) : \end{aligned} \quad (22)$$

In order to solve Eqs.(21) and (22) in the nearest-neighbor interaction case ( $w = 1$ ), we introduce two generating functions[]:

$$F_1(\lambda_1; \lambda_2; \lambda_3; t) = \sum_{i,j,k=1}^X \lambda_1^i \lambda_2^j \lambda_3^k c_{ijk}(t); \quad (23)$$

and

$$F_2(\lambda_1; \lambda_6; t) = \sum_{i,j,k;i^0,j^0,k^0=1}^X \lambda_1^i \lambda_2^j \lambda_3^k \lambda_4^{i^0} \lambda_5^{j^0} \lambda_6^{k^0} c_{ijk;i^0j^0k^0}(t); \quad (24)$$

then the Eqs.(21) and (22) turn into the following equations with respect to  $F_1$  and  $F_2$ , respectively,

$$\frac{d}{dt} F_1(\lambda_1; \lambda_2; \lambda_3; t) = \left( 1 + \frac{k}{b} \sum_{i=1}^X \lambda_1^i + \lambda_1^{-1} \right) F_1(\lambda_1; \lambda_2; \lambda_3; t); \quad (25)$$

$$\frac{d}{dt} F_2(\lambda_1; \lambda_6; t) = \left( 2 + \frac{k}{b} \sum_{i=1}^X \lambda_1^i + \lambda_1^{-1} \right) F_2(\lambda_1; \lambda_6; t); \quad (26)$$

Solving Eqs.(25) and (26), we get

$$F_1(\lambda_1; \lambda_2; \lambda_3; t) = F_1(\lambda_1; \lambda_2; \lambda_3; 0) e^t \exp \left( \frac{k}{b} \sum_{i=1}^X \lambda_1^i + \lambda_1^{-1} \right) t; \quad (27)$$

$$F_2(\mathbf{x}; \mathbf{t}) = F_2(\mathbf{x}; 0) e^{2t} \exp \left[ \sum_{i=1}^N \frac{k_i}{b} X_i^6 \right] \quad (28)$$

In terms of a generating function of the Bessel functions of imaginary argument

$$e^{x(\frac{1}{2} + v)} = \sum_{v=1}^{\infty} \frac{x^v}{v!} I_v(x); \quad (29)$$

we obtain immediately the following exact solutions

$$q_{ijk}(\mathbf{t}) = e^{2t} \sum_{n,m,l=1}^{\infty} q_{nml}(0) I_{i-n} \left( \frac{2k}{b} t \right) I_{j-m} \left( \frac{2k}{b} t \right) I_{k-l} \left( \frac{2k}{b} t \right); \quad (30)$$

$$q_{ijk;i^0 j^0 k^0}(\mathbf{t}) = e^{2t} \sum_{n,m,l;n^0,m^0,l^0=1}^{\infty} q_{nml;n^0 m^0 l^0}(0) I_{i-n} \left( \frac{2k}{b} t \right) I_{j-m} \left( \frac{2k}{b} t \right) I_{k-l} \left( \frac{2k}{b} t \right) I_{i^0-n^0} \left( \frac{2k}{b} t \right) I_{j^0-m^0} \left( \frac{2k}{b} t \right) I_{k^0-l^0} \left( \frac{2k}{b} t \right); \quad (31)$$

where  $q_{nml}(0)$  and  $q_{nml;n^0 m^0 l^0}(0)$ , respectively, correspond to their initial values.

By using of the asymptotic expansion expression of the first-kind imaginary argument Bessel function

$$I_v(x) = \frac{e^x}{\sqrt{2\pi x}} \sum_{n=0}^{\infty} \frac{(-1)^n (v;n)}{(2x)^n} + \frac{e^{-x + (v + \frac{1}{2})i}}{\sqrt{2\pi x}} \sum_{n=0}^{\infty} \frac{(v;n)}{(2x)^n}; \quad (-2 < \arg x < 3/2); \quad (32)$$

where

$$(v;n) = \frac{\frac{1}{2} + v + n}{n!} \frac{1}{\frac{1}{2} + v - n};$$

we can obtain the long-time asymptotic behavior of the local magnetization

$$q_{ijk}(\mathbf{t}) \sim \left( \frac{2k}{b} t \right)^{-3/2} e^{-(1/6k=b)t} \sum_{n,m,l=1}^{\infty} q_{nml}(0) \frac{1}{t^{3/2}} e^{-t/\tau}; \quad (33)$$

$$= \frac{1}{1 - 6k=b}; \quad (34)$$

where  $\tau$  is the relaxation time of the system. We know that the critical point of the Gaussian model is  $k_c = J/k$ ,  $T_c = b/2d$ , where  $d$  is the spatial dimension, and the correlation length critical exponent is  $\nu = 1/2$  [22]. So, by means of the following dynamical scaling hypotheses

$$T - T_c \sim \tau^{-z}; \quad (35)$$

$$\tau \sim \xi^z; \quad (36)$$

the dynamic critical exponent  $z$  of the 3D kinetic Gaussian model can be obtained

$$z = 2; \quad (37)$$

With the same treatment, we can easily solve one- and two-dimensional kinetic Gaussian model. Ignoring the process of solution, we give only the following exact results:

(1). 1D case

$$q_k(t) = e^{-t} \sum_{m=1}^X q_m(0) I_{k-m} \left( \frac{2k}{b} t \right); \quad (38)$$

$$q_{kl}(t) = e^{-2t} \sum_{n,m=1}^X q_{nm}(0) I_{k-n} \left( \frac{2k}{b} t \right) I_{l-m} \left( \frac{2k}{b} t \right); \quad (39)$$

$$= \frac{1}{1 - 2k/b}; \quad (40)$$

$$z = 2; \quad (41)$$

(2). 2D case

$$q_{nm}(t) = e^{-t} \sum_{k,l=1}^X q_{kl}(0) I_{n-k} \left( \frac{2k}{b} t \right) I_{m-l} \left( \frac{2k}{b} t \right); \quad (42)$$

$$q_{n_1 n_2 \dots n_0}(t) = e^{-2t} \sum_{i,j,i^0,j^0=1}^X q_{ij,i^0,j^0}(0) I_{n_1-i} \left( \frac{2k}{b} t \right) I_{n_2-j} \left( \frac{2k}{b} t \right) I_{n_0-i^0} \left( \frac{2k}{b} t \right) I_{n_0-j^0} \left( \frac{2k}{b} t \right) \quad (43)$$

$$= \frac{1}{1 - 4k/b}; \quad (44)$$

$$z = 2; \quad (45)$$

#### IV. CONCLUSION

In this paper, we have suggested a normalized transition probability to solve the time evolution equations of the local magnetization and spin-pair correlation function. Our treatment can in part remove the arbitrariness in Glauber dynamical theory, and makes exactly solve the time evolution equation possible.

Based on our treatment, we have re-investigated one-dimensional kinetic Ising model, resulting in same dynamical critical exponent  $z$  as Glauber[1]. We have also exactly solved the kinetic Gaussian model, and given the detail of solving process of three-dimensional case. We have in surprise found that the dynamical critical exponents have same value independent of spatial dimension, which shows the dynamical behavior has super-universality in contrast with static behavior. In fact, in equilibrium phase transition the critical exponents are strongly dependent on dimensionality.

#### ACKNOWLEDGMENTS

The work was supported by the National Basic Research Project "Nonlinear Science" and the National Natural Science Foundation of China.

#### APPENDIX A: PROOFS OF EQS. (10)-(12)

According to the definition (8) and using the master equation (1), we have

$$\begin{aligned}
\frac{d}{dt}q_k(t) &= \frac{d}{dt} \sum_k^X P(f, g; t) \\
&= \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_i(i \rightarrow \hat{i})P(f, g; t) + w_i(\hat{i} \rightarrow i)P(f, g; \hat{i}; t)] \\
&= \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_i(i \rightarrow \hat{i})P(f, g; t) + w_i(\hat{i} \rightarrow i)P(f, g; \hat{i}; t)] \\
&\quad + \sum_{i \neq k}^{f, g} \sum_k^X [w_k(k \rightarrow \hat{k})P(f, g; t) + w_k(\hat{k} \rightarrow k)P(f, g; \hat{k}; t)] \tag{A 1}
\end{aligned}$$

Looking at the first term ( $i \neq k$ ) in the latter of the last equality sign of (A 1),

$$\begin{aligned}
(i \neq k) \text{ term} &= \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_i(i \rightarrow \hat{i})P(f, g; t) + w_i(\hat{i} \rightarrow i)P(f, g; \hat{i}; t)] \\
&= \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_i(i \rightarrow \hat{i})P(f, g; t) + w_i(\hat{i} \rightarrow i)P(f, g; \hat{i}; t)] \\
&\quad + \sum_{i \neq k}^{f, g} \sum_k^X [w_k(k \rightarrow \hat{k})P(f, g; t) + w_k(\hat{k} \rightarrow k)P(f, g; \hat{k}; t)] \tag{A 2}
\end{aligned}$$

it is easy to see that this term equals to zero, as long as  $\hat{i}$  exchange with  $i$  before doing sum for  $\hat{i}$  and  $i$ . So the surplus term of (A 1) is only the last term ( $i = k$ ):

$$\begin{aligned}
\frac{d}{dt}q_k(t) &= \frac{d}{dt} \sum_k^X P(f, g; t) \\
&= \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_k(k \rightarrow \hat{k})P(f, g; t) + w_k(\hat{k} \rightarrow k)P(f, g; \hat{k}; t)] \\
&\quad + \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_k(k \rightarrow \hat{k})P(f, g; t) + w_k(\hat{k} \rightarrow k)P(f, g; \hat{k}; t)] \\
&= \sum_k^X P(f, g; t) + \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_k(k \rightarrow \hat{k})P(f, g; t) + w_k(\hat{k} \rightarrow k)P(f, g; \hat{k}; t)] \\
&= q_k(t) + \sum_k^X \sum_{i \neq k}^{f, g} \sum_i^X [w_k(k \rightarrow \hat{k})P(f, g; t) + w_k(\hat{k} \rightarrow k)P(f, g; \hat{k}; t)]
\end{aligned}$$

in which, the normalized condition and the technic of exchange of  $\hat{i}$  for  $i$  were used. Hitherto, Eq.(10) have been proven exactly. As for the proof of the Eqs.(11) and (12), it is easily accessible via the same method, so without the necessity for further proof.

- [1] R. J. Glauber, J. Math. Phys. 4, 294 (1963).
- [2] K. Kawasaki, Phys. Rev. 145, 224 (1965).
- [3] G. Forgacs, S. T. Chui and H. L. Frisch, Phys. Rev. B 22, 415 (1980).
- [4] E. J. Lage, J. Phys. A : Math. Gen. 18, 2289 (1985).
- [5] E. J. Lage, J. Phys. A : Math. Gen. 18, 2411 (1985).
- [6] E. J. Lage, Phys. Lett. A 127, 9 (1988).
- [7] Y. Achiam, Phys. Rev. B 31, 4732 (1985).
- [8] Y. Achiam, Phys. Rev. B 32, 1796 (1985).
- [9] Y. Achiam, Phys. Rev. B 33, 7762 (1986).
- [10] Jian Zhou and Z. R. Yang, Phys. Rev. B 39, 9423 (1989).
- [11] M. D. Lacasse, J. Vinals and M. Grant, Phys. Rev. B 47, 5646 (1993).
- [12] B. C. S. G. randi and W. Figueiredo, Phys. Rev. E 54, 4722 (1996).



- [13] J. Rogers and J. O. Indekeu, Phys. Rev. B 41, 6998 (1990).
- [14] J. Wang, Phys. Rev. B 47, 896 (1993).
- [15] Y. Achiam and J. M. Kosterlitz, Phys. Rev. Lett. 41, 128 (1978).
- [16] G. F. M. Azenko and O. T. Valls, Phys. Rev. B 24, 1419 (1981).
- [17] P. C. Hohenberg and B. I. Halperin, Rev. Mod. Phys. 49, 435 (1977).
- [18] Z. R. Yang, Phys. Rev. B 46, 11578 (1992).
- [19] Peiqing Tong, Phys. Rev. E 56, 1371 (1997).
- [20] T. H. Berlin and M. Kac, Phys. Rev. 86, 821 (1952).
- [21] J. J. Binney, N. J. Dowrick, A. J. Fisher and M. E. J. Newman, The Theory of Critical Phenomena, Clarendon Press, Oxford 1992.
- [22] L. E. Reichl, A Modern course in Statistical Physics, University of Texas Press, Austin, 1980.
- [23] Song Li and Z. R. Yang, Phys. Rev. E 55, 6656 (1997).