

The depinning transition of a driven interface in the random – eld Ising model around the upper critical dimension

L. Roters and K.D. Usadel[†]

Theoretische Tieftemperaturphysik, Gerhard-Mercator-Universität, Lotharstr. 1, 47048 Duisburg, Germany

S. Lubeck[‡]Weizmann Institute of Science Department of Complex Physics, 76100 Rehovot, Israel and
Theoretische Tieftemperaturphysik, Gerhard-Mercator-Universität, Lotharstr. 1, 47048 Duisburg, Germany
(dated: March 22, 2022)

We investigate the depinning transition for driven interfaces in the random – eld Ising model for various dimensions. We consider the order parameter as a function of the control parameter (driving eld) and examine the effect of thermal fluctuations. Although thermal fluctuations drive the system away from criticality the order parameter obeys a certain scaling law for sufficiently low temperatures and the corresponding exponents are determined. Our results suggest that the so-called upper critical dimension of the depinning transition is five and that the system belongs to the universality class of the quenched Edwards-Wilkinson equation.

PACS numbers: 68.35.Rh, 75.10.Hk, 75.40.Mg

I. INTRODUCTION

Driven Interfaces in quenched disordered systems display with increasing driving force a transition from a pinned interface to a moving interface (see e.g. [1] and references therein). This so-called depinning transition is caused by a competition of the driving force and the quenched disorder. The first one tends to move the interface whereas the latter one hinders the movement. Depinning transitions are observed in a large variety of physical problems, such as fluid invasion in porous materials (see, for instance, Sec. 6.2 in [2] and references therein), depinning of charge density waves [3, 4], impurity pinning of flux-line in type-II superconductors [5], contact lines [6] as well as in eld driven ferromagnets, where the interface separates regions of opposite magnetizations [7].

A well established model to investigate the depinning transition in disordered ferromagnets is the driven random – eld Ising model (RFIM) (see for instance [7, 8, 9, 10, 11]). Here, the disorder induces some effective energy barriers which suppress the interface motion. A magnetic driving eld H reduces these energy barriers but they vanish only if the driving eld exceeds the critical value H_c . The transition from the pinned to the moving interface can be described as a continuous phase transition and its velocity v is interpreted as the order parameter. Without thermal fluctuations ($T = 0$) the eld dependence of the velocity obeys the power-law behavior

$$v(h; T = 0) \propto h \quad (1)$$

for $h > 0$, where h denotes the reduced driving eld $h = H/H_c - 1$.

The depinning transition is destroyed in the presence of thermal fluctuations ($T > 0$) which may provide the energy needed to overcome local energy barriers. Although thermal fluctuations drive the system away from criticality the order parameter obeys certain scaling laws and for sufficiently low temperatures the order parameter can be described as a generalized homogenous function [12]

$$v(h; T) = v(l = h; T); \quad (2)$$

similar to usual equilibrium second order phase transitions. Setting $l = 1$ one recovers Eq. (1) for zero temperature. Choosing $T = 1$ one gets for the interface velocity at the critical eld H_c

$$v(h = 0; T) \propto T^{-1} \quad (3)$$

This power-law behavior was observed in two- and three-dimensional simulations of the driven RFIM [12, 13] as well as in charge density waves in computer simulations and mean-eld calculations [4, 14].

Furthermore, thermal fluctuations cause a creep motion of the interface for small driving elds ($H < H_c$) characterized by an Arrhenius like behavior of the velocity [15, 16]. Recently, this creep motion was observed in experiments considering magnetic domain wall motion in thin films composed of Co and Pt layers [17], in renormalization group calculations [18, 19] regarding the Edwards-Wilkinson equation with quenched disorder (QEW), as well as in numerical simulations of the RFIM [20].

In equilibrium physics a scaling ansatz according to Eq. (2) usually describes the order parameter as a function of its control parameter and of its conjugated eld. Although Eq. (2) can be applied to the depinning transition, i.e. the temperature is a relevant scaling eld, T is not conjugated to the order parameter. The conjugated eld would support the interface motion independent of its strength. But strong thermal fluctuations destroy the

[†]Electronic address: lars@thp.uni-duisburg.de

^{††}Electronic address: usadel@thp.uni-duisburg.de

[‡]Electronic address: sven@thp.uni-duisburg.de

interface instead to support the interface motion. Therefore, one has to interpret the value of the thermal exponent carefully. For instance, it is not clear whether the obtained values of ν are a characteristic feature of the whole universality class of the depinning transition or just a characteristic feature of the particular considered RFIM. This point could be important for the interpretation of experiments which naturally take place at finite temperatures.

In this paper we reinvestigate the interface dynamics of the driven RFIM and focus our attention to higher dimensions $d \geq 3$. In particular we consider the scaling behavior at the critical point and determine the exponents ν and β . Our results suggest that the so-called upper critical dimension of the depinning transition of the driven RFIM is $d_c = 5$. Above this dimension the scaling behavior is characterized by the mean-field exponents. We compare our results with those obtained from a renormalization group approach of the quenched Edward-Wilkinson equation which is expected to be in the same universality class as the driven RFIM. A summary is given at the end.

II. MODEL AND SIMULATIONS

We consider the depinning transition RFIM on cubic lattices of linear size L in higher dimensions ($d \geq 3$). The Hamiltonian of the RFIM is given by

$$H = \frac{J}{2} \sum_{\langle i,j \rangle} S_i S_j - H \sum_i S_i - \sum_i h_i S_i, \quad (4)$$

where the first term characterizes the exchange interaction of neighboring spins ($S_i = \pm 1$). The sum is taken over all pairs of neighbored spins. The spins are coupled to a homogeneous driving field H as well as to a quenched random field h_i with $\langle h_i \rangle = 0$ and $\langle h_i h_j \rangle = \delta_{ij}$. The random field is assumed to be uniformly distributed, i.e., the probability p that the random field at site i takes some value h_i is given by

$$p(h_i) = \begin{cases} (2/\pi)^{1/2} & \text{for } |h_i| < \pi/2 \\ 0 & \text{otherwise.} \end{cases} \quad (5)$$

Using antiperiodic boundary conditions an interface is induced into the system which can be driven by the field H (see [12] for details). A Glauber dynamics with random sequential update and heat-bath transition probabilities is applied to simulate the interface motion (see for instance [21]).

Due to this algorithm not only spins adjacent to the interface but throughout the whole system can flip with a finite probability at temperatures $T > 0$. In general, this may cause nucleation which always starts with an isolated spin-flip. The minimum energy required for an isolated spin-flip is $E = 2(zJ - H)$ with $z = 2^d$ nearest neighbors on a bcc lattice. Although the corresponding spin-flip probability $1 = [1 + \exp(-E/T)]$ is small

for the considered temperatures ($T = 0.2$), isolated spin-flips are possible and occur. But we observed that these spin-flips are unstable in our simulations, i.e., an isolated spin will flip back in the next update step. Thus the originally induced interfaces are stable during the simulations.

The moving interface corresponds to a magnetization M which increases in time t (given in Monte Carlo step per spin). The interface velocity, which is the basic quantity in our investigations, is obtained from the time dependence of the magnetization $v = dM/dt$ where $\langle \cdot \rangle$ denotes an appropriate disorder average. Starting with a flat interface we performed a sufficiently number of updates until the system reaches after a transient regime the steady state which is characterized by a constant average interface velocity.

As pointed out in previous works [12, 13] an appropriate choice of the interface orientation is needed in order to recover that the interface moves for arbitrarily small driving field in the absence of disorder. An appropriate choice is to consider the interface motion along the diagonal direction of a simple cubic lattice. For $d \geq 3$ an alternative is to examine the interface motion along the z -axis on a body-centered cubic (bcc) lattice. Since it is much more convenient to implement the latter case in higher dimensions we consider in this work bcc lattices, the more as the lattice structure usually does not affect the universal scaling behavior.

III. $D = 3$

In the case of the three dimensional RFIM we consider bcc lattices of linear size $L = 250$. A snapshot of a moving interface in the steady state is presented in Fig. 1. The obtained values of the interface velocities for $T = 0$




FIG. 1: Snapshot of a moving interface at $T = 0$ for $L = 128$, $\beta = 1.7$, and $H = 1.37$ in $d = 3$. In order to show the details we stretched the interface in the vertical direction by a factor 15.




FIG. 2: Dependence of the interface velocity v on the driving field H for a bcc and simple cubic (sc) lattice, respectively. The inset shows v as a function of the reduced driving field h . The dash-dotted lines are fits according to Eq. (1).

are plotted in Fig. 2. As one can see v tends to zero in the vicinity of $H_c = 1.36$. Assuming that the scaling behavior of the interface motion is given by Eq. (1) one varies H_c until one gets a straight line in a log-log plot. Convincing results are obtained for $H_c = 1.357 \pm 0.001$ and the corresponding curve is shown in the inset of Fig. 2. For lower and greater values of H_c we observe significant curvatures in the log-log plot (not shown). In this way we estimate the error-bars in the determination of the

critical field. A regression analysis yields the value of the order parameter exponent $\beta = 0.653 \pm 0.026$. This value agrees with $\beta = 0.66 \pm 0.04$ which was obtained from a similar investigation [12] where the interface moves along the diagonal direction of a simple cubic lattice (see inset of Fig. 2). Furthermore, both results are in agreement with $\beta = 0.60 \pm 0.11$ [10, 11], where in $d = 3$ the influence of helicoidal boundary conditions in one direction and periodic ones in the other direction parallel to the interface was investigated on a simple cubic lattice.

In order to determine the exponent ν we simulated the RFIM around the critical field for $T = 0.025n$ with $n = 2, 3, 4, 6, 8$. The obtained curves are shown in the inset of Fig. 3. According to Eq. (2) the interface velocity scales as

$$v(h; T) = T^{1-\nu} v(h T^{1-\nu}; 1) \quad (6)$$

Plotting $v(h; T) T^{1-\nu}$ as a function of $h T^{1-\nu}$ one varies ν , β , and H_c until one gets a data collapse of the different curves. Convincing data collapses are observed for $\beta = 0.63 \pm 0.06$, $\nu = 2.33 \pm 0.2$ and $H_c = 1.360 \pm 0.01$ and the corresponding curves are shown in Fig. 3. The obtained values of the order parameter exponent and of the critical field agree within the error-bars with the values of the $T = 0$ analysis. Furthermore our results are in agreement with similar investigations on a simple cubic lattice ($\beta = 0.63 \pm 0.07$ and $\nu = 2.38 \pm 0.2$, see [12]).

IV. $D = 4$

In order to determine the order parameter exponent of the four-dimensional driven RFIM we measured the




FIG. 3: Scaling plot of the interface velocity for $d = 3$. The data are rescaled according to Eq. (6). The inset shows the unscaled velocities for $T = 0.025n$ with $n = 2, 3, 4, 6, 8$ (solid lines) in comparison to the $T = 0$ data from Fig. 2 (dashed line).




FIG. 4: The interface velocity of the four-dimensional model at $T = 0$. For sufficiently small fields the data obey a power law according to Eq. (1) (dotted dashed line). For the two we use only those data marked by filled symbols and we find $H_c = 1.258 \pm 0.05$ and $\beta = 0.8 \pm 0.06$.




FIG. 5: Scaling plot of the interface velocity for $d = 4$ and $d = 6$, respectively (see inset). The data are rescaled according to Eq. (6).

interface velocity for bcc lattices of linear sizes $L = 140$. The obtained data for $T = 0$ are shown in a log-log plot in Fig. 4. After a transient regime which displays a finite curvature we observe an asymptotic power-law behavior for sufficiently small driving field H . A regression analysis yields $\nu = 0.8 \pm 0.06$ and $H_c = 1.258 \pm 0.002$.

To determine the exponent ν we simulated the RFIM in the vicinity of the critical field for $T = 0.025n$ where again $n = 2, 3, 4, 6, 8$ was chosen. Similar to the three-dimensional case one varies the exponents as well as the critical field until one observes a data collapse. Good results are obtained for $\nu = 0.73 \pm 0.13$, $\beta = 1.72 \pm 0.27$, and $H_c = 1.256 \pm 0.015$. The corresponding data collapse is shown in Fig. 5. Again, the obtained values of ν and H_c confirm the above presented analysis for $T = 0$.

$$\nu = D = 5$$

In the case of the d -dimensional RFIM system sizes from $L = 10$ up to $L = 30$ are simulated. Analyzing the interface motion at $T = 0$ we observe that the velocity-field dependence can not be described by a pure power-law, i.e., Eq. (1) fails. In Fig. 6 we plotted the logarithmic derivative of the velocity-field dependence

$$\beta_{\text{eff}} = \frac{\partial \ln v}{\partial \ln h} \quad (7)$$

which can be interpreted as an effective exponent. If the asymptotic scaling behavior obeys Eq. (1) the logarithmic derivative tends to the value of ν for $H \rightarrow H_c$. But as can be seen from Fig. 6 no clear saturation takes place for $d = 5$ as it is observed for the three and four-dimensional model. The lack of a clear saturation could be explained




FIG. 6: The effective exponents β_{eff} as a function of $\ln h$ for various dimensions. The figure shows that the d -dimensional exponent does not display a clear saturation as the exponent of the lower dimensions do. The data of the six-dimensional model indicate that the expected saturation value $\beta_{\text{eff}} = 1$ is reached significantly faster as compared to $d = 5$ (see text). The values of the two-dimensional system are obtained from [13].

by a too large distance from the critical point, but another reason is possible too.

Significant deviations from a pure power law behavior [Eq. (1)] occur for instance at the upper critical dimension d_c where the scaling behavior is governed by the mean-field exponents modified by logarithmic corrections. The scaling behavior around d_c is well understood within the renormalization group theory (see for instance [22, 23, 24]). For $d > d_c$ the stable fixed point of the corresponding renormalization equations is usually a trivial fixed point with classical mean-field exponents. This trivial fixed point is unstable for $d < d_c$ and a different stable fixed point exists with nonclassical exponents. These exponents can be estimated by an ϵ -expansion, for instance. For $d = d_c$ both fixed points are identical and marginally stable. In this case the asymptotic form of the thermodynamic functions is given by the mean-field power-law behavior modified by logarithmic corrections. Applying this approach to the depinning transition, the corresponding ansatz reads

$$v(h; T = 0) = h^{M_F} j \ln h^{\beta}; \quad (8)$$

where B denotes an unknown correction exponent. It is worth to mention that in contrast to the values of the critical exponents below the upper critical dimension the above scaling behavior does not rely on approximation schemes like ϵ - or $1/n$ -expansions [25]. Within the renormalization group theory it is an exact result in the limit $h \rightarrow 0$ (see e.g. [22, 26] and references therein for RG investigations and [27, 28] for measurements).




FIG. 7: The rescaled interface velocity as a function of the driving field for $d = 5$. In order to display the logarithmic corrections we plot $(v/h)^{1/B}$ vs. $\ln h$ [see Eq. (8)]. The solid line corresponds to the expected asymptotic scaling behavior for $h \rightarrow 0$ [corresponding to $\ln h \rightarrow (-\infty)$] according to Eq. (9).

The value $\nu_{MF} = 1$ is reported for depinning transitions [1, 29, 30]. Thus we analyze $v(h)h$ as a function of $\ln h$ and note again that Eq. (8) describes only the leading order of the scaling behavior, i.e., we expect that the asymptotic behavior of interface velocity obeys

$$[v(h)h]^{1/B} = \text{const } \ln h; \quad (9)$$

Therefore, we varied in our analysis the logarithmic cor-




FIG. 8: The scaling plot of the interface velocity v for the six dimensional model, i.e. at the upper critical dimension. The data are rescaled according to Eqs. (10,11) using $\nu_{MF} = 1.49$, $\nu = 1$, $\sigma = 0.22$, and $\tau = 0.19$.

rection exponent B and the critical field H_c until we get this expected asymptotic behavior. The best results are obtained for $B = 0.40 \pm 0.09$ and $H_c = 1.14235 \pm 0.001$ and the corresponding scaling plot is shown in Fig. 7. The observed asymptotic agreement with Eq. (9) corresponds to a logarithmically ($\propto \ln h$) convergence of ν to $\nu_{MF} = 1$, which explains why no clear saturation of the effective exponent could be observed for $h \rightarrow 0$ in the six dimensional model.

Similar to the $T = 0$ scaling behavior one has to modify for $T > 0$ the scaling ansatz since no data collapse could be obtained by plotting the data according to Eq. (2). Motivated by recently performed investigations of the scaling behavior of an absorbing phase transition around the upper critical dimension [31] we assume that the scaling behavior of the order parameter obeys in leading order

$$v(h; T) = T^{1-\nu_{MF}} \ln T \ln v(x; 1) \quad (10)$$

where the scaling argument x is given in leading order by

$$x = h T^{1-\nu_{MF}} \ln T \ln j \quad (11)$$

with $\nu_{MF} = 1$. In our analysis we use the value $\nu_{MF} = 1.49$ obtained from the analysis of the six dimensional RFIM (see next section). Therefore, we have to vary the exponents and in order to observe a data collapse according to Eqs. (10,11). Convincing results are obtained for $\sigma = 0.22 \pm 0.16$, $\tau = 0.19 \pm 0.12$. The corresponding data collapse is shown in Fig. 8.

$$VI. D = 6$$

Above the upper critical dimension the scaling behavior is characterized by the mean-field exponents, i.e., in leading order the interface velocity is given by

$$v(h; T = 0) \propto h; \quad (12)$$

In Fig. 9 we plot the velocity as a function of the driving field H obtained from simulations of system sizes $L \leq 14$. As one can see the velocity does not display the expected linear behavior. It seems that a linear behavior is only given for small velocities, i.e., our data do not display the pure asymptotic behavior [Eq. (12)]. This is confirmed by the behavior of the effective exponent [Eq. (7)] which increases fast for $h \rightarrow 0$ but the actual saturation to $\nu = 1$ does not take place for the considered values of h (see Fig. 6). To observe the asymptotic behavior one has to perform simulations closer to the critical point H_c which requires to simulate larger system sizes. Unfortunately, the limited CPU power makes this impossible.

An alternative is to take the curvature of the function $v(h)$ into consideration and to assume that the leading corrections to the asymptotic behavior are of the form

$$v(h; T = 0) = v_1 h + v_2 h^2 + O(h^3); \quad (13)$$




FIG. 9: The interface velocity as a function of the driving field for the six-dimensional model. The dashed line corresponds to a v according to Eq. (13). The inset displays $v(h)/h$ as a function of the reduced driving field h . The resulting linear behavior confirms that the deviations from the mean-field behavior can be described by quadratic corrections.

which recovers Eq. (12) for $h \rightarrow 0$. Fitting our data to this ansatz we get $H_c = 1.1537 \pm 0.003$, $v_1 = 0.2546$, and $v_2 = 0.0895$. The corresponding curve fits the simulation data quite well as one can see from Fig. 9. In the inset of Fig. 9 we plotted $v(h)/h$ as a function of the reduced driving field h . According to the above ansatz [Eq. (13)] one gets a linear behavior, i.e., the deviations from the pure mean-field behavior [Eq. (12)] can really be described as quadratic corrections. Thus we get that our numerical data are consistent with the assumption that the six-dimensional RFIM depinning transition is characterized by the mean-field exponent $\nu_{MF} = 1$.

Again we consider how thermal fluctuations affect the scaling behavior and analyse interface velocities obtained at different temperatures $T = 0.025n$ with $n = 2, 3, 4, 6, 8$. Similar to the situation below the upper critical dimension we assume that the scaling behavior of the interface velocity is given by Eq. (2) where the exponents are given by mean-field values. A convincing data collapse is obtained for $\nu_{MF} = 1.49 \pm 0.15$ and $H_c = 1.153 \pm 0.02$ and is plotted in the inset of Fig. 5.

VII. DISCUSSION

A well established realization of interface pinning in a disordered media is the so-called quenched Edwards-Wilkinson (QEW) equation of motion which was intensively investigated in the last decade [7, 18, 19, 29, 30, 32]. It is argued that the QEW equation as well as the driven RFIM are characterized by the same critical exponents, i.e., both models belong to the same univer-




FIG. 10: The critical exponents β and ν as a function of the dimension. The solid line corresponds to the values of an ϵ -expansion [Eq. (14)].

sality class [7, 10]. Renormalization group analyses of the quenched QEW equation [29, 30] predict, in accordance with [7], $d_c = 5$ and allow to estimate the critical exponents using an ϵ -expansion. A recently performed two-loop renormalization approach yields [30]

$$\nu_{QEW} = 1 - \frac{1}{9} \epsilon^2 + O(\epsilon^3) \quad (14)$$

where ϵ denotes the distance from the upper critical dimension, i.e. $\epsilon = 5 - d$ (unfortunately, no error-bars can be estimated from an ϵ -expansion). The corresponding values of the exponents as a function of the dimension are plotted in Fig. 10. The numerically determined exponents of the driven RFIM (listed in Table I) are in a fair agreement with the values of the ϵ -expansion.

For the QEW equation the temperature exponent is not known. Therefore, a direct comparison with the obtained values of the driven RFIM is not possible.

TABLE I: The exponents (obtained from simulations at $T = 0$ and $T > 0$, respectively) and ν of the depinning transition of the RFIM for different dimensions. The values of the two-dimensional model are obtained from [13]. The critical behavior at the upper critical dimension d_c is additionally affected by logarithmic corrections.

d	$T = 0$		$T > 0$			
2	0.35	0.04	0.33	0.02	5.00	0.3
3	0.653	0.026	0.63	0.06	2.33	0.2
4	0.80	0.06	0.73	0.13	1.72	0.27
$d_c = 5$	1		1		1.49	
6	1		1		1.49	0.15

V III. C O N C L U S I O N S

We studied numerically a field driven interface in the RFIM and determined the order parameter exponent as well as the temperature exponent. Below the upper critical dimension $d_c = 5$ the critical exponents depend on the dimension and the values of the exponents correspond to those of a two-loop renormalization group approach of the Edwards-Wilkinson equation [30]. This suggests that the depinning transition of the RFIM model belongs to the universality class of the quenched Edwards-Wilkinson equation. At the upper critical dimension $d_c = 5$ the scaling behavior is affected by logarithmic corrections. Above the upper critical dimension we ob-

serve that the scaling behavior is characterized in leading order by the corresponding mean-field exponents.

A c k n o w l e d g m e n t s

This work was supported by the Deutsche Forschungsgemeinschaft via GRK 277 Struktur und Dynamik heterogener Systeme (University of Duisburg) and SFB 491 Magnetische Heteroschichten: Struktur und elektronischer Transport (Duisburg/Bochum).

S.Lubeck wishes to thank the Minerva Foundation (Max Planck Gesellschaft) for financial support.

-
- [1] M. Kardar, Phys. Rep. 301, 85 (1998).
 - [2] H. Hinrichsen, Adv. Phys. 49, 815 (2000).
 - [3] D. S. Fisher, Phys. Rev. Lett. 50, 1486 (1983).
 - [4] D. S. Fisher, Phys. Rev. B 31, 1396 (1985).
 - [5] G. Blatter, M. V. F. and V. B. Geshkenbein, A. I. Larkin, and V. M. Vinokur, Rev. Mod. Phys. 66, 1125 (1994).
 - [6] P. G. de Gennes, Rev. Mod. Phys. 57, 827 (1985).
 - [7] R. Bruinsma and G. Aeppli, Phys. Rev. Lett. 52, 1547 (1984).
 - [8] H. Ji and M. O. Robbins, Phys. Rev. A 44, 2538 (1991).
 - [9] B. Drossel and K. Dahmen, Eur. Phys. J. B 3, 485 (1998).
 - [10] L. A. N. Amaral, A.-L. Barabasi, and H. E. Stanley, Phys. Rev. Lett. 73, 62 (1994).
 - [11] L. A. N. Amaral et al., Phys. Rev. E 52, 4087 (1995).
 - [12] L. Roters et al., Phys. Rev. E 60, 5202 (1999).
 - [13] U. Nowak and K. D. Usadel, Europhys. Lett. 44, 634 (1998).
 - [14] A. A. Middleton, Phys. Rev. B 45, 9465 (1992).
 - [15] L. B. Ioé and V. M. Vinokur, J. Phys. C 20, 6149 (1987).
 - [16] T. Nattermann, Europhys. Lett. 4, 1241 (1987).
 - [17] S. Lemelle et al., Phys. Rev. Lett. 80, 849 (1998).
 - [18] P. Chauve, T. Giamarchi, and P. L. Doussal, Europhys. Lett. 44, 110 (1998).
 - [19] P. Chauve, T. Giamarchi, and P. L. Doussal, Phys. Rev. B 62, 6241 (2000).
 - [20] L. Roters, S. Lubeck, and K. D. Usadel, Phys. Rev. E 63, 026113 (2001).
 - [21] K. Binder and D. W. Heermann, Monte Carlo Simulation in Statistical Physics (Springer, Berlin, 1997).
 - [22] F. J. Wegner, in Phase Transitions and Critical Phenomena, Vol. 6, edited by C. Domb and M. S. Green (Academic Press, London, 1976).
 - [23] E. Brezin, J. C. Le Guillou, and J. Zinn-Justin, in Phase Transitions and Critical Phenomena, Vol. 6, edited by C. Domb and M. S. Green (Academic Press, London, 1976).
 - [24] P. Pfeuty and G. Toulouse, Introduction to the renormalization group and critical phenomena (John Wiley & Sons, Chichester, 1994).
 - [25] E. Brezin and J. Zinn-Justin, Phys. Rev. B 13, 251 (1976).
 - [26] F. J. Wegner and E. K. Riedel, Phys. Rev. B 7, 248 (1973).
 - [27] J. A. Gri n, J. D. Lister, and A. Linz, Phys. Rev. Lett. 38, 251 (1977).
 - [28] J. Brinkmann, R. Courths, and H. J. Guggenheim, Phys. Rev. Lett. 40, 1286 (1978).
 - [29] T. Nattermann, S. Stepanow, L.-H. Tang, and H. Leschhorn, J. Phys. II 2, 1483 (1992).
 - [30] P. Chauve, P. L. Doussal, and K. J. Wiese, Phys. Rev. Lett. 86, 1785 (2001).
 - [31] S. Lubeck, Phys. Rev. E 65, 046150 (2002).
 - [32] O. Narayn and D. S. Fisher, Phys. Rev. B 48, 7030 (1993).