

Composite spin and orbital triplet superconductivity

Frithjof B. Anders

Institut für Festkörperphysik, Darmstadt University of Technology,
Hochschulstr. 8, 64289 Darmstadt, Germany

We show that the two-channel Anderson lattice model leads to the development of an unconventional superconductivity out of a metallic non Fermi-liquid phase. It is characterized by a composite order parameter comprising of a local spin or orbital degree of freedom bound to triplet Cooper pairs with an isotropic and a nearest neighbour form factor. The gap function is non analytic and odd in frequency, and a pseudo-gap develops in the conduction electron density of states which vanishes as $\omega \rightarrow 0$.

PACS numbers: 75.20.Hr, 75.30.Mb, 71.27.+a

1. Introduction

Heavy Fermion [1] (HF) superconductivity (SC) has drawn much attention since the discovery of superconductivity in CeCu_2Si_2 [2] which is likely characterized by an anisotropic order parameter with symmetry yet to be determined. It became apparent over the last decade that almost all HF materials are unstable with respect to magnetic or superconducting phase transitions, which either compete with each other or can even coexist as found in uranium based materials. Moreover, superconductivity in URu_2Si_2 develops out of an incoherent metallic phase with a high resistance at T_c of 100 K. The two-channel Anderson lattice model,

$$\hat{H} = \sum_{\mathbf{k}} \sum_{\alpha} \epsilon_{\mathbf{k}}^{\alpha} c_{\mathbf{k}\alpha}^{\dagger} c_{\mathbf{k}\alpha} + \sum_i E_i \sum_{\alpha} c_{i\alpha}^{\dagger} c_{i\alpha} + \sum_{i,j} \sum_{\alpha} V_{ij}^{\alpha} c_{i\alpha}^{\dagger} c_{j\alpha} + \text{h.c.} \quad (1)$$

is believed to serve as a possible description of the electronic properties of URu_2Si_2 or Pr^{3+} compounds with cubic symmetry. $\epsilon_{\mathbf{k}}$ is the band dispersion of

Presented at the Strongly Correlated Electron Systems Conference, Krakow 2002

the two conduction bands coupling to the localized doublets via hybridization $V \cdot X^{(i)}$; are the usual Hubbard operators describing local f -states at each lattice site i . One doublet is associated with a quadrupolar charge distribution ($X_{\pm}^{(i)}$) and one carries a spin ($X_{\pm}^{(i)}$). Its paramagnetic phase is characterized by a large residual resistivity and entropy, and ill-defined electronic quasi-particles. Fermi liquid physics is restored by cooperative ordering or applied magnetic field [3].

2. Composite order parameter

It was observed that the local two-particle-vertex irreducible vertex has a $1/\omega$ singularity and is odd with respect to the incoming and outgoing frequencies [4]. This is related to the fact that correlations are induced through consecutive hybridization processes on the f -shell due to the strong on-site Coulomb repulsion. Therefore, odd-frequency superconductivity [5] is favoured in this model. Assuming an order parameter with g_1 symmetry, the order parameter must have spin singlet, channel singlet (SSC) symmetry or spin triplet, channel triplet (STC) symmetry. Only for the latter case, a stable solution was found. The order parameter in the STC sector

$$O_{ij} = \text{sign}(E_h - E) \hbar s_i P_j^{st} i - \hbar s_j P_i^{ts} i; \quad (2)$$

correlates a local f -spin (quadrupolar moment) $s_i (i) - i; j = x; y; z$ - component with a itinerate Cooper pair $P^{st} (P^{ts})$ in the SSC (STC) sector:

$$P^{st} = \frac{1}{N} \sum_{\mathbf{k}} S(\mathbf{k})^T (\mathbf{k})_{i=y} i_{y=y} \sim (\mathbf{k}) \quad (3)$$

$$P^{ts} = \frac{1}{N} \sum_{\mathbf{k}} S(\mathbf{k})^T (\mathbf{k})_{i=y} i_{y=y} \sim (\mathbf{k}) : \quad (4)$$

The form factor $S(\mathbf{k})$ transforms as g_1 , and $\sim(\mathbf{k})$ is a bi-spinor in spin and channel space. $O_{i,j}$ is invariant with respect to exchanging the orbital and the magnetic doublet $j i$ and $j i$ and a simultaneous particle-hole transformation of the conduction electrons. Hence, the occurrence of superconductivity is independent of the character of the local ground state of the local ion. We investigate the superconducting phase within the dynamical mean field theory (DMFT) by self-consistently calculating the anomalous self-energy under the presence of a Cooper-pair field.

3. DMFT in the superconduction phase

The Nambu Green function is an 8×8 matrix in spin and orbital space. Assuming no directional coupling between spin and orbital degrees

of freedom, the anomalous 4×4 self-energy matrix may be written as $g(z; \mathbf{k})_{\mathbf{s}\mathbf{s}'} = \mathbf{n}_{\mathbf{s}} \mathbf{n}_{\mathbf{s}'}$ where $\mathbf{n}_{\mathbf{s}}$ and $\mathbf{n}_{\mathbf{s}'}$ are constant unit vectors in spin and channel space, and the amplitude function $g(z; \mathbf{k})$ has to be odd in frequency. This reduces the full problem to a standard 2×2 size for $g(z; \mathbf{k})$ and the diagonal self-energy. In order to derive DMFT equations with a purely local self-energy matrix $\underline{\Sigma}_c(z)$, the anomalous self-energy is restricted to isotropic pairs, e. g. $g(z; \mathbf{k}) = g(z) \cdot \underline{\underline{G}}_c(z)$ denotes the medium matrix in which the effective impurity is embedded. It is related to a generalized Anderson width matrix through $\underline{\underline{G}}_c(z) = V^2 \underline{\underline{G}}_c(z)$ and has normal and anomalous components describing quasi-particle propagation and pair-creation and annihilation, respectively. The band electron self-energy $\underline{\Sigma}_c(z) = \underline{\underline{T}} \underline{\underline{1}} + \underline{\underline{G}} \underline{\underline{T}}$ is determined by the local T-matrix $\underline{\underline{T}}$ whose diagonal elements are given by the local quasi-particle scattering matrices $\underline{\underline{T}}_{11}(z) = V^2 G_f(z)$, and its anomalous contribution stems from Cooper pairs scattered on the local shells.

As an impurity solver for effective site problem of the DMFT, the NCA [6] was extended to the superconducting phase. The additional terms (z) to the self-consistency equation for the local ionic propagators are generated by the anomalous media in the SC phase.

$$(z) = \sum_{\mathbf{k}} \int_{-1}^1 \int_{-1}^1 d\omega d\omega' f(\omega) f(\omega') (5)$$

$$(z) = \sum_{\mathbf{k}} \int_{-1}^1 \int_{-1}^1 d\omega d\omega' f(\omega) f(\omega') (6)$$

where $f(\omega) = \text{Im} \underline{\underline{G}}_{12}(i\omega)$ is the spectral function of the anomalous media. The anomalous local T-matrix is also obtained by the same additional diagram of the generating functional and reads

$$T_{12}(i\omega_n) = \sum_{\mathbf{k}} \int_{-1}^1 d\omega f(\omega) f(\omega') \frac{e^{-\omega}}{2iZ_f} P(z) (7)$$

where Z_f is the partition function of the effective site, and the contour C encircles all singularities of the integrand kernel. These coupled double integral equations are numerically solved using similar techniques as developed for the Post-NCA equations [7].

We calculated the self-consistent DMFT solution of the spectral functions close to T_c which are displayed in Fig. 1. The linear frequency dependency of the quasi-particle and anomalous Green function very close to




Fig.1. Spectral functions of the quasi-particle and the anomalous Green function in DMFT (NCA) in the superconducting phase for $1/T=T_c-1$ in the vicinity of $T_c=0$. The dotted curve shows a fit with a $\propto |\omega|$. Parameters: $\rho(\omega) = \exp(-[(\omega - W)^2]^{1/2})$ conduction band DOS, band width $W = 10$, $E_c = 2$, $E_v = V^2$ (ω),

$T_c = 0$, similar to d-wave superconductivity, is clearly visible. The opening of a pseudo-gap indicates a decrease of the energy in the superconducting phase, and, hence, its thermodynamical stability. Details of the filling dependence and the phase diagram are beyond the scope of this paper and will be published elsewhere [4].

REFERENCES

- [1] N. Grewe and F. Steglich, in Handbook on the Physics and Chemistry of Rare Earths, edited by J. K. A. G. Schneider and L. Eyring (North-Holland, Amsterdam, 1991), Vol. 14, p. 343, for a review on Heavy Fermion compounds.
- [2] F. Steglich, Phys. Rev. Lett. 43, 1892, 1979.
- [3] F. B. Anders, M. Jarrell, D. L. Cox, Phys. Rev. Lett. 78, 2000, 1997.
- [4] Details will be given in F. B. Anders, cond-mat/0204469 and Euro. Phys. J. B, 2002
- [5] R. Heid, Z. Phys. B 99, 15 (1995) and references therein.
- [6] N. Grewe, Z. Phys. B 52, 193 (1983),
- [7] F. B. Anders, J. Phys.: Condens. Matter 7, 2801 (1995)