The importance of friction in the description of low-temperature dephasing

by Florian Marquardt

Department of Physics and Astronomy, University of Basel, Klingelbergstrasse 82, CH-4056 Basel, Switzerland

(Dated: July 30th, 2002)

We discuss the importance of the real part of the Feynman-Vernon influence action for the analysis of dephasing and decay near the ground state of a system which is coupled to a bath. Using exactly solvable linear quantum dissipative systems, it is shown how the effects of the real and the imaginary part (describing friction and fluctuations, respectively) may cancel beyond lowestorder perturbation theory. The resulting picture is extended to a qualitative discussion of nonlinear systems and dephasing of degenerate fermions. We explain why dephasing rates will, in general, come out finite at zero temperature if they are deduced from the imaginary part of the action alone, a procedure which is reliable only for highly excited states.

I. INTRODUCTION

The Feynman-Vernon influence functional^{1,2} plays a prominent role in discussions of dephasing that aim to go beyond a simple master equation approach. In principle, the influence functional takes into account every effect of the environment (the "bath") on the system under consideration. This includes heating, dephasing, friction and renormalization effects (changing the external potential or the effective mass of a particle). Its popularity arises not only from the fact that it constitutes an exact approach, but also from the direct physical meaning which it acquires in some situations. In particular, this holds for typical interference experiments, where two wave packets describing a single particle follow two different trajectories in order to be recombined later on. For such a case, the influence functional is equal to the overlap between the two different bath states which result due to the particle moving along either one of the semiclassical trajectories. If the bath can distinguish between the two paths, it acts as a which-way detector, and the diminished magnitude of the overlap directly gives the ensuing decrease of the visibility of the interference pattern. Writing the overlap in the form of an exponential, S_T), one can obviously conclude that only the exp (iS_R imaginary part S_{I} of the "influence action" is responsible for this decrease and, therefore, S_{I} alone describes dephasing in such a situation. Furthermore, if the system is subject to a fluctuating external *classical* field, only S_{T} remains, while S_R vanishes identically. This is because S_I is due to the fluctuations of the bath, while S_R stems from the back-action of the bath onto the system (including friction effects), which is absent for an external noise field. The approach of evaluating S_{I} along semiclassical trajectories has been successfully applied to the calculation of dephasing rates in many situations 3,4,5 .

However, near the ground state of a system, an analysis along these lines is likely to fail. Qualitatively, this may already be deduced from the fact that dropping S_R means replacing the environment by an artificial fluctuating classical field whose correlation function includes the zero-point fluctuations of the original quantum bath. Therefore, even at zero temperature, this fluctuating field will, in general, heat the system, which directly leads to dephasing. The role of S_R is to counteract this effect. In this article, it is our aim to display explicitly, in a detailed manner, the necessity of keeping the real part $S_{\rm R}\,$ of the influence action in discussions of dephasing and decay near the ground state of a system.

This issue derives its importance partly from the fact that the evaluation of S_I along semiclassical trajectories has proven to be an efficient way of extracting dephasing rates in the problem of weak localization 3,4 in the limit of high temperatures, when zero-point fluctuations of the bath may be omitted. In contrast, at low temperatures, a single-particle semiclassical calculation may become invalid, since it neglects the Pauli principle which is known to play an important role for the inelastic scattering of electrons and is not included in the Feynman-Vernon influence functional. However, recently an extension of the influence functional to the case of a many-fermion system has been derived, using an exact procedure^{6,7,8} and including the Pauli principle. This permitted a discussion of dephasing in a disordered metal even for the case of low temperatures. Following the general strategy of earlier works⁹ dealing with the high-temperature case, the dephasing rate was deduced from S_T in a semiclassical calculation and found to be finite even at zero temperature. Since the "orthodox" theory³ had predicted a vanishing rate in the limit T ! 0, the new results prompted considerable criticism^{10,11,12,13}, which mostly emphasized technical aspects of impurity averaging or used perturbation theory to arrive at different conclusions. In the present work, we want to clarify some essential aspects of the roles of S_R and S_I , using physically much more transparent exactly solvable models.

We want to emphasize that "zero-temperature dephasing" as such is perfectly possible: If one prepares a system in any superposition of excited states and couples it to a bath, it will, in general, decay towards its ground state by spontaneous emission of energy into the bath (at T = 0). This will destroy any coherent superposition, thus leading to dephasing. Considerations of this kind are particularly relevant for quantum-information processing, where one necessarily deals with nonequilibrium situations involving excited qubit states of finite energy. The situation is different for the weak-localization problem (and similar transport interference effects): There, one is interested in the zero-frequency limit of the system's linear response, which depends on the coherence properties of arbitrarily low-lying excited states. It is the subtleties associated with a pathintegral description of these situations which we want to address in this work.

The article is organized as follows: After a brief review of the influence functional and its meaning in semiclassical situations (sec. II), we will rewrite the expressions for S_{T} and the Golden Rule decay rate, in order to compare the two (sec. III). In doing so, we will closely follow the analysis of Cohen and $Imry^{5,12}$. Then, we will show how and why the effects of S_R and S_I may compensate each other in the derivation of a decay rate starting from the path integral expression for the time-evolution of the density matrix (sec. IV), even though they cannot cancel each other in the influence action. There, it may be observed that drawing conclusions about decay directly from the ex- S_T of the influence functional is usually not ponent iS_R possible. The crucial cancellation takes place at a later stage of the calculation, after proper integration over the fluctuations around the classical paths and after averaging over the initial state. In order to prove that this compensation takes place not only in lowest order perturbation theory, we will then specialize to exactly solvable linear dissipative systems, i.e. the damped harmonic oscillator (sec. V) and the free particle (sec. VI). We will also point out that there is an important difference between the oscillator and the free particle: Dropping S_{R} has a much more drastic effect on the former, leading to an artificial finite decay rate of the ground state at zero temperature. However, in order to discuss the importance of S_R for the calculation of dephasing rates (involving decay of excited states), we have to extend our analysis to nonlinear models (sec. VII). We will explain in a more qualitative fashion why the essential insights gained from the exactly solvable models should remain valid both for the nonlinear models as well as for systems of degenerate fermions.

II. THE INFLUENCE FUNCTIONAL AND DEPHASING IN SIMPLE SITUATIONS

We are interested in the time-evolution of the reduced density matrix of a system with coordinate q which is coupled to some environment (the bath). If system and bath had been uncoupled prior to t = 0, then the density matrix at a later time t is linearly related to that at time 0:

$$(q_{1}^{2};q_{2}^{2};t) = dq_{0}^{2} dq_{0}^{2} J(q_{1}^{2};q_{2}^{2};q_{0}^{2};q_{0}^{2};t) (q_{1}^{2};q_{0}^{2};0):$$
(1)

The propagator J on the right-hand-side of this equation is given by

$$J = D q^{>} D q^{<} \exp i(S_0^{>} S_0^{<}) \exp [iS_R S_I]: (2)$$

Here we have set h 1 and introduced an abbreviated notation: The path integral extends over all "forward" paths $q^{>}$ () running from the given value of ${}_{0}^{>}q$ to $q_{L}^{>}$, likewise for the "backward" paths $q^{<}$ (). The value of the action of the uncoupled system, evaluated along $q^{>}$ (<) (); is denoted by $S_{0}^{>}$ (<). The second exponential in Eq. (2) is the Feynman-Vernon influence functional^{1,2}. Both S_R and S_I are real-valued functionals that depend on both paths simultaneously. The influence functional is the overlap of bath states which have time-evolved out of the initial bath state $_0$ under the action of either q^2 () or q ():

$$e^{iS_{R} S_{I}}$$
 $[q^{<}()]j^{\dagger}q()]:$ (3)

At zero temperature, the state $_0$ is the ground state of the unperturbed bath, while at finite temperatures an additional thermal average over $_0$ has to be performed. From this representation, it follows^{1,2} that S_R changes sign on interchanging $q^{>}$ and $q^{<}$ while S_I always remains nonnegative - the magnitude of the overlap can only be decreased compared with its initial value of one.

The meaning of S_R and S_I becomes particularly transparent for an interference setup, where two wave packets > and < travel along two different paths $q_{c1}^{>}()$ and $q_{c1}^{<}()$. We want to assume a situation which can be described semiclassically, i.e. the wave-length is supposed to be much smaller than the size of a wave packet and this again is much smaller than the typical separation between the paths and the typical dimensions over which the external potential changes. Then, it suffices to evaluate S_R and S_I just for the combination of these two paths, since the fluctuations around them are comparatively unimportant. The environment will affect the interference pattern on the screen mainly by changing the interference term⁴:

(x;x;t)
$$j_{>}(x;t)j^{2} + j_{<}(x;t)j^{2} + 2Re_{>}(x;t) < (x;t)e^{iS_{R}} \cdot s_{\perp}$$
 (4)

In this equation, (x;t) are assumed to represent the unperturbed time-evolution of the wave packets. To a first approximation, $S_{R;I}$ do not enter the "classical" terms in the first line of Eq. (4), since $S_{R,I}[q_{cl}^{>};q_{cl}^{>}] = 0$. Deviations from this first approximation stem from the integration over fluctuations away from $q_{-1}^{> (<)}$ and describe, for example, mass- and potential renormalization as well as slowing-down of the wave packet due to friction. Obviously, $S_{I}[q_{1}^{2};q_{1}^{2}]$ determines directly the decrease in visibil-Sı ity of the interference pattern while $S_R = S_R [q_{1}^2;q_{1}^2]$ only gives a phase-shift. Averaging the interference pattern over different configurations of the external potential (appropriate for impurity averaging in mesoscopic samples) may further decrease the visibility. However, if the two paths are time-reversed copies of each other (as is the case in discussions of weak-localization 3,4), the phase difference between $_{>}$ and $_{<}$ vanishes in a time-reversal invariant situation, so that the impurity-average over the corresponding phase factor does not lead to a suppression in the situation with-

out the bath. On the other hand, the average over exp (iS_R) will, in general, decrease further the magnitude of the interference term¹⁴. In any case, given the simple physical picture presented here, one would not expect S_I and S_R to be able to cancel each other's effects, since they represent, respectively, the real and imaginary part of an exponent. For the special case of a force \mathbf{F} deriving from a bath of harmonic oscillators (linear bath), with vanishing average $\mathbf{F} = 0$ and a linear interaction $\mathbf{\hat{V}} = \mathbf{\hat{T}}$, we have:

$$S_{I} = \begin{array}{ccc} & Z_{t} & Z_{t_{1}} & & D_{t_{1}} \\ S_{I} = & dt_{1} & dt_{2} & q_{1}^{2} & q_{1}^{2} & Re & F_{1}F_{2} & q_{2}^{2} & q_{2}^{2} \end{array}$$
(5)

$$S_{R} = \begin{array}{ccc} Z_{t} & Z_{t_{1}} & D & E \\ dt_{1} & dt_{2} & q_{1}^{2} & q_{1}^{2} & Im & \hat{F_{1}}\hat{F_{2}} & q_{2}^{2} + q_{2}^{2} & (6) \end{array}$$

Here we have used the notation $\vec{q_1} \quad \vec{q}$ (t₁). The angular brackets denote averaging over the equilibrium state of the unperturbed bath. S_I only depends on the symmetrized part of the bath correlator Re \vec{F} (\vec{F} (0) = Dn OE \vec{F} () \vec{F} (0) =2, which becomes the classical correlator FF () \vec{F} (0) = 2, which becomes the classical correlator FF () \vec{F} (0) if or the case of classical Gaussian random noise. In the latter case, S_R vanishes and the influence functional exp(S_I) is simply the classical average of the phase factor

$$\sum_{0}^{Z_{t}} (q^{<}()) \dot{q}())F()d:$$
(7)

If one drops S_R in a dephasing calculation (noting that it does not enter dephasing in semiclassical situations such as those that can be described by Eq. (4)), one effectively replaces the quantum bath by a classical fluctuating force whose correlator is determined by the symmetrized part of the quantum correlator. This contains the zero-point fluctuations, since

$$\begin{array}{c} Dn & \circ E \\ \hat{Q}() \hat{\chi}(0) & / \end{array}$$

$$(2n (!) + 1) \cos(!) = \operatorname{coth} (! = 2T) \cos(!) \quad (8)$$

for the coordinate \hat{Q} of a single bath oscillator of frequency !, with n (!) being the Bose distribution function that vanishes at T = 0.

Although replacing the quantum bath by a classical noise force seems to be a drastic step, it can lead to correct predictions for the dephasing rate in semiclassical situations such as the one discussed above, even at zero temperature. It is instructive to observe how this comes about in an exactly solvable model. A particularly simple situation is the one analyzed by Caldeira and Leggett^{15} (see also Ref. 16). They considered a damped quantum harmonic oscillator, where the initial state consisted of a superposition of Gaussian wavepackets, one centered at the origin, the other at a distance z. In the course of time, the displaced wavepacket oscillates back and forth in the oscillator potential well. Whenever the packets overlap, an interference pattern results (due to the difference in the respective momenta). The environment leads both to damping and dephasing, where the latter typically proceeds at a much faster rate. For our purposes, we are interested in the limit of small damping, where, to a first approximation, the center-of-mass motion of the wave packets is not appreciably altered by friction in the period of the oscillation. In this case, it turns out that, indeed, the result predicted by the approximation Eq. (4) for the attenuation of the interference pattern is correct. This can be seen in Ref. 15 by taking the limit of weak damping $(\ ! \ 0; ! \ ! \ !_R \,)$ in their exact result for the exponent of the attenuation factor (see Eq. (2.13) of Ref. 15) and comparing this to S_{I} (see Eq. (5)), which is to be evaluated for the pair of classical paths followed by the two wave packets, $q_{c1}^{>}(t) = z \cos(!t)$ and $q_{c1}^{<}(t) = 0$. In terms of the quantity $C_{\,00}$ listed in appendix A of the present work, both results may be obtained by multiplying C_{00} by sin $(+t)^2$ and setting = 0; $\stackrel{!}{\leftarrow}$ = !. In addition, in order to obtain $S_{I}[q_{cl}^{>};q_{cl}^{<}]$, the factors sin (! (t 支)) and sin (! (t 支)) must be replaced by $\cos(!t_1)$ and $\cos(!t_2)$. The results obviously coincide for the points in time when the wave-packets meet (!t = -2 + n). We emphasize that the correspondence to the semiclassical result holds only for a situation far away from the ground state of the system, where at least one of the wave packets is in a superposition of highly excited oscillator states.

At zero bath temperature, the dephasing is purely due to spontaneous emission, as pointed out in the discussion of Ref. 15. Energy is transferred from the oscillating wavepacket into the bath and the system relaxes towards lowerenergy states. The dephasing rate is proportional to the total rate _{out} at which the system leaves a given energy level¹⁵. Of course, this rate may be obtained using the Golden Rule (i.e. a master equation description) only for weak coupling, but the qualitative picture seems to be general¹⁵. In the correct description of the physical situation considered here, the rate _{out} is given entirely by the rate of spontaneous emission, ^{em}_{Sp}. However, if S_R is neglected, then S_I describes a classical noise force (equivalent to the zero-point fluctuations of the bath at T = 0) and there will be both induced emission and absorption, proceeding at equal rates ^{em}_{ind} = ^{abs}_{ind}. Therefore, the system is also excited by the bath in that approximation. Nevertheless, we have pointed out above that the total dephasing rate comes out right. The reason is the following: The rate

_{out} is the same in both cases, because the rate of spontaneous emission in the correct description is exactly twice that of induced emission in the approximation:

$$p_{ut} = \mathop{\stackrel{\text{em}}{}}_{\text{sp}} \quad \mathop{\stackrel{\text{em}}{\inf}}_{\text{ind}} + \mathop{\stackrel{\text{abs}}{\inf}}_{\text{ind}} : \qquad (9)$$

At this point, it is easy to see the physical reason why such an approximation may fail near the ground state of the system. Then, the transitions downwards in energy may be blocked, which completely suppresses $\stackrel{\text{em}}{\text{sp}}$, but not $\stackrel{\text{abs}}{\text{ind}}$. We will make this argument more precise in the following sections.

III. DECAY RATES FROM S_I AND GOLDEN RULE: DEPENDENCE ON THE SPECTRA OF BATH AND SYSTEM MOTION

The growth of S_{I} with time depends on the spectral density of the bath fluctuations at frequencies which appear in

the system's motion. Formally, this can be seen by introducing the spectrum of the system motion related to the given pair of paths $q^{>}$ (<) (following the ideas of Refs. 5,12):

$$P(!;s) = \frac{1}{2} \int_{1}^{2} de^{i!} (q^{2}(s + \frac{1}{2}) + q^{2}(s + \frac{1}{2})) (q^{2}(s + \frac{1}{2})) (q^{2}(s - \frac{1}{2}) + q^{2}(s - \frac{1}{2})) (q^{2}(s - \frac{1$$

 $(t_1 + t_2)=2$ and た す are sum and dif-Here, s ference times. Therefore, P is the Fourier transform with of the q-dependent terms in S_T (Eq. (5)). We respect to take $q^{(<)}$ (t⁰) to be zero whenever t⁰ falls outside the range 0:t]

Furthermore, we define hFFi, to be the Fouriertransform of the symmetrized correlator of $\hat{\mathbf{F}}$,

$$hFFi_{!} = \frac{1}{4} \begin{bmatrix} Z_{+1} & Dn & OE \\ & d & e^{i!} & F() \neq (0) \end{bmatrix}; (11)$$

so it is real and symmetric in frequency. The same holds for the system spectrum P (!;s).

Using this, S_{I} can be expressed in the following way:

$$S_{I} = ds \quad d! \text{ hF F } i_{!} P (!;s): (12)$$

If the dependence of P on s is not essential, then S_{T} grows linearly with time t, at a rate given by the "overlap of bath and system spectra". Similar expressions can be derived for a spatially inhomogeneous fluctuating force^{5,12}, leading to an additional k-dependence.

Applying this kind of reasoning to a *single* electron moving in a dirty metal⁵, the system motion is found to contain frequencies up to (at least) $1 = e_1$, such that the growth of S_I with time depends on the bath-spectrum up to this cutoff, including the zero-point fluctuations of the corresponding high-frequency bath modes (which become important at low temperatures). As explained above (sec. II), the description of dephasing in terms of S_{I} alone may be trusted in a semiclassical situation, where the electron is in a highly excited state. This holds even at zero temperature, when the qualitative physical picture is essentially the same as in the model of oscillating wave packets due to Caldeira and Leggett, discussed above: Dephasing is due to spontaneous emission of energy into the bath. The contribution of frequencies up to $1 = e^{1}$ then implies that the spontaneous emission (and the resulting dephasing) is facilitated by the impurity scattering. The physics behind this is well-known in another context: In quantum electrodynamics, the emitted radiation would be called "bremsstrahlung", since it is the scattering off an external potential that induces the electron to emit radiation.

On the other hand, one can express in a similar manner decay rates from a simple Golden Rule calculation¹². For the decay of an initial state نتز, we have

$$a_{i} = 2 \quad d! \quad F F \quad hqqi^{(i)} : \qquad (13)$$

Here $\stackrel{D}{F}\stackrel{E}{F}_{!}$ and $\stackrel{had}{P}\stackrel{(i)}{}_{!}$ are the Fourier transforms of the unsymmetrized correlators of $\mathbf{\hat{F}}$ and $\mathbf{\hat{q}}$, taken in the equilibrium state of the bath or the initial state of the system, respectively:

$$\begin{array}{cccc} D & E & 1 & Z & D & E \\ \widehat{F}\widehat{F} & \frac{1}{2} & d & e^{i!} & \widehat{F}() & \widehat{F}(0) \\ & & & & \\ hqqi_{!}^{(i)} & \frac{1}{2} & d & e^{i!} & hijq()q(0) \\ \end{array}$$
 (14)

The important point to notice is that, near the ground state of the system, the quantum correlator hqqi⁽ⁱ⁾ is very asymmetric in frequency space, as the system can mostly be excited only (! > 0). At low temperatures, the same holds for the bath. Thus, the decay rate (13), containing the product of correlators evaluated at ! and !. is very much suppressed below the value that it would acquire if either the bath-correlator or the system-correlator were symmetrized (becoming symmetric both in time and frequency). Since dropping S_R is equivalent to symmetrizing the bath correlator and, furthermore, semiclassical calculations give a symmetric spectrum P (!;s) of the system motion as well, it becomes clear why there are situations when the decay rate, as deduced from S_I , is finite at low temperatures, while the Golden-rule decay rate vanishes. The question then arises whether any procedure that amounts to symmetrization of the correlators, thus leading to drastically wrong results for Golden Rule decay rates at zero temperature, may be justified to discuss dephasing using a path-integral approach. Observations such as this have led Cohen and Imry to conclude that, in their semiclassical analysis of electron dephasing inside a $metal^{12}$, the contribution of the bath's zero-point fluctuations to the dephasing rate should be dropped. Their argument was not a mathematical proof, but rather drawn from physical intuition and analogies. In the following sections, we try to elucidate the importance of S_R in descriptions of dephasing and decay near the ground state of a system, demonstrating exactly how a cancellation between S_R and S_I may arise.

IV. CANCELLATION OF S_R AND S_I IN LOWEST-ORDER PERTURBATION THEORY

We consider a system which is in its ground state at t = 0before being coupled to a linear bath. From the Golden Rule, we expect there to be no finite decay-rate for the ground state at zero-temperature. It is well-known² how to derive a master equation from the influence functional, by expanding it to lowest order in the exponent, iS_{R} Sr. We display that calculation here in order to point out where and how S_R and S_I do cancel. The time-evolution of the probability to find the system in its unperturbed ground state is:

$$Z = dq_{L}^{2} dq_{L}^{4} \circ (q_{L}^{4}; q_{L}^{2}) (q_{L}^{2}; q_{L}^{4}; t)$$

$$Z = Dq_{L}^{2} Dq_{L}^{4} \circ (q_{L}^{4}; q_{L}^{2}) e^{i(S_{0}^{2} - S_{0}^{2}) + iS_{R} - S_{1}} \circ (q_{0}^{2}; q_{0}^{4})$$

$$Z = 1 + Dq_{L}^{2} Dq_{L}^{4} e^{i(S_{0}^{2} - S_{0}^{2})} \circ (q_{L}^{4}; q_{L}^{2})$$

$$(iS_{R} - S_{L}) \circ (q_{0}^{2}; q_{0}^{4}) : (15)$$

Here $_0$ is the density matrix of the unperturbed ground state, and the path integration also includes integrals over the initial and final coordinates. Now one can insert the expressions for S_R and S_I from Eq. (6). The integration over the trajectories and the endpoints produces the (unsymmetrized) correlators of the system coordinate, such as (for $t_1 > t_2$):

Z

$$D q^{2} e^{iS_{0}^{2}} = 0 (q_{L}^{2})q^{2} (t_{1})q^{2} (t_{2}) = 0 (q_{0}^{2}) = 0$$

 $h_{0} jq(t_{1})q(t_{2}) j_{0} ie^{iE_{0}t} : (16)$

Evidently, both the unperturbed action S_0 and the integrations involving the state $_0$ are essential for obtaining the correct correlator. In the long-time limit, we obtain the decay rate which also follows from the Golden Rule (13):

Only the term growing linearly in time has been retained at the end of this equation.

Note that, of course, there will always be a small reduction in the probability of finding the unperturbed ground state after switching on the interaction, since the ground state of the coupled system contains contributions from other system states as well. This point has been discussed in more detail recently for the case of the damped harmonic oscillator¹⁷. It does not contribute to $_0$ which describes the decay linear in time.

We emphasize that the decay of P_0 (t) is not governed by S_I alone: If S_R were dropped, then only the (real-valued) symmetrized version of the bath correlator would appear in Eq. (17). In that case, the decay rate would only depend on the (real-valued) symmetric part of the system correlator as well. Therefore, a finite decay rate of the ground state would result even at zero temperature, where really there would have been none. The physical reason is the heating introduced by the classical noise field, whose correlator contains the zero-point fluctuations of the original quantum bath, as we have discussed before.

On a formal level, we may argue that the decay of the density matrix is not governed by S_{I} alone since the "weighting factor" that is used when "averaging" over many paths contains the *phase* factor $\exp(i(S_0^{>}))$ $S_0^{<}$)). Therefore, such an average is not the same as a classical average, for which the decay could not be overestimated by dropping S_R , since then we could use $\pm xp$ (iS_R) S_T)ij hexp (S_T) i Furthermore, the integration over the density matrix of the initial state is obviously essential, as it is needed to produce the correct form of the system correlator. Both facts mean that it would be premature to draw any conclusions about dephasing and decay at an early stage of the calculation, by merely looking at the influence action.

V. EXACTLY SOLVABLE LINEAR SYSTEMS: "CANCELLATION TO ALL ORDERS" FOR THE DAMPED OSCILLATOR

In order to show how S_R and S_I can cancel also beyond lowest-order perturbation theory, we will now turn to linear quantum dissipative systems. Although the exact solutions for the damped harmonic oscillator and the free particle have been well-known for a long time¹⁸, we will review the essential steps in the derivation, pointing out where S_R and S_I do enter. We will first turn to the oscillator, for which the Golden Rule result suggests that one would obtain an artificial decay of the ground state at zero temperature if S_R were neglected. For simplicity, our discussion is restricted to T = 0, since that is the limit where these effects show up most clearly.

For any linear system which is linearly coupled to a bath of oscillators, the propagator J for the density matrix (Eq. (2)) can be evaluated exactly, since the integration over system paths is Gaussian. In fact, J is found to be given by the "semiclassical" result, i.e. an exponential containing the action along stationary paths, multiplied by a prefactor which does not contain the endpoints of the paths (a specialty of linear systems):

$$J(q_{t}^{>};q_{t}^{<};q_{0}^{<};q_{0}^{<};t) = \frac{1}{N(t)}e^{iS[q_{c1}^{>};q_{c1}^{<}]}; \quad (18)$$

with S = $S_0^>$ + $S_0^<$ + S_R + $iS_I.$ The paths $q_{cl}^{>\,(<\,)}$ make the full action stationary,

$$\frac{S}{q_{1}^{2}} = \frac{S}{q_{1}^{2}} = 0; \qquad (19)$$

and fulfill boundary conditions of the form $q_{cl}^{>}(0) = q_{0}^{>}$. The prefactor N (t) can be obtained most easily from the condition for normalization of the density matrix,

Z

$$dq_{\pm} J (q_{\pm};q_{\pm};q_{0};q_{0};t) = 1:$$
 (20)

N (t) is found to be independent of $S_{\tt I},$ but it does depend on $S_{\tt R}$ (see, for example, the general proof in app. E of Ref. 7).

In the following, we will turn to the special case of the *Ohmic* bath, which leads to a velocity-proportional friction force and has a power spectrum rising linearly at low frequencies (for T = 0):

$$hFFi_{!} = \frac{j!}{2}; \qquad (21)$$

We will argue below that, in the case of the damped harmonic oscillator, no essential qualitative result will be changed by using other bath spectra, unless these have an excitation gap exceeding the oscillator frequency.

As usual, we introduce the center-of-mass coordinate R() ($\dot{q}() + \dot{q}()$)=2 and the difference coordinate r() $\dot{q}()$ $\dot{q}()$ in order to write down the equations obtained from (19), for the case of the Ohmic bath:

$$\frac{d^{2}r}{d^{2}} = \frac{dr}{d} + !_{0}^{2}r = 0$$
(22)
$$\frac{d^{2}R}{d^{2}} + \frac{dR}{d} + !_{0}^{2}R = i_{0}d^{0}Re \hat{F}()\hat{F}()r()^{0}(23)$$

Here =m is the damping rate and b is the unperturbed frequency of the oscillator. Note that the second equation leads to a complex-valued solution R (). It is also possible to formulate the calculation slightly differently^{18,19}, by using stationary solutions with respect to the real part of S only. In any case, inserting the solutions R (R_{cl}) and r (r_{cl}) into the action S shows that the imaginary part of the action $S_{cl} = S[k_{ll}^2; q_{cl}^2]$ is determined directly only by S_I . As expected, S_{cl} turns out to be a bilinear expression in the endpoints $R_t; R_0; r_t; r_0$:

$$\begin{aligned} \text{ReS}_{\text{cl}} &= S_0^{>} + S_0^{<} + S_{\text{R}} = \\ &\quad \text{R}_{\text{t}} r_{\text{t}} \text{L}_{\text{tt}} + \text{R}_{\text{t}} r_0 \text{L}_{\text{t0}} + \text{R}_0 r_{\text{t}} \text{L}_{0\text{t}} + \text{R}_0 r_0 \text{L}_{00}(24) \\ \text{Im S}_{\text{cl}} &= S_{\text{I}} = r_0^2 \text{C}_{00} + 2 r_0 r_{\text{t}} \text{C}_{0\text{t}} + r_{\text{t}}^2 \text{C}_{\text{tt}} : \end{aligned}$$
(25)

All the coefficients are real-valued functions of time. For our purposes, it is sufficient to know that the entries of the matrix L are independent of S_{I} (while they do depend on the damping from S_{R}). In contrast, the entries of C depend both on S_{I} and S_{R} . The dependence on S_{R} arises only because the path r, which is inserted into Eq. (5), is affected by the friction described by S_{R} ; see Eq. (22). The entries of C are proportional to the strength of the bath fluctuations (i.e. the magnitude of hF F i_{I}). Explicit expressions for all of these coefficients²⁰ can be found in Ref. 18 (cf. their section 6). For reference, the quantities used in the following discussion have been listed in appendix A. At this point, evaluation of the coefficients for Im S_{cl} shows that they will grow in time beyond all bounds, regardless of whether the path r is calculated by taking into account S_R or neglecting it (setting = 0 in Eq. (22)). However, no conclusion about dephasing and decay can be drawn from this, since the foregoing discussions lead us to expect that any potential cancellation between S_R and S_I will take place only *after* proper integration over the initial density matrix. This can be deduced from the derivation of the Golden Rule decay rate given in the preceding section. It is also very reasonable that S_I grows without bounds for any semiclassical path r(), since all paths contribute more or less to the time-evolution of all eigenstates, and decay of excited states is perfectly correct in the model of the damped harmonic oscillator.

The initial ground state density matrix has the form

$$\begin{array}{ll} {}_{0}(\mathbf{q}_{0}^{<};\mathbf{q}_{0}^{<}) \ / \ \exp & (\mathbf{q}_{0}^{<\,2} + \mathbf{q}_{0}^{<\,2}) = (4 \ \mathbf{q}_{0}^{2} \ _{0}) \\ \\ = \ \exp & (\mathbf{R}_{0}^{2} + \mathbf{r}_{0}^{2} = 4) = (2 \ \mathbf{q}_{0}^{2} \ _{0}) \ ; \qquad (26) \end{array}$$

with $\dot{q}^2_{0} = 1 = (2m !_0)$ as the square of the ground state width. After performing the Gaussian integrals over R₀ and r₀, which are needed to evaluate Eq. (1), we obtain the final result for the density matrix ($\dot{q}^2; q_1^{<}; t$) at a time t after switching on the interaction with the bath. It has the general form

$$\frac{1}{N'(t)} \exp (aR_t^2 + bR_t r_t + cr_t^2) : (27)$$

In particular, the prefactor of R_t^2 gives the width of the probability distribution at time t,

$$\hat{q}^2$$
 (t) = (2a)¹ : (28)

Let us now look at the behaviour of the width in order to analyze the effect of dropping S_R in the calculation. The expression for q^2 (t) has been derived already by Caldeira and Leggett¹⁸ (cf. their Eq. 6.34):

$$q_1^2$$
 (t) = 2C₀₀ + m !₀=2 + L₀₀² = (2m !₀) = L_{t0}² : (29)

This expression clearly contains quantities both from the imaginary part (C_{00}) and from the real part (L_{00} ; L_{t0}) of the action. Formally speaking, real and imaginary parts have become intermixed when integrating over R_0 , due to quadratic completion of the expression iR_0 ($r_tL_{0t} + r_0L_{00}$) m $!_0R_0^2$ in the exponent (the former term is from Eq. (24), while the latter term is from the initial density matrix, Eq. (26)). Physically, this result is to be expected, since the growth of the width of the probability distribution will be governed by the balance between the bath fluctuations (S_I) and the friction (S_R).

The complete time-evolution of q^2 (t) after switching on the interaction at t = 0 is displayed in Fig. 1. This includes cases where the value of the damping constant has been artificially set to zero or to other values different from that prescribed by the fluctuation-dissipation theorem (FDT), which connects to the strength of the bath fluctuations described by hFFi, (see Eq. (21) for the case of T = 0). If S_R is neglected completely (no friction, = 0), heating takes place, causing the variance to grow linearly in time, at a rate given by the Golden Rule expression involving the symmetrized bath correlator. Formally, the growth of C_{00} with time cannot be compensated in that case, since L_{00} and L_{t0} acquire their original unperturbed values. For small > 0, the width saturates at a value much larger than that of the ground state. On the other hand, if the friction is too strong, the width saturates at a value below that of the ground state, and the Heisenberg uncertainty relation is violated (since p^2 also shrinks). This point is discussed in the book of Milonni²¹ in the context of an atom interacting with the vacuum electromagnetic field. If, however. has the correct value prescribed by the FDT, q² changes only slightly from its unperturbed value.

In fact, for a linear system like the quantum harmonic oscillator, its behaviour under the action of the linear bath can be described and understood entirely using a classical picture²². The reason for this is as follows: one starts out with an initial state for system and bath that has a positive definite Gaussian Wigner-density, which can be interpreted directly as a classical phase space density whose time-evolution then corresponds one-to-one to the evolution of the Wigner-density (see appendix B). Thus, the quantum dissipative dynamics may be described by a *classical* Langevin equation including friction and a fluctuating driving force:

$$\frac{d^2q}{dt^2} + \frac{dq}{dt} + !_0^2 q = \frac{1}{m} F (t) :$$
 (30)

In this equation, the damping term incorporates the effects of S_R . It counteracts the fluctuations of F () that are described by S_I in the path-integral picture and whose correlation function is, therefore, given by the *symmetrized* part of the bath correlator.

The behaviour of the width has been discussed above using the influence functional, but it can be understood more easily in this picture. We can use the susceptibility of the damped oscillator to find its response to the force fluctuations F, which, at zero temperature, are entirely due to the zero-point fluctuations. In particular, for the limit t ! 1, we obtain

$$q_{1}^{2}$$
 (t) ! d! j (!) j² hFF i₁; (31)

where

$$(!) = \frac{1=m}{!_0^2 \quad !^2 \quad i!}$$
(32)

is the susceptibility of the damped oscillator, and hFF i_1 is the symmetrized power spectrum (see Eq. (21)). For the



Figure 1: The variance ² (t) q^2 (t) of the probability density (q;q;t) for the damped harmonic oscillator, plotted in units of the unperturbed ground state variance 2_0 , as a function of time t for the following friction strengths (top to bottom): =(2 hFF i₁₌₁) = 0; 0:5; 1; 2 (other parameters are m = !_0 = 1). A ratio of 1 corresponds to the correct value prescribed by the FDT, while = 0 means S_R has been neglected completely, such that 2 (t) grows without bound. The dashed lines give the limit for t ! 1 (see Eq. (31)).

correct equilibrium state (obtained by fully keeping S_R and S_I), the width at t! 1 can also be found by applying the FDT to the oscillator coordinate (cf. Eq. 6.37 of Ref. 18). In contrast, the effect of dropping S_R is obtained by letting ! 0 in Eq. (30), while hr F i, is kept fixed.

The example of the damped harmonic oscillator demonstrates that the proper behaviour near the ground state of the system cannot be observed at the early stages of the calculation, by looking at the action evaluated along classical paths (Eqs. (24), (25)), regardless of whether these paths properly include the damping or not. Only after correct integration over the initial state density matrix the contributions from S_R and S_I can compensate each other (in Eq. (29)), such that no artificial finite decay rate of the ground state results, if S_R is kept.

VI. POWER-LAW BEHAVIOUR IN QUANTUM BROWNIAN MOTION

We now turn to a discussion of the free particle, i.e. the Caldeira-Leggett model of quantum Brownian motion. There, the reasoning is the same as before in principle: Neglecting S_R means that the zero-point fluctuations of the bath contained in S_I may heat up the particle. However, there is an important difference: The spectrum of the free motion does not contain any resonance peak, unlike that of

the harmonic oscillator. Therefore, only the low-frequency components (! ! 0) of the bath spectrum will contribute to heating and decay (compare Eq. (12)), which results in a peculiar long-time behaviour^{18,19,23} that cannot be captured using the Golden Rule, as has been emphasized in Ref. 7.

One can observe the distinction between oscillator and free particle most easily in the picture of the classical Langevin equation (30) introduced above. In appendix B, it is shown how the density matrix propagator J may be obtained in general via the time-evolution of the Wigner density. This approach is physically more transparent than the path-integral calculation. For the purposes of our present discussion, however, it will suffice to consider the timeevolution of the diagonal elements of the density matrix in momentum space. These have also been analyzed in Ref. 7, in order to demonstrate the failure of the Golden Rule calculation to describe the decay of the original ground state at zero temperature. We will confirm and explain the outcome of this analysis using a simple argument based on the Langevin equation (30). Then we will point out the role of $S_{\mathsf{R}}\,$ and describe the important difference to the damped oscillator.

If the particle has momentum p_0 at time 0, its momentum p at time t will be determined by the fluctuating force F () in the following way (by solving Eq. (30) for d = 0):

$$p = p_0 e^{t} + ds e^{(t s)} F(s):$$
 (33)

Since F () is a Gaussian random process, the probability density of finding a momentum p at time t is a Gaussian, of variance

$$p^{2} \qquad \begin{array}{cccc} Z_{t} & Z_{t} \\ p^{2} & ds_{1} & ds_{2}e \end{array} \qquad \begin{array}{cccc} (2t \ s_{1} \ s_{2}) \\ bF \ (s_{1})F \ (s_{2})i: (34) \end{array}$$

The behaviour of p^2 as a function of time is as follows (at T = 0): As long as t 1= , friction is unimportant and it is the double time-integral over the correlator of F which is to be evaluated. For the Ohmic bath considered here, the power spectrum is relatively strong at low frequencies (rising linearly with !, see Eq. (21)). This leads to a slow decay in time, hF (t)F (0)i / 1=t², which results in a logarithmic growth of p^2 ,

$$p^2 = 2 - \ln(!_c t)$$
: (35)

Here $!_{c}$ is the cutoff frequency of the bath spectrum and we have assumed $!_{ct}$ 1. It is this logarithmic behaviour which cannot be obtained using the Golden Rule approximation, to be discussed below. At later times, t 1= , the growth saturates at a constant value,

$$p^2 ! - \ln (!_c =) :$$
 (36)

The perturbation expansion in the coupling between system and bath can be carried out by expanding the influence action in the propagator J (Eq. (2)) in powers of iS_R S_I (see Eq. (15) in the present article or Eq. (E13) of Ref. 7). We want to discuss the time-evolution of the diagonal elements $_{pp}$ (t) of the density matrix in momentum space. The evolution starts from the equilibrium density matrix, which is a Gaussian momentum distribution of variance $p_0^2 = m T$ (the following formulas will be analyzed for arbitrary T). After a time t, the density matrix is still a Gaussian, but of variance (compare Eq. (33)):

$$p^2 = p_0^2 e^{2t} + p^2$$
: (37)

From this we can easily derive the result for $_{pp}^{(1)}$ (t) which could alternatively be obtained by first expanding J to first order in iS_R S_I , going over to the momentum representation and finally integrating over the initial density matrix (as was done in Ref. 7; see Eq. (E18), which is, however, still written in terms of matrix elements of the coordinate operator):

$$pp (t) = \frac{1}{2 p^{2} i} \exp \left(\frac{p^{2}}{2 p^{2} i}\right)$$

$$pp (0) \left(1 + \frac{p^{2} p_{0}^{2}}{2 p_{0}^{2} i} - \frac{p^{2}}{p_{0}^{2} i}\right) + (38)$$

In evaluating p^2 , only the terms up to first order in and hFFi, must be kept for the purposes of this expansion. At finite temperature T, the behaviour of p^2 at long times t 1=T is governed by the linear decrease stemming from $p_0^2 e^{2t}$ p_0^2 (1 2 t) and the linear increase from p^2 2 Tt If the initial density matrix really describes the equilibrium distribution, then detailed balance holds and these terms cancel. For times t 1=T (or arbitrary times at T = 0), the time-evolution of p^2 is governed by \overrightarrow{p} , evaluated for the correlator hFF i taken at T = 0 (which only contains the zero-point fluctuations of the bath). For consistency of the expansion, p^2 in Eq. (34) has to be evaluated by setting = 0. Therefore, only the discussion given above for times t 1= turns out to be relevant. The logarithmic growth of p given in Eq. (35) corresponds to a power-law behaviour of the (exact) time-evolution of $_{pp}$ (t) for t 1= . This logarithm, that appears in the perturbation expansion (38), is "overlooked" in the Golden Rule approximation, where only terms growing linearly with time are kept. Therefore, the Golden Rule rate turns out to vanish⁷. These powerlaws are characteristic of the density matrix propagator J of quantum Brownian motion at zero temperature (see appendix B).

On the other hand, the effect of neglecting S_R may be analyzed by setting the friction constant to zero in Eq. (34). Then, the spread p^2 of the momentum distribution grows without bounds, which is obviously not the correct physical behaviour. It is qualitatively similar to the heating produced in the damped oscillator model. However, since the growth proceeds only logarithmically in time, the artefacts of this approximation, when applied to the free particle, are not nearly as drastic as the finite decay rate of the ground state observed for the oscillator. In particular, within perturbation theory, no qualitative change is obtained by dropping S_R for the free particle (no finite decay rate is produced).

Furthermore, if we had considered a super-Ohmic bath (e.g. due to phonons), whose spectrum decays faster than $!^{1}$ for ! ! 0, the growth of p^{2} would saturate even without friction (as it should, since there is no velocitydependent friction for such a bath). In contrast, the behaviour of the harmonic oscillator discussed above would remain qualitatively the same as for the Ohmic bath, since its decay depends on the bath spectrum at the resonance frequency of the oscillator and is not affected in any essential way by the details of the spectrum at low frequencies. This is consistent with the fact that in the oscillator case already the Golden Rule is sufficient to obtain an essentially correct picture of the artificial decay produced by dropping S_R , while, for quantum Brownian motion, it fails to describe the subtle power-laws associated with the lowfrequency (long-time) properties of the Ohmic bath at zero temperature.

The difference between oscillator and free particle also shows up clearly in the behaviour of the imaginary part of the influence action: For the free particle, S_I always grows only logarithmically with time t (at T = 0), independent of whether one keeps S_R or sets it to zero (i.e. 0 in the equations of motion). This can be derived from the formulas given in appendix B or those of Ref. 7, Appendix E (or Ref. 18, section 6, with the proper limit $!_R$! 0 for the free particle).

Hence, we conclude that for the oscillator (or any motion containing an extended spectrum) the effects of dropping S_R are much more pronounced than for the free motion, since they lead to an artificial finite decay rate. We have also observed that this effect can already be understood within the framework of the Golden Rule approximation, whereas that approximation is incapable of describing the more subtle "power-law decay" found for the free particle.

VII. QUALITATIVE DISCUSSION OF OTHER MODELS: NONLINEAR COUPLING AND PAULI PRINCIPLE

It could be argued that our discussion of the damped oscillator has only demonstrated the importance of S_R in preventing an artificial decay of the *ground state* and has little to do with dephasing. Indeed, if one wants to discuss dephasing, one should rather consider the decay of a coherent superposition of the ground state and some excited state. This is relevant both for arbitrary nonequilibrium situations as well as for the calculation of the system's linear response, where the perturbation creates a superposition of excited states and the ground state. In the example of the damped harmonic oscillator, full relaxation into the ground state. This holds regardless of the bath spectrum, provided the latter does not vanish around the resonance frequency of the oscillator. For weak coupling, the decay

of the density matrix (and, therefore, the linear response) may be described conveniently using a simple master equation approach. What is more, even if one neglects S_R in a path-integral calculation, the predicted decay rates will be correct, excepting only that for the ground state, which we have discussed above. The reason for this, however, is a specialty of the linearly damped harmonic oscillator. Since the bath couples to the coordinate operator, whose matrix elements only connect adjacent oscillator levels, the relevant "system correlator" (in Eq. (10)) turns out to be symmetric for any state but the lowest one. Therefore, the decay rate will depend only on the symmetric part of the bath correlator as well, see Eq. (17). Hence, for all the excited states, the evaluation of the total decay rate is not affected by dropping $S_{\ensuremath{\mathsf{R}}}$. This point has already been discussed in connection with the model of two oscillating wave packets, at the end of section II.

Since we are interested in demonstrating the importance of S_R for the dephasing of a superposition of low-lying levels, the natural choice is to consider a bath with an excitation gap larger than the resonance frequency of the oscillator (see Fig. 2). In that case, we expect the lowest levels to remain coherent in the correct description. Unfortunately, in the model of the linearly damped oscillator discussed above, we would not obtain any finite decay rate, *regardless* of whether S_R is kept or not. This is because the corresponding classical noise force is out of resonance and cannot heat the oscillator.

Therefore, in order to analyze a situation where the importance of retaining S_R is displayed even for excited states, we have to go beyond exactly solvable linear models. As a consequence, the following discussion is necessarily incomplete insofar as we cannot give exact proofs of the statements to be made below. These statements will be based on the experience acquired in the simpler models discussed above. We will also make use of some Golden Rule type arguments, noting that the Golden Rule has been sufficient to understand the behaviour of the damped oscillator in a qualitatively correct way. In any case, we believe it is useful to contrast the results obtained by dropping S_R with the "commonly accepted" picture.

Let us, therefore, consider the following model: We retain the harmonic oscillator, but change the coupling between system and bath from $\hat{\nabla} = \hat{\nabla} \hat{\nabla} = f(\hat{\nabla} \hat{\nabla})$ with a *nonlinear* function f. This is called state-dependent friction²². In that case, the transition matrix elements of the system operator f(q) will, in general, be nonzero between any two states, which is the important difference to the linear case. If we introduce a bath spectrum with a gap larger than n!₀, then, for sufficiently weak coupling and according to a simple Golden Rule calculation, the first n excited states will not decay at zero temperature. Any coherent superposition of these states will therefore remain coherent. The full, non-perturbative picture will be slightly more complicated, but the essential point should remain the same: The first excited states will acquire an admixture of other levels and become entangled with the states of the bath. They will experience some frequency shifts and the transition matrix elements will be renormalized. Of course, the new eigenstates of the coupled system will



Figure 2: The models described in the text. Left: A bath with a gap would lead to the upward transitions indicated by grey arrows if (and only if) S_R were neglected. Right: A sea of fermions may block downward transitions (dashed) even though they are possible in the single-particle picture.

remain in a coherent superposition forever (by definition). The n + 1 lowest ones still have a discrete spectrum and are in direct correspondence to the initial unperturbed states. When switching on the interaction at t = 0, a partial decay will result. Physically, this corresponds to the relaxation of the initial state into the selfconsistent coupled state of system and bath (see Ref. 19 for a discussion of these issues in the case of the free particle). However, the initial decay will saturate in time on a short time-scale. During this transient adjustment, the reduced system density matrix becomes mixed to a small extent (if the coupling is weak), but no non-saturated long-term decay (indicated by a finite decay rate) results. In any case, the transient decay is an artefact of the procedure of suddenly switching on the interaction. A full calculation of, for example, the linear response properties of the dissipative system would start out with the selfconsistent ground state of the fully interacting system (as was done in Ref. 23 for the free particle).

In contrast, dropping S_R would lead to a completely different picture, since then S_I would correspond to a classical noise force. Starting from any of the first n excited states, the fluctuating field would always be able to induce transitions up in energy, thus leading to a finite decay rate. The role of S_R in the correct description of a quantum noise force is precisely to cancel these upward transitions, as has been demonstrated in the case of the linearly damped oscillator. Due to this finite decay rate, any initially coherent superposition would get destroyed, leading to complete dephasing. No saturation of the decay could be observed, in spite of the gap in the bath spectrum.

It is this behaviour that is described qualitatively correctly by evaluating S_{I} along semiclassical paths of the oscillator. The expression for S_{I} , Eq. (5), is now to be changed simply by replacing q by f (q). While an (unperturbed) oscillator path of the form q() / $\cos(!_0t)$ only contains frequencies $!_0$ which cannot couple to the gapped bath, the function f (q()) will, in general, contain all higher harmonics as well. This directly corresponds to the character of the symmetrized system spectrum, i.e. the Fourier transform of hff (q()); f (q(0))gi. These frequencies contained in the system motion then will couple to the bath spectrum, leading to an unbounded growth of S_{I} with time. Similarly, if we were to use a bath without a gap, but where the spectrum falls off fast towards low frequencies, then the correct description would also yield decay rates that quickly become smaller when going towards the ground state, while they would saturate at a finite, comparatively large level if S_R were dropped.

The present work has been motivated in part by the ongoing controversial debate on low-temperature dephasing in an interacting disordered fermion system 6,7,8,10,11,12,13,24 , which has led to a revival of interest for the general question of dephasing in mesoscopic systems^{17,25,26,27,28,29}. Since the usual Feynman-Vernon influence functional deals only with a single-particle situation, it cannot be applied directly to this problem. However, the authors of Refs. 6,7,8 have succeeded in deriving an extension of the usual influence functional to the many-fermion situation, using an exact procedure. In their result, the Fermi distribution (and, therefore, the Pauli principle) only enters S_R , while S_I is unaffected (Eqs. (54) and (55) of Ref. 6). Therefore, the dephasing rate, as read off from S_I , is found to equal the rate which would also be obtained in a purely single-particle calculation. Although we do not analyze the full problem of weak-localization here, we will use the insights gained above in order to explain why, in our opinion, the assumption that the dephasing rate can be derived from S_{T} alone must be proven instead of being taken for granted. This holds even when S_R vanishes on the relevant pairs of classical time-reversed trajectories, since, in general, the integration over the fluctuations away from these trajectories will be essential for obtaining a cancellation between the effects of S_R and S_I (see sec. IV), in addition to properly taking into account the initial density matrix. Furthermore, we note that in these calculations the detailed form of the bath spectrum at low frequencies turns out to be unimportant for the essential result of a finite zero-temperature dephasing rate. This is in marked contrast to the case of the free particle (where the Ohmic bath plays a special role²³; see also Refs. 28,29), but similar to what is observed for the damped harmonic oscillator when S_R is neglected. It is for this reason that we have chosen a bath with an excitation gap in order to demonstrate the importance of S_R , since there the effects come out most clearly, although they are also present for other bath spectra (compare the discussion above).

The model situation of the damped oscillator described above already contains one key ingredient related to the description of dephasing for an electron moving inside a disordered metal, namely an extended system spectrum: Due to the impurity scattering, the system spectrum (e.g., of the velocity operator) contains frequencies up to (at least) the elastic scattering rate, which may couple to the zeropoint fluctuations contained in the bath spectrum. This is in contrast to the free particle, which can only couple to the low-frequency bath modes (see the discussion at the end of section VI). As before, the importance of S_R will be visible most easily for a bath containing an excitation gap. For such a bath, the *free* motion (without impurity scattering) will not show any nontransient decay, *regardless* of whether S_R is taken into account or not. In contrast, the model to be discussed below, which is more similar to the situation in the disordered metal, will show a finite decay rate of low-lying levels if and only if S_R is neglected.

Apart from the extended system spectrum, we have to take into account another important feature of the calculations^{3,6,7,8,12} concerning electrons in a disordered metal: the use of the semiclassical analysis. For linear systems the semiclassical result for the path-integral is exact. such that it can even be used near the ground state of the system, as we have done it here. However, in such a case, the size of the fluctuations around the path is comparable to the amplitude of the path itself, which is not the situation in which semiclassics is usually applicable. In order to gain intuition for a typical semiclassical situation, we could reconsider the example with the nonlinear coupling to a gapped bath given above: Imagine an initial superposition of fast wavepackets, whose wavelength is much smaller than the packet size but whose excitation energies are still below the bath threshold. They would show a finite decay or dephasing rate (for the reasons given above), if only S_{T} were used for the analysis, but not in the full calculation. Still, for this scenario one might argue that it has been clear from the outset that the semiclassical analysis cannot be trusted, since those high-frequency components in the system spectrum that are responsible for the decay are necessarily larger than the energetic distance to the ground state.

It is only in a system of degenerate fermions that the following two conditions can be fulfilled all at once: On the one hand, the semiclassical analysis of a single (noninteracting) electron moving at the Fermi level is valid for an external potential which is sufficiently slowly varying, such that the system spectrum (concerning the motion of the single electron) only contains frequencies much smaller than the Fermi energy. On the other hand, the whole many-particle system may be near (or in) its ground state.

While the first feature would lead one to believe that S_R may be omitted, we have learned from the examples discussed before that this is likely to be incorrect whenever the second condition holds as well. We stress once again that the two conditions are mutually exclusive in a single-particle problem, which is the reason why such considerations have not played any role in influence-functional calculations up to now.

In order to render the discussion concrete, we can once again make use of the example with a nonlinear coupling f (q) given above, provided we suppose the N lowest oscillator states to be filled up with fermions initially (see Fig. 2). For a relatively smooth function f, the matrix elements of f (q) only connect states within a range much smaller than $_{\rm F} = N !_0$, which, in our example, should still be larger than the gap $n!_0$ of the bath. Dephasing of a *single* particle near $_{\rm F}$ can then certainly be described fully within the semiclassical analysis, and, to a good approximation, using S_T alone, as has been discussed above. The same holds for the many-particle system, if one explicitly considers a classical noise force, where S_R is absent. In that case, the many-particle problem can really be treated as a collection of independent single-particle problems, as is the case for any external time-dependent potential. Only in the end

an average of the full Slater determinant over all possible realizations of the external noise has to be carried out.

If the dephasing rate is calculated solely from S_{I} , it turns out to be finite^{6,7,8} and not to depend at all on the distance to the Fermi surface. This is consistent with the fact that the the value of $_{\rm F}$ does not even appear in the singleparticle calculation. On the other hand, for a quantum bath S_R does not vanish and should be included in the influence action. It is true that for the case of a highly excited single particle the dephasing rate comes out correct, regardless of whether $S_{\ensuremath{\mathbb{R}}}$ is kept or not, as we have discussed before. However, we have also pointed out that there is an important physical difference between the two calculations. Since in the correct approach (including S_{R}), the transitions induced by the bath are purely downwards in energy (accompanied by spontaneous emission into the bath), it is reasonable to expect that, in the many-particle problem, they will be blocked by the Pauli principle. In any case, this is what is found using the Golden Rule. Judging from these arguments, the contribution to S_{I} from the nonzero overlap of the (symmetrized) system spectrum with the high frequencies of the bath spectrum does not imply dephasing and decay, but rather a renormalization, similar to that obtained for an electron interacting with optical phonon modes (a gapped bath), leading to the formation of a polaron. The formation of the polaron will be visible as an initial transient decay of the single-electron density matrix (for the artificial case of factorized initial conditions), which saturates on a short time-scale. Therefore, to lowest order, for a bath with a gap no finite relaxation rate of the lowest-lying single-particle excitations above the Fermi sea is expected, just as for the lower levels of the harmonic oscillator in the single-particle model with nonlinear coupling to a gapped bath.

We have to qualify this statement by taking into account the fact that the coupling between electrons and bath always also induces an effective interaction between the electrons. In this way, a given electron becomes coupled indirectly to the bath of other electrons, such that scattering processes will, indeed, lead to a finite decay rate even for those low-lying excited levels. This precludes any rigorous proof demonstrating the complete absence of decay and dephasing for low-lying levels even in such a rather simple situation, in spite of the assumption of an excitation gap in the bath spectrum. However, the consequences of the effective interaction can be distinguished easily from the finite dephasing rate that would be predicted by looking at S_{I} alone. If the coupling to the bath is of strength g, then the latter rate would go as q^2 . In contrast, the effective interaction (obtained after integrating out the bath) will itself be of strength q^2 , such that the resulting relaxation rate (due to coupling of an electron to the density fluctuations of other electrons) will be of fourth order in g. In addition, of course, the rate will depend strongly on the distance to the Fermi surface, vanishing when the Fermi energy is approached.

VIII. CONCLUSIONS

We have tried to demonstrate that the real part S_R of the Feynman-Vernon influence action cannot be neglected in an analysis of dephasing and decay near the ground state of a system, in spite of the fact that, in the simplest situations (involving highly excited states and the semiclassical analysis), it is only the imaginary part S_{I} which enters the dephasing rate. To this end, we have discussed how S_R and S_I may cancel each other's effects not only in lowest-order perturbation theory but also to all orders, by examining exactly solvable linear quantum dissipative systems. In general, the cancellation is only found after proper integration over the fluctuations away from semiclassical trajectories, taking into account the action of the unperturbed system and its initial density matrix. Furthermore, we have pointed out an essential difference between the damped oscillator and the free particle with respect to this issue. We have argued that the insight obtained in the case of the damped oscillator is also applicable to nonlinear systems and important for discussions of dephasing in systems of disordered degenerate fermions.

In summary, it may be possible to discuss dephasing and decay without considering S_R either if the noise is nearly classical (external nonequilibrium radiation or bath in the high-temperature limit) or if the system itself is in a highly excited state. Otherwise, dephasing rates obtained solely from S_I are bound to come out finite at zero temperature in most cases, even when a simpler Golden Rule calculation gives vanishing results. Judging from the examples discussed above, this is probably not because such a non-perturbative procedure goes beyond the Golden Rule approximation, but because it neglects some physics already contained even within this approximation.

Still, although the primary message of this paper is that special care has to be taken in extracting dephasing rates from calculations using the influence functional, we emphasize at the same time that there is no general proof showing the impossibility of zero-temperature dephasing near the ground state. There cannot be such a proof, since there is evidently at least one counter-example (the Caldeira-Leggett model of quantum Brownian motion and similar models^{18,28,29}), and, furthermore, there is no generally applicable definition of "dephasing" that is useful under all conceivable circumstances.

Acknowledgments

I would like to thank Christoph Bruder, Jan von Delft, Dimitri Golubev and Andrei Zaikin for stimulating discussions. This work has been funded by the Swiss National Science Foundation.

Appendix A: SOME QUANTITIES FOR THE DAMPED OSCILLATOR

In this appendix, we list, for purposes of reference, the quantities relevant to our discussion of the time-evolution of the quantum damped harmonic oscillator. These can also be found in Ref. 18 (in a slightly different notation).

The quantity C_{00} arises in evaluating S_{I} along a pair of semiclassical paths, by inserting the solution r() of Eq. (22) for the boundary conditions given by r_{t} ; r_{0} and determining the coefficient of r_{0}^{2} in the result (see Eq. (25)). Necessarily, C_{00} contains only the symmetric part of the bath correlator, since S_{I} depends only on that:

$$C_{00} = \frac{1}{4} (\sin (t))^{2} dt_{1} dt_{2} exp_{1} \frac{1}{2} (t_{1} + t_{2}))^{0} dt_{1} dt_{2} exp_{1} \frac{1}{2} (t_{1} + t_{2}) dt_{1} dt_{2} exp_{1} \frac{1}{2} (t_{1} + t_{2})) ff^{2} (t_{1}); f^{2} (t_{2}) g sin (t + t_{2})) : (A1)$$

Here \div $p \xrightarrow{p} (=2)$ is the renormalized frequency of the (underdamped, $< 2!_0$) oscillator.

The other quantities needed for calculating q^2 (t) arise from the evaluation of ReS_{cl} (Eq. (24)). In contrast to C_{00} , which has to be evaluated numerically, they can be given in closed form:

$$L_{00} = m (=2 + 2 \cot(2 t))$$
 (A2)

$$L_{t0} = \frac{m \div e^{t=2}}{\sin(\div t)}$$
(A3)

The exponential increase of L_{t0} cancels that of C_{00} when calculating the width in Eq. (29). However, if the damping rate is set to zero, C_{00} still grows beyond all bounds while L_{t0} remains bounded.

Appendix B: DENSITY MATRIX PROPAGATOR FROM THE WIGNER DENSITY EVOLUTION

Using the classical Langevin equation (30), the kernel J which relates the reduced density matrix of a linear damped system at time t to that at time 0 can be found in a way which is physically more transparent than the corresponding derivation using path-integrals (see Eqs. (1), (2), (18),(24) and (25)). One first solves for the time-evolution of the classical phase space density (i.e. the Wigner density) under the action of friction and the Gaussian random force F (). Starting from a peak located in phase space, the phase space density evolves into a two-dimensional Gaussian distribution, whose covariances are related to the correlator of the force. This gives the propagator J^{W} of the Wigner density, which only needs to be Fourier transformed with respect to the momenta in order to obtain the density matrix propagator J, expressed via center-of-mass and difference coordinates R and r:

$$Z \qquad J(R_{t};r_{t}R_{0};r_{0};t) = dp_{t}dp_{0} e^{i(p_{t}r_{t} p_{0}r_{0})}J^{W}(R_{t};p_{t}R_{0};p_{0};t):$$
(B1)

Here we show how the propagator J for the density matrix of a free damped particle subject to the Ohmic bath may be obtained in this way. Everything works the same for the damped oscillator, only the resulting expressions are slightly more lengthy.

The propagator J^{W} (\mathbb{R}_{t} ; p_{t} ; p_{0} ; p_{0} ;t) of the Wigner density is found by solving the classical equations of motion for R and p for a given initial condition (\mathbb{R}_{0} ; p_{0}), taking into account friction and the action of the force F:

$$\frac{dp}{dt} = p + F(t)$$
(B2)

$$\frac{dR}{dt} = \frac{p}{m}$$
(B3)

This yields the solutions

$$p_t = p_0 e^{-t} + p \tag{B4}$$

$$R_t = R_0 + \frac{P_0}{2} (1 e^t) + R;$$
 (B5)

where $_{p}$ and $_{R}$ are given as integrals over the force F ():

$$A_{p} = \int_{0}^{Z_{t}} ds e^{-(t-s)} F(s)$$
(B6)

$$_{R} = \frac{1}{2} \int_{0}^{2} ds (1 e^{(t s)}) F(s) :$$
 (B7)

Since F () is a Gaussian random process, $_{p}$ and $_{R}$ are Gaussian random variables as well. Therefore, the phase space density evolving out of (R R₀) (p p) is a two-dimensional Gaussian distribution in phase space (R;p):

$$J^{W} (R_{t}; p_{t}; p_{0}; p_{0}; t) = (R_{t} R_{t} R_{t}) (p_{0}; p_{0}; t) = (R_{t} R_{t} R_{t}) (p_{0}; p_{0}; t) = (B8)$$

The average values p_t and R_t may be read off from eqs. (B4) and (B5). J^W has to be Fourier transformed with respect to p_t and p_0 in order to arrive at the density matrix propagator $J(R_t; r_t \mathfrak{R}_0; r_0; t)$ (see Eq. (B1)). In order to do this, we express J^W as a Gaussian density in terms of $p_t; p_0$, for fixed $R_t; R_0$:

$$J^{W} / \exp \frac{1}{2} P^{t}K^{1} P ;$$
 (B9)

with

$$P = \begin{array}{cc} P_t & P_t \\ P_0 & P_0 \end{array}$$
(B10)

and the covariance matrix

$$K = \frac{p_{t}^{2} \quad h \ p \ p_{t}}{h \ p \ p_{t}} : \qquad (B11)$$

Making use of (B4) and (B5), the average values and the deviations of p_t and p_0 are found to be given by:

$$p_t = (R_t \quad R_0) \quad (B12)$$

$$p_0 = e^{\tau} (R_t R_0)$$
 (B13)

$$p_{\rm e} = {}_{\rm p} {}_{\rm R} \tag{B14}$$

$$\mathbf{p} = \mathbf{e}^{\mathsf{T}}_{\mathsf{R}} \tag{B15}$$

$$\frac{1}{e^{t}}$$
(B16)

 p_0 and p_t are the momenta at time t and 0 which the particle must have if it is to go from R_0 to R_t in time t, provided no fluctuating force acts. p_t and p_0 are the deviations from these values needed to compensate the effects of F ().

Quadratic completion in the exponent immediately yields the result of the Fourier integration over p_t and p_0 , which is the desired density matrix propagator J:

$$\begin{array}{c} J(\mathcal{R}_{t};r_{t}\mathcal{R}_{0};r_{0};t) \neq \\ \text{exp i}(\mathcal{R}_{t} \ \mathcal{R}_{0}) (r_{t} \ e^{t}r_{0}) \quad \frac{1}{2} D \\ \left(r_{t} \ \mathcal{R}_{t} \ \mathcal{R}_{0}\right)^{2} (B17) \end{array}$$

This reproduces the result given in Ref. 18 (or Ref. 7, App. E). The prefactor can be determined from the normalization condition, Eq. (20), and only depends on the time t The term in angular brackets still has to be averaged over the force F (). The resulting real part of the exponent equals S_T evaluated along the semiclassical paths (compare the general structure given in Eq. (25)). In terms of F (), tpand po read explicitly:

$$p_{t} = \int_{a}^{Z_{t}} ds \frac{e^{s}}{e^{t}} \frac{1}{1} F (s)$$
(B18)

$$p_0 = \int_{0}^{\Delta_t} ds \frac{e^s}{e^t} \frac{e^t}{1} F$$
 (s): (B19)

Note that

$$p_{e} \quad p_{e} = \int_{0}^{Z_{t}} ds F(s) : \qquad (B20)$$

The averages p_1^2 , p_0^2 and h p pito be evaluated in Eq. (B17) contain time-integrals over the force correlator hF (s₁)F (s₂)i At zero temperature, these integrals lead to terms growing logarithmically in time, which are characteristic for the free particle coupled to an Ohmic bath (compare the discussion in the main text, sec. VI).

- Electronic address: Florian.Marquardt@unibas.ch
- ¹ R. P. Feynman and F. L. Vernon, Ann. Phys. (N.Y.) **24**, 118 (1963).
- ² R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, (McGraw-Hill, New York, 1965).
- ³ B. L. Altshuler, A. G. Aronov, and D. E. Khmelnitsky, J. Phys. C Solid State **15**, 7367 (1982).
- ⁴ A. Stern, Y. Aharonov, and Y. Imry, Phys. Rev. A **41**, 3436 (1990).
- ⁵ D. Cohen, Phys. Rev. E 55, 1422 (1997); D. Cohen, J. Phys. A 31, 8199 (1998).
- ⁶ D. S. Golubev and A. Zaikin, Phys. Rev. B **59**, 9195 (1999).
- ⁷ D. S. Golubev and A. Zaikin, Phys. Rev. B **62**, 14061 (2000).
- ⁸ D. S. Golubev and A. Zaikin, Physica B **280**, 453 (2000).
- ⁹ S. Chakravarty and A. Schmid, Phys. Rep. **140**, 195 (1986).
- ¹⁰ I. L. Aleiner, B. L. Altshuler, and M. E. Gershenson, Waves in Random Media 9, 201 (1999).
- ¹¹ M. Vavilov and V. Ambegaokar, cond-mat/9902127 (1999).
- ¹² D. Cohen and Y. Imry, Phys. Rev. B **59**, 11143 (1999).
- ¹³ T. R. Kirckpatrick and D. Belitz, Phys. Rev. B 65, 195123 (2002); Comment by D. S. Golubev, A. Zaikin and G. Schön in cond-mat/0111527, and reply in cond-mat/0112063.
- ¹⁴ Note that the impurity average of S_R (which corresponds to a properly weighted average over all possible pairs of trajectories) vanishes *in general*, simply because S_R [\vec{q} ; $\vec{q}^<$] = S_R [$\vec{q}^<$; $\vec{q}^<$].

- ¹⁵ A. O. Caldeira and A. J. Leggett, Phys. Rev. A **31**, 1059 (1985).
- ¹⁶ D. Loss and K. Mullen, Phys. Rev. B **43**, 13252 (1991).
- ¹⁷ K. E. Nagaev and M. Büttiker, Europhys. Lett. 58, 475 (2002).
- ¹⁸ A. O. Caldeira and A. J. Leggett, Physica **121**A, 587 (1983).
- ¹⁹ V. Hakim and V. Ambegaokar, Phys. Rev. A **32**, 423 (1985).
- ²⁰ Note that the damping constant used here corresponds to
 2 in Ref. 18, starting from their Eq. (6.8).
- ²¹ P. W. Milonni, The Quantum Vacuum, (Academic Press, San Diego, 1994).
- ²² U. Weiss, Quantum Dissipative Systems, (World Scientific, Singapore, 2000).
- ²³ P. Schramm and H. Grabert, Journal of Stat. Phys. 49, 767 (1987).
- ²⁴ P. Mohanty, E. M. Q. Jariwala, and R. A. Webb, Phys. Rev. Lett. **78**, 3366 (1997).
- ²⁵ P. Cedraschi, V. V. Ponomarenko, and M. Büttiker, Phys. Rev. Lett. 84, 346 (2000).
- ²⁶ F. Marquardt and C. Bruder, Phys. Rev. B **65**, 125315 (2002).
- ²⁷ G. Seelig and M. Büttiker, Phys. Rev. B **64**, 245313 (2001).
- ²⁸ F. Guinea, Phys. Rev. B 65, 205317 (2002).
- ²⁹ D. S. Golubev, C. P. Herrero, and A. D. Zaikin, condmat/0205549.