

Creation of a dipolar superfluid in optical lattices

B. Damski^{1,2}, L. Santos¹, E. Timmermann³, M. Lewenstein¹, S. Kotochigova⁴, P. Julienne⁴ and P. Zoller⁵

(1) Institut für Theoretische Physik, Universität Hannover, D-30167 Hannover, Germany

(2) Instytut Fizyki, Uniwersytet Jagielloński, Reymonta 4, PL-30 059 Kraków, Poland

(3) Institut für Quantenoptik, Universität Hannover, D-30167 Hannover, Germany

(4) National Institute of Standards and Technology, Gaithersburg, MD 20899, USA.

(5) Institut für Theoretische Physik, Universität Innsbruck, A-6020 Innsbruck, Austria

We show that by loading a Bose-Einstein condensate (BEC) of two different atomic species into an optical lattice, it is possible to achieve a Mott-insulator phase with exactly one atom of each species per lattice site. A subsequent photo-association leads to the formation of one heteronuclear molecule with a large electric dipole moment, at each lattice site. The melting of such dipolar Mott-insulator creates a dipolar superfluid, and eventually a dipolar molecular BEC.

The physics of strongly correlated systems in ultra-cold trapped quantum gases has attracted a growing interest recently. The experimental progress on the trapping and cooling techniques, and the control of the interaction potentials by means of Feshbach resonances allow to analyze situations beyond the validity of the mean-field approach. In particular, the possibility to achieve 1D bosonic gases opens the pathways toward a gas of impenetrable bosons [1]. Also, a rapidly rotating trapped bosonic gas should exhibit effects similar to the fractional quantum Hall effect [2]. The field of strongly correlated atomic systems concerns also ultra-cold Bose gases with large scattering lengths [3,4], and the Mott-insulator (MI) to superfluid (SF) phase transition [5] in cold bosonic gases in optical lattices [6,7]. The latter has been recently observed in the remarkable experiment [8].

The physics of ultra-cold molecular gases is also a very active research area. Cooling and trapping of molecules can be achieved by means of buffer-gas techniques [9], by employing the dipole moments of polar molecules [10], and by means of photo-association of already existing ultra-cold atomic gases [11]. Jaksch et al. [12] have recently proposed that a bosonic atomic gas, which is in a MI phase with exactly two atoms per site in an optical lattice, can be used to obtain via photo-association a superfluid gas of homonuclear molecules, and eventually a molecular BEC.

The influence of dipole-dipole forces on the properties of ultra-cold gases has also drawn a considerable attention. It has been shown that these forces significantly modify the ground state and excitations of trapped dipolar BEC's [13-18]. In addition, since dipole-dipole interactions can be quite strong relative to the short-range (contact) interactions, dipolar particles are considered to be promising candidates for the implementation of fast and robust quantum computing schemes [19,20]. Several sources of cold dipolar bosons have been proposed, including atoms with large magnetic moments [4], and externally induced electric dipoles [13,15,19]. Perhaps

the most promising perspective in this sense is provided by ultra-cold gases of polar molecules with large electric dipole moments, which could eventually dominate the physics of the system [15,16].

This Letter is devoted to the analysis of the generation of ultra-cold polar molecules in an optical lattice. We consider the loading of the lattice by a superfluid of two different atomic species, and analyze the transition into a MI phase with exactly one atom of each species per site. This transition is followed by the formation of polar dimers by photo-association, or adiabatic passage. Finally, the quantum melting of the molecular Mott phase into a polar molecular superfluid takes place.

We consider a miscible gas of two trapped atomic bosonic species at very low temperature T . Although our calculations could in principle be employed for arbitrary atoms, we have assumed in the calculations below the particular mixture of ^{41}K and ^{87}Rb , whose simultaneous Bose-Einstein condensation has been recently realized [21]. We additionally consider that the binary gas is confined in a 2D optical lattice, similarly to the 3D one-component case recently observed in Ref. [8]. The transverse dimension is considered as harmonically confined. The physics of the system is governed by the two-species generalization of the Bose-Hubbard (BH) model, described by the Hamiltonian:

$$H = \sum_{\langle i,j \rangle} [J_a a_i^\dagger a_j + J_b b_i^\dagger b_j] + U_{ab} \sum_i n_{ai} n_{bi} + \frac{1}{2} \sum_i [U_{0a} n_{ai} (n_{ai} - 1) + U_{0b} n_{bi} (n_{bi} - 1)]; \quad (1)$$

where a_i, b_i are the annihilation operators of ^{41}K and ^{87}Rb atoms at the lattice site i , which occupy a state described by the Wannier functions $w_a(\mathbf{r} - \mathbf{r}_i), w_b(\mathbf{r} - \mathbf{r}_i)$ of the lowest energy band, localized on this site. This implies that energies involved in the system are small compared to the excitation energies to the second band. We denote the position of the local min-

in um of the optical potential as r_i , and the number operator of K (Rb) atoms at the site i as $n_{ai} = a_{ai}^\dagger a_i$ ($n_{bi} = b_{bi}^\dagger b_i$). In Eq. (1) only the nearest-neighbor tunneling is considered, which is described by the parameter $J_{a,b} = \int d^3r \psi_{a,b}^\dagger(r-r_i) \left[\frac{\hbar^2}{2m} r^2 + V_L(r) \right] \psi_{a,b}(r-r_j) d^3r$, where j and i are the indices of neighboring sites, and $V_L(r) = \sum_{x,y} V^0 \cos^2(k \cdot r)$ is the optical lattice potential (with wavevector k) which we consider to be the same for both species. The optical potential for K and Rb is almost the same for the detunings corresponding to the wavelength 1064nm of a Nd:Yag laser [22]. The same species interact via a short range pseudopotential, which is described by the coefficients $U_{0a}; U_{0b}$: $U_{0a,b} = 4 \frac{\hbar^2}{m_{a,b}} a_{K,Rb}^2 \int d^3r \psi_{a,b}^\dagger(r-r_i) \psi_{a,b}(r-r_j) d^3r$, where a_K, m_K (a_{Rb}, m_{Rb}) are the scattering length and atomic mass of potassium (rubidium). The interactions between different species are characterized by the coefficient $U_{ab} = 2 \frac{\hbar^2}{m_{ab}} a_{KRb} \int d^3r \psi_a(r-r_i) \psi_b(r-r_j) d^3r$, where a_{KRb} is the K-Rb scattering length, and m_{ab} is the reduced mass for K and Rb atoms. The scattering length a_{KRb} has been estimated recently to be of the order of 8.6nm [21], that is the repulsive interactions between K and Rb are stronger than those between K-K and Rb-Rb, which are characterized by $a_K = 3.17\text{nm}$ and $a_{Rb} = 5.24\text{nm}$, respectively. As a result, it is energetically favorable for K and Rb particles to remain immiscible, i.e. to stay at different lattice sites, rather than in the same ones. This situation can be however modified by means of Feshbach resonances, which can strongly modify the value of a_{KRb} . We have thus assumed that the value of a_{KRb} can be reduced to $0.5 \frac{a_K a_{Rb}}{a_K + a_{Rb}}$. In that case the two components acquire a miscible phase.

In the 2D calculations presented below, we have considered square lattices up to 80×80 sites, with periodic boundary conditions. In the first stage, we demonstrate the possibility of performing a transition from a state with two immiscible superfluids to a Mott insulator phase with exactly one Rb and one K atom per site. To this aim we first obtain the ground state of the system using a variational Gutzwiller ansatz [6].

We approximate the ground-state wave function by $\prod_{i,j} j_{MF} i = \prod_i j_i i$, where the product is over all lattice sites. The functions $j_i i$ for each site are expressed in the Fock basis, $j_i i = \sum_{n,m} \frac{1}{n!m!} f_{n,m}^{(i)} j_n m i$, where n, m indicate the occupation number for the K and Rb atoms, respectively. For numerical reasons we have assumed that maximally 3 atoms of each species can be present at any lattice site, and checked the validity of this assumption self-consistently. The coefficients $f_{n,m}^{(i)}$ are found by minimization of $\langle \prod_{i,j} j_{MF} i | H | \prod_{i,j} j_{MF} i \rangle$, where the chemical potentials $\mu_a; \mu_b$ were chosen so that the mean numbers of K, Rb atoms were equal to the number of lattice sites.

The ground state for weak lattice potentials corresponds to the coexistence of superfluids for both atomic

species, and it was used as the initial condition for subsequent time-dependent calculations. This state is characterized by a non-vanishing value of the superfluid order parameters $\langle j_{ai} i, j_{bi} i \rangle$ (Fig. 1). To drive the system from superfluid to Mott phase we slowly increase the lattice potential. This process can be described using a dynamical Gutzwiller method, as discussed in Ref. [12]. To this aim one allows the coefficients $f_{n,m}^{(i)}$ to be time dependent and to follow the dynamics derived from the time dependent variational principle. The resulting equation becomes:

$$i \dot{f}_{n,m}^{(i)} = \frac{U_{0a}}{2} n(n+1) + \frac{U_{0b}}{2} m(m+1) f_{n,m}^{(i)} + J_a \frac{a_i}{h} \frac{p}{n+1} f_{n+1,m}^{(i)} + \frac{a_i}{h} \frac{p}{n} f_{n-1,m}^{(i)} + J_b \frac{b_i}{h} \frac{p}{m+1} f_{n,m+1}^{(i)} + \frac{b_i}{h} \frac{p}{m} f_{n,m-1}^{(i)} + U_{ab} n m f_{n,m}^{(i)}; \quad (2)$$

where

$$\frac{a_i}{h} = \sum_{\langle ij \rangle} h_{MF} j_{MF} i; \quad \frac{b_i}{h} = \sum_{\langle ij \rangle} h_{MF} j_{MF} i;$$

When the lattice potential increases, the system follows the changes quasi-adiabatically, and the two species enter sequentially the Mott phase with one atom per site. As shown in Fig. 1 this occurs first for the heavier Rb atom at $t = 0.75s$, and then at $t = 1.5s$ for K atoms. The ratio $U_{0a} = J_a$ ($U_{0b} = J_b$) at the transition point equals approximately 23.5 (26), which interestingly compares very well with the expected value, 23.2, for a 2D single-component gas. Therefore the transition point is rather unaffected by the interactions between different species.

At the end of the process each site contains one Rb and one K atom, with a relative atom number fluctuations less than 3%. This, however, requires a sufficiently slow modification of the lattice potential, typically within a time scale of 2 seconds for a lattice wavelength $\lambda = 1064\text{nm}$. Note that the time scale is provided by the inverse of the recoil energy of the lattice, and therefore it scales as λ^{-2} . We note also at this point that a transition 3 times faster leads to about two times larger number fluctuation in the final state. We want to stress that during the whole time evolution the mean number of particles was constant, and equal to the number of lattice sites, which gives us the most complete analogy, within the Gutzwiller approach, to the experimental situation.

Having obtained a Mott phase with one Rb and one K atom per site, a heteronuclear molecule can be generated at each lattice site. One possible method would be two-color Raman photoassociation to make ground state $1 +$ or $3 +$ dimers. Since heteronuclear species do not have gerade/ungerade symmetry, levels of either symmetry could be formed via intermediate excited states of $\Sigma = 0^+$ or 1 symmetry, where labels projection of

electronic plus spin angular momentum on the molecular axis. Since the states excited at long range typically have mixed singlet and triplet character, photoassociation to ground 1^+ levels is even possible for collision of two doubly spin-polarized atoms, which collide via the ground 3^+ potential. The $1=R^6$ dependence of the excited molecular potential for the case of heteronuclear molecules, in contrast to the $1=R^3$ dependence for the homonuclear case, leads to excitation at relatively short internuclear separation R . This difference may lead to reduced Franck-Condon factors for the first excitation step but enhanced factors for the second de-excitation step [24]. In any case, these can be worked out once the potential energy curves of the K-Rb dimer are better known. A second possible method of making ground state molecules is available for heteronuclear dimers, namely, use of a pulse of microwave radiation to directly associate two colliding atoms in a one-color transition from a scattering state to a bound dimer state. However, the dipole matrix elements within the 3^+ state are small, so this will need fairly high microwave power. Either method will result in an excited vibrational level not far below the dissociation limit. It should be possible to use a succession of Raman pulses to transfer the population from such a level to the ground $v = 0$ vibrational level of the 1^+ or 3^+ state, in a manner similar to that described by Ref. [12]; microwave/infrared pulses may also be feasible for this heteronuclear species. The $3^+ v = 0$ level has a small dipole moment of 0.0046 atomic units (1 atomic unit is $8.478 \cdot 10^{-30}$ Cm), whereas it is much larger, 0.30 atomic units, for the $1^+ v = 0$ level [23]. In the latter case, special care should be taken to prevent the heating due to black-body radiation, which could be avoided by an appropriate shielding.

After the creation of the K-Rb dimers, the system becomes single-component. However, since the molecules have a large dipole moment and are considered to be oriented by an external electric field (100 V/cm), we have to include the dipole-dipole interactions in our calculations. The corresponding Hamiltonian reads [7]:

$$H = J \sum_{\langle i,j \rangle} b_i^\dagger b_j + \frac{1}{2} U_0 \sum_i n_i (n_i - 1) + \frac{1}{2} U_1 \sum_{\langle i,j \rangle} n_i n_j + \frac{1}{2} U_2 \sum_{\langle \langle i,j \rangle \rangle} n_i n_j + \dots; \quad (3)$$

where b_i is the annihilation operator of a heteronuclear molecule at the lattice site i . The number operator n_i , the Wannier basis $w(\mathbf{r} - \mathbf{r}_i)$ and the tunneling coefficient J for molecules are defined similarly as in Eq. (1). However, the interaction part significantly differs from that of Eq. (1). The coefficients U are defined as follows: $U = \int d^3r d^3r' \langle \mathbf{r} - \mathbf{r}_i | \hat{V}_{\text{int}}(\mathbf{r} - \mathbf{r}') | \mathbf{r} - \mathbf{r}_j \rangle \int d^3r d^3r' \langle \mathbf{r} - \mathbf{r}_j | \hat{V}_{\text{int}}(\mathbf{r} - \mathbf{r}') | \mathbf{r} - \mathbf{r}_i \rangle$, where $\langle \mathbf{r} - \mathbf{r}_i | \hat{V}_{\text{int}}(\mathbf{r} - \mathbf{r}') | \mathbf{r} - \mathbf{r}_j \rangle$ is the dimensionless intersite distance. In particular, U_0 determines the on-site interactions, U_1 the nearest-neighbor interactions, U_2 the interactions

between the next-nearest neighbors, etc. Consequently, the respective summations in Eq. (3) must be carried out over appropriate pairs of sites which are marked by $\langle \dots \rangle$ for the nearest neighbors, $\langle \langle \dots \rangle \rangle$ for the next-nearest neighbors, etc. In the 2D calculations presented below, we have taken into account $\frac{1}{2}$ interactions with up to 4 neighbors ($U_1 = 1$, $U_2 = \frac{1}{2}$, $U_3 = 2$, $U_4 = \frac{1}{5}$), since the effects of interactions of a longer range are negligible. In the case of polarized dipoles the interaction potential is

$$V_{\text{int}} = d^2 \frac{1 - 3 \cos^2 \theta}{r^3} + \frac{4 \hbar^2 a}{M} (r - r^0); \quad (4)$$

where the first part of Eq. (4) provides the dipole-dipole interaction characterized by the dipole moment d and the angle θ between the dipole direction and the vector $\mathbf{r} - \mathbf{r}^0$. The second part of Eq. (4) represents the short-range interactions given by the s-wave scattering length a and the molecular mass M . We use the dipole moment as 0.3 a.u. [23], and assume the s-wave scattering length of K-Rb molecule as $a = \frac{1}{2} \sqrt{\frac{\hbar^2}{\mu k_{\text{Rb}}}}$ [25].




FIG. 1. Dynamical transition from the SF phase of ^{41}K (solid line) and ^{87}Rb (dashed line) atoms to the MI phase. The upper plot shows the value of the order parameters $\langle n_i \rangle$, $\langle n_i^2 \rangle$ (constant for all lattice sites) for both species, while the lower one depicts the variance $\text{Var}(n) = \langle n_i^2 \rangle - \langle n_i \rangle^2$ of the on-site occupation. The lattice potential was changed as $V_0(t) = V_{\text{SF}} + (V_{\text{MI}} - V_{\text{SF}})(t - t_0)^3 \exp(1 - (t - t_0)^3)$, where $V_{\text{SF}} = 4$; $V_{\text{MI}} = 19$ in units of recoil energy of ^{41}K atoms, $t_0 = 1.8\text{ s}$ for the case of lattice wavelength 1064 nm.

We have checked that the MI phase of these heteronuclear molecules is indeed the ground state of the system for a mean lattice filling factor equal to 1. We employ this state as the initial condition for our dynamical calculations of the melting of the Mott phase towards a molecular superfluid. It is, however, interesting to note, that by itself the MI phase of dipolar particles offers promising perspectives as a quantum computation device [20]. The melting dynamics is analyzed by employing again a dynamical Gutzwiller ansatz, but with the Hamiltonian (3). During the dynamics we consider a first stage in which we reduce the lattice trapping potential up to 4 in

recoil energy units of K ^{87}Rb molecules. This first stage allows us to reduce the value of $U_0=J$, but the system remains still in the MI state mostly due to the large contribution of the repulsive dipolar on-site interactions to the U_0 coefficient. This process can be performed in a relatively fast time scale compared to the total duration of the melting process, since the MI gap guarantees the adiabaticity at this stage.




FIG. 2. Melting of ^{41}K ^{87}Rb dimers initially in the MI phase. The plot shows the time evolution of the molecular superfluid order parameter $\rho_{\text{bf},i}$ (solid line), which is the same for all lattice sites. The dashed line refers to static calculations of the ground state of the dipolar molecules placed in the lattice. Note a very sharp transition from the MI phase to the SF one, that occurs at $t \approx 0.2\text{s}$. The frequency of the transverse confinement was changed as $\omega_z(t) = \omega_{\text{MI}} + (\omega_{\text{SF}} - \omega_{\text{MI}})t/t_0$, where $\omega_{\text{MI}} = 2\pi \times 8\text{kHz}$, $\omega_{\text{SF}} = 2\pi \times 6\text{kHz}$, and $t_0 = 0.65\text{s}$.

Later on, we have observed that the dynamics can be better controlled by modifying the transverse confinement. As shown in Ref. [7,15], the properties of the dipolar gas strongly depend on the aspect ratio of the on-site wavefunction. In particular, by reducing the transverse confinement, the on-site wavefunction becomes more elongated in transverse direction. As a consequence, U_0 decreases [7], and the system enters into the superfluid phase. Fig. 2 shows this process. Due to the non-completely adiabatic evolution the gas does not exactly follow the expected stationary result, although very clearly a superfluid molecular phase is accomplished. The ratio $U_0=J \approx 23.2$ at which the transition from MI to superfluid equals surprisingly to the value expected for a 2D single-component gas without a permanent dipole moment. As shown in Fig. 2 this dynamical transition into the superfluid phase occurs for realistic parameters in a time scale of 1s. This transition can be realized, 50 times faster, although the dynamics of the superfluid order parameter is in such case not as smooth as in Fig. 2.

In this Letter we have analyzed the formation of a superfluid of polar molecules. We have considered the loading of a BEC of ^{87}Rb and ^{41}K atomic species into an optical lattice. The modification of the lattice potential produces the dynamical transition into a MI phase

with only one atom of each species per site. The atoms can then be photo-associated on-site. Once the polar molecules are created, a molecular MI is formed. By reducing the lattice potential and modifying the transverse confinement, such MI evolves into a molecular superfluid within a feasible time scale. Eventually, a molecular BEC is formed. The mechanism studied in this Letter, provides a perspective towards a dipolar BEC.

We acknowledge support from the Alexander von Humboldt Stiftung, the Deutsche Forschungsgemeinschaft, EU RTN Network "Cold Quantum Gases", the Polish KBN (grant no 2 P 03B 124 22), ESF PESC Program BEC2000+ and the US Office of Naval Research. We thank G. Birkel, G. V. Shlyapnikov and E. Tiesinga for discussions.

-
- [1] E. H. Lieb and W. Liniger, Phys. Rev. 130, 1605 (1963); M. Girardeau, J. Math. Phys. 1, 516 (1960).
 - [2] B. Paredes et al, Phys. Rev. Lett. 87, 010402 (2001).
 - [3] S. C. Cowell et al, Phys. Rev. Lett. 88, 210403 (2002).
 - [4] N. R. Claussen et al, Phys. Rev. Lett. 89, 010401 (2002).
 - [5] See e.g., M. P. A. Fisher et al, Phys. Rev. B 40, 546 (1989).
 - [6] D. Jaksch et al, Phys. Rev. Lett. 81, 3108 (1998).
 - [7] K. Goral, L. Santos and M. Lewenstein, Phys. Rev. Lett. 88, 170406 (2002).
 - [8] M. Greiner et al, Nature (London) 415 39 (2002).
 - [9] J. D. Weinstein et al, Nature (London) 395, 148 (1998).
 - [10] H. L. Bethlem et al, Nature (London) 406, 491 (2000).
 - [11] R. Wynar et al, Science 287, 1016 (2000).
 - [12] D. Jaksch et al, Phys. Rev. Lett. 89, 040402 (2002).
 - [13] S. Yi and L. You, Phys. Rev. A 61, 041604 (2000).
 - [14] K. Goral, K. Rzaewski, and T. Pfau, ibid. 61, 051601 (2000).
 - [15] L. Santos et al, Phys. Rev. Lett. 85, 1791 (2000).
 - [16] K. Goral and L. Santos, cond-mat/0203542.
 - [17] H. Pu, W. Zhang, and P. M. Dyre, Phys. Rev. Lett. 87, 140405 (2001).
 - [18] S. Giovanazzi, D. O'Dell, and G. Kurizki, Phys. Rev. Lett. 88, 130402 (2002).
 - [19] D. Jaksch et al, Phys. Rev. Lett. 85, 2208 (2000).
 - [20] G. K. Brennen et al, Phys. Rev. Lett. 82, 1060 (1999); G. K. Brennen et al, Phys. Rev. A 65, 022313 (2002); D. DeMille, Phys. Rev. Lett. 88, 067901 (2002).
 - [21] G. M. Odungo et al, cond-mat/0205485; G. M. Odungo et al, Science 294 1320 (2001).
 - [22] R. Grimm, M. Weidemüller, and Y. B. Ovchinnikov, Adv. At. Mol. Opt. Phys. 42, 95 (2000).
 - [23] P. Julienne and S. Kotochigova, to be published.
 - [24] Also, the on-site densities (10^{15} cm^{-3}) are 10000 times larger than in traditional photoassociation experiments.
 - [25] The value of a is not known, but does not affect our results, since it can always be compensated by transverse confinement, i.e. dipolar forces.