

# Feshbach-resonance-induced atomic filamentation and quantum pair correlation in atom-laser-beam propagation

Wei-ping Zhang, Chris P. Search, Han Pu, Pierre Meystre and Ewan M. Wright  
Optical Sciences Center, University of Arizona, Tucson, AZ 85721

(Dated: April 14, 2024)

We study the propagation of an atom-laser beam through a spatial region with a magnetic field tuned to a Feshbach resonance. Tuning the magnetic field below the resonance produces an effective focusing Kerr medium that causes a modulational instability of the atomic beam. Under appropriate circumstances, this results in beam breakup and filamentation seeded by quasi-particle fluctuations, and in the generation of correlated atomic pairs.

PACS numbers: PACS numbers: 03.75.Fi, 05.30.Fk, 67.40.Db

It has been recognized for quite some time that the two-body interactions that govern the dynamics of atomic Bose-Einstein condensates (BECs), and which are expressed most simply in terms of the s-wave scattering length  $a$ , play a role for matter waves analogous to that of a cubic (Kerr) nonlinearity in optics. This was illustrated in a number of experiments that include matter-wave four-wave mixing [1], the generation of dark [2] and bright [3] atomic solitons, coherent matter-wave amplification [4], and the generation of quantum-correlated atomic pairs [5]. It has also been demonstrated that the strength of this nonlinearity can be varied, and its sign changed, using in particular Feshbach resonances [6, 7]. In the language of nonlinear optics, changing the sign of the scattering length amounts to switching from a defocusing (for repulsive interactions,  $a > 0$ ) to a focusing (for attractive interactions,  $a < 0$ ) nonlinear medium. The tunability of the scattering length offers a degree of flexibility of considerable interest in the study of the many-body properties of atomic condensates, as well as for nonlinear atom-optics applications.

As pointed out by Timmermans et al. [8], caution is in order when considering the dynamics of a condensate in the vicinity of a Feshbach resonance. In the atomic systems under consideration, they correspond to a situation where the total energy of two colliding atoms equals the energy of a bound molecular state. In the vicinity of the resonance, it is essential to properly account for the molecular dynamics. Indeed, it has recently been unambiguously established experimentally that a proper description of the system requires that one accounts for the coherent coupling between the atomic and molecular field, resulting in the onset of atom-molecule coherence in BECs [9]. The presence of the molecular state is also predicted to lead to "resonant superfluidity" in quantum-degenerate atomic Fermi systems [10].

Far from resonance, however, it is possible to adiabatically eliminate the molecular field from the system's dynamics, in which case the atomic condensate is subject to an effective, tunable scattering length that describes the binary atomic collisions. The goal of this letter is to illustrate how this tunable scattering length can be ex-

ploited to manipulate and control the propagation of a continuous-wave atom-laser beam. The specific geometry that we consider consists of an atom-laser beam propagating through a region where a spatially dependent magnetic field is applied to switch the nature of the binary collisions from being repulsive, or defocusing, to attractive, or focusing. This results in a spatial change of the interatomic interaction similar to the presence of a Kerr medium in nonlinear optics, albeit with some important differences. We show that under appropriate circumstances, the quasi-particle fluctuations in the initial atom-laser beam can seed the filamentation of the atom-laser beam, with the generation of pair-correlated atoms in spatial sidebands. This is similar to the amplification of squeezed light by parametric processes in a nonlinear optical Kerr medium, but with the important difference that in the present situation it is not necessary to start from a squeezed input.

We recall that the generation of correlated atomic pairs from BECs has recently been studied by a number of authors [5, 11]. In these works, correlated atomic pairs are generated either from vacuum fluctuations [11], or grow via four-wave mixing from the combined effects of a seed beam and vacuum fluctuations [5]. In contrast, the proposed scheme involves the generation of a pair-correlated atomic source from quasi-particle vacuum fluctuations, with an atom-laser beam as the sole pump beam.

A Feshbach resonance is a scattering resonance where two particles in an open channel collide to form a bound state in the closed channel. In the case of quantum-degenerate atomic fields, such as Bose-Einstein condensates or atom-laser beams, the bound state in the closed channel is a coherent molecular state. The dynamical equations describing the atom-molecule coherent coupling can be written as [8]

$$\begin{aligned} i\hbar \frac{\partial \hat{a}}{\partial t} &= \frac{\hbar^2 \mathbf{r}^2}{2m} + U_a \hat{a}^\dagger \hat{a} + U_{am} \hat{a}^\dagger \hat{m} + \hat{m}^\dagger \hat{a} \\ i\hbar \frac{\partial \hat{m}}{\partial t} &= \frac{\hbar^2 \mathbf{r}^2}{4m} + U_m \hat{m}^\dagger \hat{m} + U_{am} \hat{a}^\dagger \hat{a} + \hat{a}^\dagger \hat{m} \end{aligned}$$

$$+ \hat{a}_a^2; \quad (1)$$

Here  $\hat{a}_{a,m}$  are the atomic and molecular field operators,  $m$  is the atomic mass,  $U_a = (4\hbar^2/m)a_a$ ,  $U_m = (2\hbar^2/m)a_m$ , and  $U_{am} = (3\hbar^2/2m)a_{am}$  are the atom-atom, molecule-molecule, and atom-molecule binary interaction strengths, with  $a_a$ ,  $a_m$ , and  $a_{am}$  the corresponding scattering lengths, and  $h$  is the coupling strength the Feshbach resonance. The parameter  $\epsilon$  is the energy difference between the molecular state and the atomic state.

For large resonance detunings  $j \gg \epsilon$ , it is possible to adiabatically eliminate the molecular field, leading to the effective Heisenberg equation of motion

$$i\hbar \frac{\partial \hat{a}_a}{\partial t} = \frac{\hbar^2 \nabla^2}{2m} + h \hat{a}_a^\dagger \hat{a}_a \hat{a}_a; \quad (2)$$

for the atomic Schrödinger field. In this expression, the effective coupling constant  $h = U_a/2$  accounts for the change of the strength of the atomic binary collisions resulting from the Feshbach resonance.

We now proceed to study the propagation of an atom laser through a spatially-dependent effective nonlinearity, which for simplicity we take to be of the form

$$h(z) = U_a/2 \Theta(z);$$

where  $\Theta(z)$  is a Heaviside step function. In particular, we consider the situation in which the binary collisions change from repulsive for  $z < 0$ , to attractive for  $z > 0$ . The continuous wave atom laser beam is incident from  $z < 0$ , and we describe the incident atomic field as a coherent beam of Bose-condensed atoms propagating along the  $z$ -axis with wave vector  $k_0 \hat{z} = (m v_0/\hbar)\hat{z}$ . We also ignore any atoms that are back reflected at the interface: If we take account of the fact that the magnetic field actually varies over a length  $L$  around  $z = 0$ , then using a WKB analysis we find that back reflections are negligible if  $(2\hbar^2/mk_0)(2\hbar^2/mj)n_0 = L$ ,  $v_0^2 = 2m$  where  $n_0$  is the density of atoms in the condensate.

An important distinction between this and the generic situation considered in nonlinear optics, where the beam is incident from vacuum, is that the incident atomic field is self-interacting. Hence, a proper description must include the quasiparticle vacuum fluctuations propagating together with the coherent BEC component. As such, a realistic atom laser beam consists of the sum of a moving condensed component, described classically, and quantum fluctuations [12],

$$\hat{a}_a(r;t) = \sqrt{\frac{P}{n_0}} + \hat{\alpha}(r;t) e^{i(k_0 k_n)z} e^{i\epsilon_0 t}; \quad z < 0; \quad (3)$$

where  $\sqrt{\frac{P}{n_0}}$  describes the coherent BEC component (taken real without loss of generality),  $\hbar^2 k_0^2 = 2m$ , and  $k_n = n_0 v_0$  is a nonlinear phase shift. The noise operator  $\hat{\alpha}(r;t)$  obeys bosonic commutation relations.

From the corresponding Heisenberg equation of motion (2), one observes that the binary collisions contribute a term of the form

$$i\hbar \frac{\partial \hat{\alpha}(r;t)}{\partial t} = U_a n_0 \hat{\alpha}(r;t) + \hat{\alpha}^\dagger(r;t)$$

to the evolution of the noise operator  $\hat{\alpha}(r;t)$ . This form of evolution is quite familiar in quantum optics, where it is known to lead to the generation of squeezed states. Thus, the repulsive self-interaction of the incident atom laser beam for  $z < 0$  is sufficient to provide initial squeezed state vacuum fluctuations at the input port of the (focusing) nonlinear medium  $z = 0$ .

We proceed by expanding the noise operator  $\hat{\alpha}(r;t)$  in terms of plane-waves as

$$\hat{\alpha}(r;t) = \int \frac{d\mathbf{q}}{(2\pi)^2} \int \frac{dk_z}{2\pi} \hat{a}_0(\mathbf{q};k_z;t) e^{i\mathbf{q} \cdot (\mathbf{r} - \mathbf{v}_0 t)} + \int \frac{d\mathbf{q}}{(2\pi)^2} \int \frac{dk_z}{2\pi} \hat{b}_0(\mathbf{q};k_z;t) e^{i\mathbf{q} \cdot (\mathbf{r} - \mathbf{v}_0 t)} e^{i\mathbf{q} \cdot \mathbf{r}_0}; \quad (4)$$

where  $\mathbf{q}$  is the transverse wave vector and the operators  $\hat{a}_0(\mathbf{q};k_z;t)$  and  $\hat{b}_0(\mathbf{q};k_z;t) = \hat{a}_0(\mathbf{q};-k_z;t)$  describe forward and backward propagating modes in the rest frame of the atomic beam, respectively.

In the incident region with repulsive interactions,  $z < 0$ , it is convenient to express the small quantum fluctuations about the condensate wave function in terms of quasiparticles characterized by bosonic annihilation operators  $\hat{\alpha}(\mathbf{q};k_z;t)$  and  $\hat{\beta}(\mathbf{q};k_z;t) = \hat{\alpha}^\dagger(-\mathbf{q};-k_z;t)$ , given by [12]

$$\begin{aligned} \hat{a}_0(\mathbf{q};k_z;t) &= u_k \hat{\alpha}(\mathbf{q};k_z;t) + v_k \hat{\beta}(\mathbf{q};k_z;t); \\ \hat{b}_0(\mathbf{q};k_z;t) &= u_k \hat{\alpha}(-\mathbf{q};k_z;t) + v_k \hat{\beta}(-\mathbf{q};k_z;t); \end{aligned}$$

where

$$u_k^2 = v_k^2 + 1 = \frac{1}{2} \left( \frac{k + \hbar n_0}{E_k} + 1 \right);$$

with  $k = \sqrt{\frac{P}{n_0} + k_z^2}$ ,  $E_k = \hbar^2 k^2/2m$ , and  $E_k = \frac{\hbar^2 k^2}{2} + 2k\hbar n_0$  being the quasiparticle energy associated with  $\hat{\alpha}(\mathbf{q};k_z;t)$  and  $\hat{\beta}(\mathbf{q};k_z;t)$ .

We are now in a position to study the propagation of the atom laser beam as it enters the region of Feshbach resonance,  $z > 0$ . Specifically, in this paper we focus on the initiation of the filamentation process before appreciable depletion. For this purpose, it is sufficient to consider a linearized perturbation theory in the slowly varying amplitude approximation. For  $z > 0$  we decompose the atomic field operator in the same manner as Eq. (3) with the substitution of  $\hat{\alpha}(r;t)$  in place of  $\hat{\alpha}(r;t)$  for the fluctuation operator. Furthermore, analogous to Eq. (4), we also decompose the noise operator  $\hat{\alpha}(r;t)$  in terms of quasiplane waves,

$$\hat{\alpha}(r;t) = \int \frac{d\mathbf{q}}{(2\pi)^2} \hat{\alpha}(\mathbf{q};t) e^{i\mathbf{q} \cdot \mathbf{r}}; \quad (5)$$

where

$$\hat{a}_q(z;t) = \int_0^{z/v_0} dk_z \hat{a}(q;k_z;z) e^{ik_z(z-v_0t)} + \hat{b}(q;k_z;z) e^{ik_z(z-v_0t)} \quad (6)$$

Here we allow for a slow  $z$ -dependence of the operators  $\hat{a}$  and  $\hat{b}$ . This approach, familiar from quantum optics,

allows one to account for the possible amplification of the quantum noise. Spatial variation of the quasiparticle operators was ignored in the region  $z < 0$  due to the absence of sideband gain for repulsive binary collisions.

Substituting Eq. (5) into Eq. (2) yields pairs of coupled operator equations that describe the linearized spatial evolution of the condensate sidebands,

$$i \frac{\partial}{\partial T} \hat{a}(q;k_z;z) = \frac{q + n_0}{n_0} \hat{a}(q;k_z;z) + \frac{n_0}{q + n_0} \hat{b}^*(q;k_z;z) \quad (7)$$

where  $T = t + z/v_0$ . Assuming a solution to Eq. (7) of the form  $e^{gT}$  and matching the field operator solutions in the two regions at  $z = 0$  we obtain finally

$$\begin{aligned} \hat{a}(q;k_z;z) &= \int_0^n U(q;z) u_k + V(q;z) v_k \hat{a}(q;k_z) + [U(q;z) v_k + V(q;z) u_k] \hat{b}^*(q;k_z) ; \\ \hat{b}(q;k_z;z) &= \int_0^n V(q;z) u_k + U(q;z) v_k \hat{a}^*(q;k_z) + [V(q;z) v_k + U(q;z) u_k] \hat{b}(q;k_z) ; \end{aligned} \quad (8)$$

where the coefficients  $U$  and  $V$  are given by

$$\begin{aligned} U(q;z) &= \cosh g \frac{z}{v_0} - \frac{i}{g} (q + n_0) \sinh g \frac{z}{v_0} ; \\ V(q;z) &= i \frac{n_0}{g} \sinh g \frac{z}{v_0} ; \end{aligned} \quad (9)$$

with the gain parameter

$$g = \frac{q}{(n_0)^2 - (q + n_0)^2} \quad (10)$$

Without loss of generality we have taken  $t = 0$  in order to focus on how the filamentation develops with propagation distance.

We observe that the linearized solution of Eqs. (8) can exhibit gain in the case of a focusing Feshbach resonance,  $\mu < 0$ . Specifically, the gain coefficient  $g$  is real for  $q < 2m j \hbar_0 = \hbar$ , with a maximum,  $q_{max} = j \hbar_0$ , for the transverse wavevector

$$q_{max} = \frac{\hbar}{2m j \hbar_0} \quad (11)$$

Physically, this growth of a range of coupled pairs of spatial sidebands  $q$  around the condensate corresponds to a transverse modulational instability with a cone of unstable transverse momenta peaked around  $|j| = q_{max}$ . If allowed to grow this instability will ultimately lead to the breakup of the incident laser beam into transverse structures or filaments similarly to the situation in nonlinear optics [13, 14]. If instead of using a plane-wave condensate we consider a beam with widths  $w_{xy}$  in the  $x$  and  $y$  transverse directions, we can eliminate the instability in the  $y$ -direction, for example, by choosing  $q_{max} w_y < 1$ ;  $q_{max} w_x \ll 1$ , so that the most unstable

wavevector cannot be supported in the  $y$ -direction. In that case the cone of unstable wavevectors becomes a pair of sidebands that are peaked at  $q_{max}$  along the  $x$ -direction. This quasi-one-dimensional geometry may be better to isolate correlated atomic pairs.

This modulational instability is intimately related to the collapse of  $^{85}\text{Rb}$  condensates seen in recent experiments [7]. In these experiments, stable repulsive BEC's with densities of  $n_0 \sim 10^{22} \text{ cm}^{-3}$  are produced in the presence of a magnetic field tuned close to a Feshbach resonance ( $B = 160 \text{ G}$ ) and then the magnetic field is tuned above  $167 \text{ G}$  where the interaction becomes attractive. Far from the resonance the interactions are attractive with a background scattering length of  $a_{bg} = 23.85 \text{ nm}$  [9]. Using these parameters gives a  $q_{max} = 7.75 \times 10^6 \text{ m}^{-1}$  or a transverse velocity of  $0.6 \text{ cm/s}$  for the sideband atoms.

Proceeding with the example of a plane-wave condensate, to quantify the statistical properties of the generated atomic sidebands, we evaluate the field correlation and intensity correlation functions for  $\hat{a}_q(z)$  defined in Eq. (6). Assuming that the quantum fluctuations in the incident atom laser beam are in the quasi-particle vacuum state  $|j_0\rangle$ , i.e.,  $\hat{a}|j_0\rangle = \hat{b}|j_0\rangle = 0$ , and with Eqs. (8) and (9) we find,

$$\langle \hat{a}_q^y \hat{a}_{q^0}^y \rangle = \langle q \rangle_{q,q^0} \quad (12)$$

$$\langle \hat{a}_q^y \hat{a}_q^y \hat{a}_{q^0}^y \hat{a}_{q^0}^y \rangle = \langle q \rangle \langle q \rangle + \mathcal{D} \langle q \rangle^2_{q,q^0} \quad (13)$$

where all the operators are evaluated at the same longitudinal distance  $z$  and in Eq. (13), we have taken  $q \neq q^0$ . Physically,  $\langle q \rangle$  is the probability density in momentum

space of finding an atom with a transverse momentum  $\hbar q$  after propagating a distance  $z$  in the Feshbach region, and  $\mathcal{Q}(q)^2$  is the joint probability that if one atom

is detected with momentum  $\hbar q$  a second one will be detected with opposite momentum  $-\hbar q$ . They are given by,

$$\begin{aligned} \rho(q) &= 2 \int_0^{z/2} dk_z \left[ v_k^2 + \frac{2n_0}{g^2} (u_k^2 + v_k^2) \sinh^2 \left( g \frac{z}{v_0} - \frac{2n_0}{g^2} u_k v_k \sinh^2 \left( g \frac{z}{v_0} \right) \right) \right]; \\ \mathcal{Q}(q) &= 2 \int_0^{z/2} dk_z \left[ u_k v_k + \frac{n_0}{g^2} (u_k^2 + v_k^2) \right] \frac{2}{g^2} u_k v_k \left[ u_k \sinh^2 \left( g \frac{z}{v_0} + \frac{ig}{2} \sinh \left( 2g \frac{z}{v_0} \right) \right) \right]; \end{aligned} \quad (14)$$




FIG. 1: Correlation spectrum of the correlated atomic pairs calculated from Eq. (14). Due to the divergence at large wave number, the upper limit of  $k_z$  is cut off at  $k_s = m c_0 = \hbar$  the wave number corresponding to the Bogoliubov sound velocity. Here the wave numbers, energy and length are in units of  $k_s$ ,  $\hbar^2 k_s^2 = (2m)$  and  $1/k_s$ , respectively. The parameters used in the calculation are  $v_0 = 5c_0$ ,  $n_0 = 0.5$ . The plots show that both  $\rho$  and  $\mathcal{Q}$  grow fastest near  $q_{\text{ax}} = 1 = \frac{1}{2}$  [see Eq. (11)].

with  $u_q = q + n_0$ .

As an illustration we assume that atoms are detected after having propagated a distance  $z$  in the Feshbach region such that the linearized analysis is still valid. The rate at which atoms are detected at an angle  $\theta = \tan^{-1}(q/k_0)$  relative to the  $z$ -axis will be proportional to  $\rho(q)$  while the rate of coincidence detections at  $+\theta$  and  $-\theta$  will be proportional to  $\mathcal{Q}(q)^2$ . The corresponding probability densities are plotted in Fig. 1, from which we see that, as expected physically, with increasing propagation distance  $z$  the most probable transverse wavevector for detecting either an on-axis atom or a correlated atomic

pair at  $q$  is increasingly  $q = q_{\text{ax}}$ . These results show that the transverse modulational instability can indeed be an effective source of correlated atomic pairs.

In conclusion, we have studied the propagation of an atom laser beam through a Feshbach resonance region where the effective interaction between atoms is tuned from being repulsive to being attractive. The attractive interaction is equivalent to a self-focusing Kerr medium for the atom laser beam which can generate quantum-pair-correlated atomic filament beams by a modulational instability. This opens up new window to create non-classical ultracold atomic sources by coherent control of the atomic interactions. The quantum-pair-correlated atomic beam may have potential applications in both fundamental research and applications in quantum information and computation.

This work is supported in part by the US Office of Naval Research under Contract No. 14-91-J1205 and No. N00014-99-1-0806, by the National Science Foundation under Grant No. PHY 98-01099, by the US Army Research Office, and by the Joint Services Optics Program.

- 
- [1] L. Deng et al, Nature 398, 218 (1999).
  - [2] S. Burger et al, Phys. Rev. Lett. 83, 5198 (1999); J. Denzschlag et al, Science 287, 97 (2000).
  - [3] K. E. Strecker et al, Nature 417, 150 (2002).
  - [4] S. Inouye et al, Nature 402, 641 (1999); M. Kozuma et al, Science 286, 2309 (1999).
  - [5] J. M. Vogels, K. Xu and W. Ketterle, Phys. Rev. Lett. 89, 020401 (2002).
  - [6] S. Inouye et al, Nature 392, 151 (1998).
  - [7] S. L. Comish et al, Phys. Rev. Lett. 85, 1795 (2000).
  - [8] E. T. In ’t Hout et al, Phys. Rev. Lett. 83, 2691 (1999).
  - [9] E. A. Donley et al, Nature 417, 529 (2002).
  - [10] M. Holland et al, Phys. Rev. Lett. 87, 120406 (2001).
  - [11] H. Pu and P. Meystre, Phys. Rev. Lett. 85, 3987 (2000); L.-M. Duan et al, Phys. Rev. Lett. 85, 3991 (2000); A. Vardi and M. G. Moore, Phys. Rev. Lett. 89, 090403 (2002).
  - [12] A. L. Fetter, Ann. Phys. 70, 67 (1972).
  - [13] A. J. Campbell et al, Appl. Phys. Lett. 23, 628 (1973); ibid. 24, 178 (1974).

- [14] E . M . Nagasako, R . W . Boyd, and G . S . Agarwal, Phys. Rev. A 55, 1412 (1997).