

## Erratum : Low-dimensional Bose gases [Phys. Rev. A 66, 013615 (2002)]

U. AlKhawaja<sup>1</sup>, J.O. Andersen<sup>1</sup>, N.P. Proukakis<sup>1,2</sup>, and H.T.C. Stoof<sup>1</sup><sup>1</sup>Institute for Theoretical Physics,

Utrecht University, Leuvenlaan 4, 3584 CE Utrecht, The Netherlands

<sup>2</sup>Foundation for Research and Technology Hellas,

Institute of Electronic Structure and Laser, P.O. Box 1527, Heraklion 71 110, Crete, Greece

(November 10, 2021)

PACS numbers: 03.75.Fi, 67.40.-w, 32.80.Pj

In Ref [1], we presented a modified Popov theory that is capable of calculating the quasicondensate density  $n_0$  in arbitrary dimensions and at all temperatures. When a condensate exists, i.e., in three dimensions at sufficiently low temperatures and in two dimensions at zero temperature, the condensate density  $n_c$  is defined by the off-diagonal long-range order of the one-particle density matrix. In particular, in the modified Popov theory we have that

$$n_c = \lim_{x \rightarrow 1} n_0 e^{\frac{1}{2} [\hat{h}(x) - \hat{h}(0)]^2 i}; \quad (1)$$

where  $n_0$  satisfies the equations

$$n = n_0 + \frac{1}{V} \sum_k \frac{k^d h^d}{2h^d k^d} + \frac{n_0 T^{2B}(2)}{2k + 2} + \frac{k^d N(h^d k^d)}{h^d k^d}; \quad (2)$$

$$= (2n - n_0) T^{2B}(2); \quad (3)$$

and the phase fluctuations are determined by

$$\hat{h}(x) - \hat{h}(0) i = \frac{T^{2B}(2)}{V} \sum_k \frac{1}{2h^d k^d} [1 + 2N(h^d k^d)] \frac{1}{2k + 2} e^{ik \cdot x}; \quad (4)$$

Here  $n$  is the total density of the gas,  $T^{2B}(2)$  is the two-body matrix at energy  $2$ , and  $\mu$  is the chemical potential.  $k = \hbar^2 k^2 = 2m$  and  $h^d k^d = \frac{2}{k} + 2n_0 T^{2B}(2) k^{1-2}$ . In Ref. [1], we assumed that the exponent in Eq. (1) vanishes in the limit  $x \rightarrow 1$  and thus found that at zero temperature the depletion of the condensate in two dimensions equals

$$\frac{n - n_0}{n} = \frac{1}{4} (1 - \ln 2) T^{2B}(2); \quad (5)$$

whereas in three dimensions it equals

$$\frac{n - n_0}{n} = \frac{32}{3} \left( \frac{p}{2} \right)^r \frac{1}{na^3}; \quad (6)$$

where  $a$  is the scattering length and we have used  $T^{2B}(2) = 4 \hbar^2 a^2 m$ . A careful analysis shows, however, that for a consistent calculation of the quantum depletion the above assumption is incorrect and we have to take the contribution from the phase fluctuations into account. In the limit  $x \rightarrow 1$ , we obtain in two and three dimensions

$$[\hat{h}(x) - \hat{h}(0)]^2 i = \frac{1}{2} (\ln 2) T^{2B}(2); \quad (7)$$

and

$$[\hat{h}(x) - \hat{h}(0)]^2 i = 2 \left( \frac{p}{2} \right)^r \frac{1}{na^3}; \quad (8)$$

respectively. Since we consider a weakly-interacting Bose gas, the right-hand sides of Eqs. (7)–(8) are small and we can expand the exponents in Eq. (1) to first order. Combining this with Eqs. (5)–(6), we finally obtain

$$\frac{n - n_c}{n} = \frac{1}{4} T^{2B}(2); \quad (9)$$

in two dimensions and

$$\frac{n - n_c}{n} = \frac{32}{3} \frac{r}{na^3}; \quad (10)$$

in three dimensions. These results are in perfect agreement with the results obtained from Popov theory.

We would like to point out that none of the other results presented in Ref. [1] are affected by the above, since the theory is always applied in a regime where the gas parameter is small and the subtlety discussed here can be safely ignored.

#### ACKNOWLEDGMENTS

The authors would like to thank Stefano Giorgini for a useful discussion that led us to this observation.

---

[1] U. AlKhawaja, J.O. Andersen, N.P. Proukakis, and H.T.C. Stoof, Phys. Rev. A 66, 013615 (2002).