

Non-universal critical Casimir force in confined ^4He near the superfluid transition

X. S. Chen^{a,b}, V. Dohm^b

^a Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100080, China

^b Institut für Theoretische Physik, Technische Hochschule Aachen, D-52056 Aachen, Germany

Abstract

We present the results of a one-loop calculation of the effect of a van der Waals type interaction potential $\propto x^{-d}$ on the critical Casimir force and specific heat of confined ^4He near the superfluid transition. We consider a ^4He film of thickness L . In the region $L \ll \xi$ (correlation length) we find that the van der Waals interaction causes a leading non-universal non-scaling contribution of $O(\xi^{-2}L^{-d})$ to the critical temperature dependence of the Casimir force above T^* that dominates the universal scaling contribution $\propto e^{L/\xi}$ predicted by earlier theories. For the specific heat we find subleading non-scaling contributions of $O(L^{-1})$ and $O(L^{-d})$.

Key words: critical phenomena, finite-size effects, helium 4, Casimir force

In the last decade the critical Casimir effect in ^4He films has been an interesting topic of theoretical [1,2,3] and experimental [4] research. On the basis of field-theoretic calculations for purely short-range interaction it was predicted that the critical temperature dependence of the Casimir force is governed by a universal scaling function $X(L/\xi)$ where L is the film thickness and $\xi = \xi_0 t^{-\nu}$, $t = (T - T^*)/T^*$, is the bulk correlation length. The universal behavior for large L/ξ was predicted to be $\propto e^{L/\xi}$. Here we investigate the question which finite-size effect arises from van der Waals type interactions. They do not change the leading universal bulk critical behavior of the free energy and specific heat. The finite-size effects of these interactions were discussed in Refs. [1,2,3] where it was assumed that they yield only non-singular (temperature-independent) contributions. Here we shall show that these interactions yield a leading singular non-scaling temperature dependence of the Casimir force that is non-universal and dominates the exponential scaling contribution above T^* for large L/ξ . We also present predictions for subleading non-scaling contributions to the critical specific heat of ^4He films above T^* that may be non-negligible in the analysis of experimental data.

The present paper is a continuation of recent work where leading non-scaling cutoff and lattice effects [5,6,7,8] and non-scaling effects of van der Waals type interactions [8,9,10] have been found both for the finite-size susceptibility and for the bulk order-parameter correlation function. These effects were not taken into account in the previous theory [1,2,3] that was based on the usual ^4He Hamiltonian in d dimensions

$$H = \int d^d x \left[\frac{1}{2} (\nabla \phi)^2 + \frac{1}{2} (r' \phi)^2 + u_0 (\phi'^2)^2 \right] \quad (1)$$

for the n -component field $\phi(\mathbf{x})$. Here the short-range interaction term $(r' \phi)^2$ has the Fourier representation $k^2 \phi_{\mathbf{k}}' \phi_{-\mathbf{k}}'$. In the following we assume the existence of an additional subleading long-range interaction term $\propto k^{-2} \phi_{\mathbf{k}}' \phi_{-\mathbf{k}}'$ with $2 < \alpha < 4$ which corresponds to a van der Waals type spatial interaction potential $U(x) \propto x^{-d}$. For the case of ^4He ($n = 2$) we shall consider Im geometry with Dirichlet boundary conditions.

We have performed a one-loop calculation of the critical Casimir force $F = F_s + F_{ns}$. For the singular part F_s we have found

$$F_s(L; b) = -b L^{d+2} B(L/\xi) + L^{-d} X(L/\xi) \quad (2)$$

where the last term L^{-d} is the universal scaling part due to the short-range interaction. For $L \ll \xi$ this scaling part is dominant. For large $L = \xi$, however, it becomes exponentially small, $X(L = \xi) \sim e^{-L/\xi} \ll 1$.

The non-universal non-scaling term b due to the van der Waals type interaction reads for $L = \xi$ & 1

$$B(L = \xi) = (d-3+\epsilon)(L = \xi)^{-(d-3+\epsilon)}(L = \xi)^0(L = \xi); \quad (3)$$

$$(L = \xi) = \frac{1}{2}(2-\epsilon)^{-4} \int_{(L = \xi)^2}^{\xi} dx [1 + x \frac{\partial}{\partial x}] e^{-x}; \quad (4)$$

$$e(x) = \int_0^{\xi} dy y^{(2-\epsilon)/2} e^{-xy/4} \frac{r-d-1}{y} \tilde{W}_1(y) \quad (5)$$

where $(z;x) = x^{-z} \int_0^x dt e^{-t} t^{z-1} = \frac{R_1}{0} \int_0^x dt e^{-t} t^{z-1}$ is the incomplete Gamma function and

$$\tilde{W}_1(y) = \frac{r-d-1}{y} \frac{1}{2} \sum_{n=1}^{\infty} \exp\left(-\frac{y}{4} n^2\right); \quad (6)$$

The large $L = \xi$ behavior is $(L = \xi)^{-2} L^{-2}$. Thus the leading critical temperature dependence $F_s(\xi; L; b) \sim L^{-2} L^{-d}$ is nonuniversal for $L \ll \xi$ and dominates the exponentially small scaling part.

Corresponding non-universal non-scaling effects exist for the free energy [8] which implies a significant restriction for the range of validity of the universal Priman-Fisher scaling form [11,12].

We have also calculated the finite-size effect of the van der Waals interaction on the specific heat in one-loop order for $L \ll \xi$ above T_c . The result reads

$$C(\xi; L; b) = C_b + 2C_{\text{surface}} + 1 + b \mathcal{F}(\xi; d)^2 \frac{1}{L} + C + 2ba_0^2 \mathcal{F}(\xi; d)^6 L^{-d} \quad (7)$$

with $a_0 = (r_0 - r_c)/t$ where C_b, C_{surface} and $C = L^{-d} g(L = \xi)$ are the known bulk, surface and finite-size parts of the specific heat due to the short range interaction [13]. For the relation between a_0 and r_0 see [14]. The amplitudes of the non-universal non-scaling terms b due to the van der Waals type interaction are

$$\mathcal{F}(\xi; d) = \frac{5}{2} \frac{d}{2} \int_0^{\xi} ds s^{(3-d)/2} e^{-s} f(s) \quad (8)$$

with

$$f(s) = \frac{1}{4} (2-\epsilon) \frac{2}{2} \frac{1}{s} + \frac{1}{2} \frac{1}{s}$$

$$+ \frac{1}{2} \frac{1}{s}; \quad (9)$$

and

$$\mathcal{F}(\xi; d) = \frac{(2-\epsilon)}{2^{d+2}} \frac{d+2}{d-2} \frac{d+2}{2} (d+2) \quad (10)$$

where ζ is Riemann's zeta function.

The next step of our theory will be to specify the nonuniversal strength b and the exponent of the van der Waals type interaction in H_e . Since these parameters are as yet unknown the range of applicability of the earlier predictions for the universal finite-size scaling functions of the Casimir force and of the specific heat above T_c is not yet established.

Finally we note that our one-loop results b were obtained by a calculation at the Gaussian level (with $u_0 = 0; r_{0c} = 0$ and at infinite cutoff) where part of the non-Gaussian (higher-loop) effects were taken into account by the replacement $r_0 \rightarrow r_0 + \frac{1}{2}$. It remains to be seen to what extent a full renormalization-group treatment with $u_0 > 0$ at the two-loop level confirms this procedure. In this sense our predictions of the temperature dependence of the terms b in Eqs. (2) and (7) should be considered only as preliminary. A full renormalization-group treatment will be given elsewhere.

Acknowledgements

We acknowledge support by DLR and by NASA under grant numbers 50WM 9911 and 1226553.

References

- [1] M. Knecht, S. Dietrich, Phys. Rev. Lett. 66 (1991) 345; Phys. Rev. A 46 (1992) 1886; 46 (1992) 1922; J. Low Temp. Phys. 89 (1992) 145.
- [2] M. Knecht, The Casimir Effect in Critical Systems (World Scientific, Singapore, 1994).
- [3] M. Knecht, J. Phys. Cond. Mat. 11 (1999) R 391.
- [4] R. Garcia, M. H. W. Chan, Phys. Rev. Lett. 83 (1999) 1187; 88 (2002) 086101.
- [5] X. S. Chen, V. Dohm, Eur. Phys. J. B 7 (1999) 183.
- [6] X. S. Chen, V. Dohm, Eur. Phys. J. B 10 (1999) 687.
- [7] X. S. Chen, V. Dohm, Eur. Phys. J. B 15 (2000) 283.
- [8] X. S. Chen, V. Dohm, Phys. Rev. E 66 (2002) 016102.
- [9] D. Dantchev, J. Rudnick, Eur. Phys. J. B 21 (2001) 251.
- [10] D. Dantchev, Eur. Phys. J. B 23 (2001) 211.
- [11] V. Priman, M. E. Fisher, Phys. Rev. B 30 (1984) 322.

- [12] V. Privman, Phys. Rev. B 38 (1988) 9261.
- [13] V. Dohm, Phys. Scripta 49 (1993) 46.
- [14] R. Schloms, V. Dohm, Phys. Rev. B 42 (1990) 6142.