

Comment on “Universal Fluctuations in Correlated systems”

In [1] the “universal” probability density function (PDF) of various global measures x from correlated equilibrium and non-equilibrium physics-inspired models was shown (figure 2 of [1]) to be in reasonable agreement with $\Pi(y) = K(e^{y-e^y})^a$, where $y = b(x-s)$. This Bramwell-Holdsworth Pinton (BHP) PDF with shape parameter $a = \pi/2$, was contrasted with the Fisher-Tippett-Gumbel (FTG) PDF where $a = 1$, one of 3 possible asymptotes for the maxima of uncorrelated random variables (r.v.’s) [2]. In [1] the BHP PDF was also seen to describe the distribution of maxima ξ_{max}^L from L trials of length N vectors ξ of correlated r.v.’s, generated by $\xi = \mathbf{M}\chi$. The elements of χ were i.i.d. exponentially distributed, and \mathbf{M} was “an $N \times N$ matrix with random but fixed elements”. The applicability of the BHP PDF to all the considered processes in [1] was inferred to be due to “finite size, *strong correlations* (our italics) and self-similarity”. In this Comment we show that this inference is not supported by the extremal behaviour of ξ_{max}^L : because 1) while a suitable choice of N ensures that ξ_{max}^L falls on the BHP curve, other equally arbitrary values of N give maxima which don’t; and 2) ξ is in fact not strongly correlated, and the PDF of ξ_{max}^L change little for decorrelated ξ . We thus suggest that a differs from its FTG asymptote of 1 here largely due to slow convergence of maxima of the nearly Gaussian ξ .

In [1], “random” implies “i.i.d Gaussian, mean 0, variance 1” i.e. $P(M) = \frac{1}{\sqrt{2\pi}}e^{-M^2/2}$ and “random but fixed” implies \mathbf{M} has random elements held constant between trials. ξ is correlated but the fact that \mathbf{M} and χ are statistically independent means that for the simplest case of $N = 2$, $\xi_1 (= M_{11}\chi_1 + M_{12}\chi_2)$ has PDF $P(\xi_1) = \int dM_{11}dM_{12}d\chi_1d\chi_2\delta(\xi_1 - M_{11}\chi_1 - M_{12}\chi_2)P(M_{11})P(M_{12})P(\chi_1)P(\chi_2)$. Taking exponential $P(\chi) = \lambda e^{-\lambda\chi}$ as in [1] we generalise to $N \times N$ matrix \mathbf{M}

so $P(\xi) = \frac{1}{2\pi} \int_{-\infty}^{\infty} I(k)^N e^{ik\xi} dk$. Here $I(k)^N = \lambda^N \left[\int_{-\infty}^{\infty} \frac{P(M)}{\lambda + ikM} dM \right]^N$ is the generating

function of $P(\xi)$. We then have $I(k) = \sqrt{\frac{\pi}{2}} \frac{\lambda}{k} e^{\lambda^2/2k^2} (1 - \text{erf}(\lambda/\sqrt{2}k))$.

The maxima of ξ are dominated by the tail behaviour of $P(\xi)$, which from a small k expansion of I is seen to be Gaussian $\sim (\lambda/\sqrt{N}) \exp(-\xi^2\lambda^2/4N)$. The maxima of ξ will thus be in the basin of attraction of the FTG distribution. The asymptotic shape parameter a corresponding to $L, N \rightarrow \infty$ cannot be changed by short range dependence [2], but one expects corrections to a for finite N, L because of the logarithmically slow convergence of the maxima of Gaussians, and, sometimes, a change in

effective degrees of freedom N due to correlation [2].

Figure 1 shows the PDF of ξ_{max}^L for $L = 10^5$ trials in 2 cases: $N = 40$ and 250, and for comparison the PDF of $\chi_{max}^L = \max(\chi^L)$, when $N = 40$, to illustrate the expected rapid convergence of χ_{max}^L to the FTG curve. Convergence of ξ_{max}^L however slows with increasing N . By $N = 40$, convergence of ξ_{max}^L is noticeably slower than χ_{max}^L and the PDF follows the BHP curve; while for $N = 250$ even slower convergence in χ_{max}^L gives obvious deviation from the BHP form resembling that seen in figure 2 of [1]. For $N = 40$ the autocorrelation function of χ (not shown) drops fast by the first lag. Weak (less than ± 0.2) long ranged oscillations *are* seen at all lags. Despite this, decorrelating ξ by shuffling the $L \times N$ matrix before taking the maximum of ξ changes the PDF of its maxima in a relatively minor way (see Figure 1), confirming that correlation cannot explain most of the change from the FTG PDF here. Our result implies that, even though subsequent results may show that the BHP curve *can* result from strong correlation, it *need not*.

NWW thanks Mervyn Freeman, Clare Watt, Pat Espy and David Stephenson for valuable suggestions.




FIG. 1. $L = 10^5$. Middle plot shows PDF of ξ_{max}^L (circles) compared with PDF of maxima of decorrelated process (crosses), $N = 40$. Top plot (stars) is PDF of maxima χ_{max}^L of exponentially distributed r.v.’s, $N = 40$. Lowest plot (circles) shows ξ_{max}^L in $N = 250$ case. Also shown on each plot are the BHP curve (solid line), and the $a = 1$ FTG plot (dashed line). Displacements of $10^{-1.5}$ and 10^{-3} for the middle and lower plots, respectively, were applied for clarity.

N. W. Watkins

British Antarctic Survey, Cambridge, CB3 0ET, U.K.

S. C. Chapman and G. Rowlands

University of Warwick, Coventry, CV4 7AL, U.K.

Received February 5, 2022

PACS numbers: 02.50.-r, 05.40.-a

- [1] S. T. Bramwell *et al.*, Phys. Rev. Lett. **84**, 3744 (2000).
- [2] S. Coles, Extreme Value Theory and Applications, 1999; <http://www.stats.bris.ac.uk/~masgc/extremes/exnewnotes2.ps>