

Non-universality in Ising models with four spin interaction.

V. Mastropietro

Dipartimento di Matematica,

Università di Roma "Tor Vergata"

Via della Ricerca Scientifica, I-00133, Roma

Abstract. We consider two bidimensional classical Ising models, coupled by a weak interaction bilinear in the energy densities of the two systems; the model contains, as limiting cases, the Ashkin-Teller and the Eight-vertex models for certain values of their parameters. We write the energy-energy correlations and the specific heat as Grassmann integrals formally describing Dirac 1+1 dimensional interacting massive fermions on a lattice, and an expansion based on Renormalization Group is written for them, convergent up to temperatures very close to the critical temperature for small coupling. The asymptotic behaviour is determined by critical indices which are continuous functions of the coupling.

Key words Coupled Ising models, non-universality, Renormalization Group, fermions, critical indices.

1. Introduction and main results

1.1 Motivations. It is well known that the partition function and many correlations of the $d = 2$ classical Ising model can be written as Grassmann integrals formally describing a system of $d = 1 + 1$ free fermions, see [H], [SML], [K], [MW], [S], [ID]. This mapping of the Ising model in terms of a fermionic theory is quite useful; as the corresponding action is quadratic in the Grassmann variables, it can be diagonalized by a Bogoliubov transformation so that the partition function and many observables can be exactly computed. However, even a slight perturbation of the Ising model makes the corresponding fermionic action non-quadratic. The relationship between spin Ising-like models and interacting fermionic models is deeply investigated in the literature; it is for instance claimed that models of coupled Ising models with an interaction quartic in the spins, like the Eight-vertex model, are in the same class of universality of models of interacting $d = 1 + 1$ fermions in the continuum, like the massive Thirring model or the massive Luttinger model, see for instance [N], [LP], [PB]. This relationship is however found under many approximations. In this paper we consider two Ising models coupled by a general interaction bilinear in the spin densities; the model contains as particular cases the Ashkin-Teller and Eight-vertex model for certain values of their parameters. We clarify the relation between this general spin model and systems of interacting fermions by writing the partition function and some correlations as Grassmann integrals with a formal action resembling but not coinciding with the action of the massive Thirring or Luttinger model; the main difference (in addition to the fact that the formal action describes fermions in a lattice, not in the continuum) is that it does not verify some special symmetries (like invariance under local gauge transformations) which are generally verified in models of interacting fermions and which play an important role in their analysis. The main interest in this representation in terms of Grassmann integrals is that we can apply the Renormalization Group techniques developed for interacting fermions (see for instance [BG] or [GM]) for writing a convergent perturbative expansion for the partition function and some correlations of the spin model we are considering. A naive power series expansion in the coupling is not expected to be convergent close to the critical temperature (the free and interacting model are not analytically close) and one needs a resummed or renormalized expansion which is provided by Renormalization Group.

1.2 Spin systems with quartic interactions. We consider two Ising models coupled via a four spin interaction bilinear in the energy densities of the two sublattices. Given $M \in 2\mathbb{Z}^2$ a square lattice with side M and periodic boundary condition, we call $x = (x; x_0)$ a site of M . If $x^{(1)} = 1$ and $x^{(2)} = 1$, we write the following Hamiltonian

$$H((^{(1)}; ^{(2)})) = H_I(^{(1)}) + H_I(^{(2)}) + V(^{(1)}; ^{(2)}) \quad (1.1)$$

where, if $\alpha = 1; 2$

$$H_I(\alpha) = \sum_{x, x_0=1}^M [J_1^{(\alpha)}(x, x_0) + J_2^{(\alpha)}(x, x_0+1)] \quad (1.2)$$

$$\begin{aligned} V(\alpha^{(1)}; \alpha^{(2)}) = & \sum_{x, x_0=1}^M f a [\binom{(1)}{x, x_0} \binom{(1)}{x+1, x_0} \binom{(2)}{x, x_0} \binom{(2)}{x+1, x_0} + \binom{(1)}{x, x_0} \binom{(1)}{x, x_0+1} \binom{(2)}{x, x_0} \binom{(2)}{x, x_0+1}] \\ & + b [\binom{(1)}{x, x_0} \binom{(1)}{x+1, x_0} \binom{(2)}{x, x_0} \binom{(2)}{x, x_0+1} + \binom{(1)}{x, x_0} \binom{(1)}{x, x_0+1} \binom{(2)}{x+1, x_0+1} \binom{(2)}{x, x_0+1}] \quad (1.3) \\ & + c \sum_{\alpha=1;2}^X [\binom{(\alpha)}{x, x_0} \binom{(\alpha)}{x+1, x_0} \binom{(\alpha)}{x, x_0} \binom{(\alpha)}{x, x_0+1} + \binom{(\alpha)}{x, x_0} \binom{(\alpha)}{x+1, x_0} \binom{(\alpha)}{x+1, x_0+1} \binom{(\alpha)}{x, x_0+1}] \lg \end{aligned}$$

The above Hamiltonian can be a model for a number of physical problems; for instance two magnetic spin planes coupled by an interplanar interaction quartic in the spins (in this interpretation $\alpha^{(1)}$ and $\alpha^{(2)}$ are the spin of the first or second plane). Moreover it contains, as limiting cases, two well known models, the Ashkin-Teller and the Eight-vertex model.

1.3 The Ashkin-Teller model. Ashkin and Teller introduced their model as a generalization of the Ising model to a four state system. Each site x is occupied by one of the four kinds of atom, said A, B, C, D , and two neighboring atoms interact with an energy ϵ_0 for AA, BB, CC, DD ; ϵ_1 for AB, CD , ϵ_2 for AC, BD , ϵ_3 for AD, BC . It is quite easy to express this model in terms of Ising spins, see [F]. One associates to each site of the lattice two spins $\alpha_x^{(1)}$ and $\alpha_x^{(2)}$, see Fig.1; then $\alpha_x^{(1)}; \alpha_x^{(2)} = (+; +)$ if there is an A atom at site x , $(+; -)$ if there is an atom B , $(-; +)$ if there is an atom C and $(-; -)$ if there is an atom D . Then the Hamiltonian is given by

$$H(\alpha^{(1)}; \alpha^{(2)}) = H_I(\alpha^{(1)}) + H_I(\alpha^{(2)}) + \sum_{x, x_0=1}^M [J_0^{(2)} \binom{(2)}{x, x_0} \binom{(2)}{x, x_0+1} \binom{(1)}{x, x_0} \binom{(1)}{x, x_0+1} + J_0^{(1)} \binom{(2)}{x, x_0} \binom{(2)}{x+1, x_0} \binom{(1)}{x, x_0} \binom{(1)}{x+1, x_0+1}] \quad (1.4)$$

where $H_I(\alpha^{(\alpha)})$ is given by (1.2) with $J_1^{(\alpha)} = J_2^{(\alpha)} = J^{(\alpha)}$ and

$$\begin{aligned} J^{(1)} &= (\epsilon_0 + \epsilon_1 - \epsilon_2 - \epsilon_3) = 4J & J^{(2)} &= (\epsilon_0 + \epsilon_2 - \epsilon_3 - \epsilon_1) = 4J \\ J^{(0)} &= (\epsilon_0 + \epsilon_3 - \epsilon_1 - \epsilon_2) = 4J & J_0 &= (\epsilon_0 + \epsilon_1 + \epsilon_2 + \epsilon_3) = 4J \end{aligned} \quad (1.5)$$

The model is not soluble, except in some particular case, and we refer to [Ba] chapt.12 and references therein for a review of the main results about it; it is of the form (1.1) with $b = c = 0$.

1.4 The Eight-vertex model. In such model one associates to each site of a square lattice a vertex with four arrows with different orientations, see [Ba] chapt. 10.

Among the sixteen possibilities, only eight kind of vertices are allowed, and an energy $\epsilon_i, i = 1; \dots; 8$, is associated to each of them. It was solved in [Ba] if $\epsilon_1 = \epsilon_2, \epsilon_3 = \epsilon_4, \epsilon_5 = \epsilon_6, \epsilon_7 = \epsilon_8$ and for suitable values of the ϵ_i it reduces to the ice-model, solved in [Li]. Such models are called vertex models and were introduced to describe crystal with hydrogen bonding. As explained in [Ba] or [W] they can be written in terms of the following Hamiltonian, see Fig. 2

$$H(\sigma^{(1)}; \sigma^{(2)}) = \sum_{x, x_0} [J_{x, x_0+1 \ x+1, x_0} + J^0_{x, x_0 \ x+1, x_0+1} + J^0_{x, x_0 \ x+1, x_0+1} + J^0_{x, x_0+1 \ x+1, x_0}] \quad (1.6)$$

with

$$\begin{aligned} \epsilon_1 = \epsilon_2 = J - J^0 - J^0, \quad \epsilon_3 = \epsilon_4 = J + J^0 - J^0 \\ \epsilon_5 = \epsilon_6 = J^0 - J + J^0, \quad \epsilon_7 = \epsilon_8 = J^0 + J + J^0 \end{aligned} \quad (1.7)$$

If $J^0 = 0$ the above Hamiltonian decouples in the Hamiltonian of two Ising models, involving spins located in two sublattices, one with $x + x_0$ equal to an even number and the other to an odd number. Each site of the sublattices is in the center of the unit square of the other one. Calling $\sigma^{(1)}$ and $\sigma^{(2)}$ the spin located in the first or the second sublattice, relabeling the spins and performing a rotation of $\frac{\pi}{4}$ the Hamiltonian can be written as, see Fig.3

$$H(\sigma^{(1)}; \sigma^{(2)}) = H_I(\sigma^{(1)}) + H_I(\sigma^{(2)})$$

$$J^0 \sum_{x, x_0=1}^M [\sigma^{(2)}_{x, x_0} \sigma^{(2)}_{x, x_0+1} \sigma^{(1)}_{x, x_0} \sigma^{(1)}_{x+1, x_0} + \sigma^{(1)}_{x, x_0} \sigma^{(1)}_{x, x_0+1} \sigma^{(2)}_{x+1, x_0+1} \sigma^{(2)}_{x, x_0+1}] \quad (1.8)$$

where

$$H_I(\sigma^{(i)}) = \sum_{x, x_0=1}^M J [\sigma^{(i)}_{x, x_0} \sigma^{(i)}_{x+1, x_0} + \sigma^{(i)}_{x, x_0} \sigma^{(i)}_{x, x_0+1}]$$

which is again of the form (1.1) with $a = c = 0$ (of course this identification is exact if boundary conditions in the Eight-vertex model are chosen properly).

1.5 Main results. We will consider in this paper the case

$$J_1^{(2)} = J_2^{(1)} = J_1^{(2)} = J_2^{(2)} = J \quad (1.9)$$

If $\epsilon = 0$ the problem reduces to the Ising model whose critical temperature is given by $\tanh \epsilon J_0 = \frac{1}{2} \frac{1}{1 - t_c}$ where $J = J_0$ and t_c is the temperature. The interaction changes in general the value of the critical temperature by terms $O(\epsilon)$, and it is instead technically convenient to fix its value still in correspondence of $t_c = \frac{1}{2} \frac{1}{1 - t_c}$, by choosing properly the molecular energy parameter J as a function of ϵ ; in this way the critical temperature of the system with $\epsilon = 0$ and $\epsilon \neq 0$ is the

same. We consider then the model (1.1) with J_r replacing J , and we will choose $J_r = J + O(j^{-1})$ so that the critical temperature is in correspondence $t_c = \frac{1}{2} - 1$. We define

$$\langle O(x)O(y) \rangle = \frac{1}{Z} \sum_{\substack{x^{(1)}, x^{(2)} = \pm 1 \\ x^2 = M}}^X O(x)O(y) e^{H(x^{(1)}, x^{(2)})} \quad (1.10)$$

where $Z = \sum_{\substack{x^{(1)}, x^{(2)} = \pm 1 \\ x^2 = M}}^P e^{H(x^{(1)}, x^{(2)})}$ is the partition function,

$f(J) = \lim_{j \rightarrow 1} \frac{1}{j} \log Z$ is the free energy, $\frac{\partial^2}{\partial J^2} f(J)$ is the specific heat, and the correlation function of the observable $O(x)$ is

$$\langle O(x)O(y) \rangle_{;T} = \langle O(x)O(y) \rangle - \langle O(x) \rangle \langle O(y) \rangle : \quad (1.11)$$

In the Ising model close to the critical temperature $\frac{\partial^2}{\partial J^2} f^0(J) \sim C \log j - t_c j^{-1}$ and for large distances $j \langle O(x)O(y) \rangle_{;T} \sim j^{-\frac{A}{2}} \frac{C e^{-A j t_c j^{\frac{1}{2}} x} y^j}{j^{\frac{1}{2}} y^j}$, with A, C suitable constants.

We will prove the following theorem.

Theorem 1. Consider the hamiltonian (1.1), with J_r replacing J , and assume that $ja + bj > 0$ and (1.9). There exist $\epsilon > 0; a_3 > 0; a > 0$ constants and a function $\phi(j) = \tanh J_r - \tanh J$ such that, for $j \geq \epsilon$ and $a - j t_c j^{-\frac{1}{2}} e^{-\frac{1}{a_3 j^2}}$, the energy-energy correlation verifies

$$\lim_{j \rightarrow 1} \langle \phi(x) \phi(x^0) \phi(y) \phi(y^0) \rangle_{;T} = \phi^{(1)a}(x; y) + \phi^{(1)b}(x; y) + \phi^{(1)c}(x; y) \quad (1.12)$$

where $x; x^0$ and $y; y^0$ are nearest neighbor pairs respectively and, for any integer positive N

$$\begin{aligned} j \phi^{(1)a}(x; y) &\sim \frac{1}{j^{\frac{1}{2}} y^{\frac{1}{2} + 1}} \frac{C_N}{1 + (\frac{j}{y})^N} \\ j \phi^{(1)b}(x; y) &\sim \frac{1}{j^{\frac{1}{2}} y^{\frac{1}{2} + 3}} \frac{C_N}{1 + (\frac{j}{y})^N} \\ j \phi^{(1)c}(x; y) &\sim \frac{1}{j^{\frac{1}{2}} y^{\frac{1}{2} + 1}} \frac{C_N}{1 + (\frac{j}{y})^N} \end{aligned} \quad (1.13)$$

where $C_N; \epsilon$ are positive constants, $\epsilon = j t_c j^{\frac{1}{2} + 2}$, and $\phi(j) = O(j^{-1})$,

$$\phi_1(j) = a_1 + O(j^{-2}) \quad \phi_2(j) = a_2 + O(j^{-2}) \quad \phi_3(j) = a_1 + O(j^{-2}) \quad (1.14)$$

with $a_1 > 0; a_2 > 0$ constants. If $1 - \frac{j}{y} \sim 1$

$$\phi^{(1)a}(x; y) = \frac{1 + \phi^{(1)a}(x; y)}{Z_0^2 (x_0^2 + x^2)^{1 + \frac{1}{2}}} \quad \phi^{(1)b}(x; y) = \frac{1 + \phi^{(1)b}(x; y)}{Z_0^2 (x_0^2 + x^2)^{1 + \frac{3}{2}}} \quad (1.15)$$

and $j^{-1/c}(x;y)j \sim \frac{C}{|x-y|^{2+c}}$, where $Z_0 > 0$ is a constant and $f(x;y)$ are smooth bounded $O(1)$ functions. Finally the specific heat verifies

$$C_1 j^{-\frac{1}{1-\beta}} [1 - j^{-2-1}] + \frac{1}{3} [1 - j^{-2-3}] j^{-\frac{\partial^2}{\partial J^2} f(J)} \leq C_2 j^{-\frac{1}{1-\beta}} [1 - j^{-2-1}] + \frac{1}{3} [1 - j^{-2-3}] j^{-\frac{\partial^2}{\partial J^2} f(J)} \quad (1.16)$$

where C_1, C_2 are positive constants.

1.6 Remarks. The above theorem describes the behaviour of the energy-energy correlation and the specific heat near the critical temperature of Ising models in two dimensions weakly coupled by a four spin interaction. One can distinguish two different regimes in the asymptotic behaviour of the energy-energy correlation function, discriminated by an intrinsic correlation length of order $j - t_c j^{1-\beta}$ with $\beta = O(1)$. If $1 < |x-y| < \infty$, the bounds for the correlation function is power-like (with a β dependent exponent) while if $|x-y| < \infty$, there is a faster than any power decay with rate of order j^{-1} . In the first region we can obtain the exact large distance asymptotic behaviour of the energy-energy correlation function, see (1.15), while in the second region only an upper bound is obtained. The logarithmic behaviour of the specific heat in the Ising model getting closer and closer to the critical temperature is changed by the four spin interaction in a power law, at least up to temperatures very close to t_c for small four spin coupling. A bound like $C_1 \log(j - t_c j^{1-\beta}) - j^{-\frac{\partial^2}{\partial J^2} f(J)} \leq C_2 \log(j - t_c j^{1-\beta})$, for some positive β -independent constants C_1 and C_2 , which would be true if the model would be in the same universality class as the Ising model, cannot be true by (1.16) at least for $j - t_c j \sim e^{-\frac{1}{a_3-2}}$.

In the particular case $a = c = 0$ our model reduces to the Eight-vertex model, which is exactly soluble [Ba]; our results agree with the informations obtained by the exact solution, in which non universal critical behaviour is found and it is believed that the specific heat diverges with a power law [W]. In the case $a = b = 0$ our model reduces to a model of two non interacting Ising models with nearest neighbor and four spin interaction within each copy, which was studied in [PS] (which is indeed the first paper in which fermionic RG methods were applied to classical Ising-like models, and it is a major source of inspiration for the present work). In this case $C_1 \log(j - t_c j^{1-\beta}) - j^{-\frac{\partial^2}{\partial J^2} f(J)} \leq C_2 \log(j - t_c j^{1-\beta})$ (up to $t = t_c$) so that universality indeed holds.

1.7 Sketch of the proof. The proof of the theorem is based on Grassmann variables combined with renormalization group techniques. In x2 we briefly recall the wellknown representation of the bidimensional Ising model in terms of Grassmann variables, mainly due to [K], [H], [MW], [S] and recently rederived in a more cohesive way in [PS]. The partition function and the correlation can be written in terms of Grassmann integrals with a quadratic action which can be explicitly computed in terms of Pfaffians. The exact solvability of the Ising model is hence related to

the fact that it can be expressed in terms of free fermions, as it was first noted in [SML]. In the Grassmann representation there are four independent Grassmann variables associated to each point of the lattice, two with a large $O(1)$ mass and the other two with a continuously vanishing mass at the critical temperature. In x2 we show that two Ising models weakly coupled by four spin interactions, with hamiltonian (1.1), can be also written in terms of Grassmann integrals, with the difference that there are now eight independent Grassmann variables (four for each Ising model) and that the formal action contains now terms which are quartic in the fields, corresponding to a short range interaction among fermions. By performing a suitable linear transformation in the Grassmann variables one obtains that in the new variables the quadratic part of the action strongly resembles the action of two massive Dirac fermions in $d = 1 + 1$ dimensions (it would coincide with that if the continuum limit would be taken); again one Dirac fermion has a large $O(1)$ mass and the other is vanishing at t_c , and we call them heavy and light fermions. We are essentially exploiting in this transformation the relation $P^2 A = \det A$.

In x3 we integrate out the heavy fields in the Grassmann integral for the partition function, so obtaining a Grassmann integral whose formal action contains monomials of every degree in the light Dirac fermions. In order to do this we use the representation in [Le] of fermionic truncated expectations and Gram-Hadamard inequality.

In x4 we apply renormalization group methods to integrate the light fermions, which in a sense are the critical modes. We will use a suitable modification of the multiscale expansion used in [BM] (see also [BG] or [GM] for a general introduction to the formalism) to study the correlation functions of the Heisenberg-Ising XYZ chain; the close relationship between Ising models with quartic coupling and the XYZ times has been pointed out many times in the literature, see for instance [Ba]. A power counting analysis says that the terms bilinear in the Grassmann variables are relevant in a RG sense, while the quartic terms, or the bilinear with an extra derivative are marginal. One can understand here why universality is still present if one considers decoupled Ising models with nearest neighbor and four spin interaction within each copy, and why on the contrary it is lost if the Ising models are coupled. In the first case (which is the one treated in [SP]) one can easily check that the local part of the quartic terms is vanishing, so the effective interaction is indeed irrelevant in the RG sense. On the other hand in the second case the quartic interactions is truly marginal and this produces a line of fixed points for the RG transformation (instead of the gaussian fixed point as in the previous case) continuously depending on the coupling λ . We decompose the Grassmann integration $P(d)$ as a product of independent Grassmann integrations $P(d^{(h)})$, with covariance with non vanishing support only for momenta with modulus between $h-1$ and $h+1$, with $\lambda > 1$ and $h = 0; 1; 2; 3; \dots$. We integrate each $P(d^{(h)})$ iteratively starting from $P(d^{(1)})$ obtaining a sequence

of effective potentials $V^{(h)}$ describing the theory at momentum scale h ; at each step new contributions to the mass and the wave function renormalization are obtained which are included in the fermionic integration; hence $P(d^{(h)})$ has a covariance with mass m_h and wave function renormalization Z_h with a non trivial dependence on h , i.e. $m_h \propto t_c j^{-2h}$ and $Z_h \propto j^{-4h}$, with $j_4 = O(j^2)$. The iteration stops as soon as the mass m_h becomes of order h ; we will call h the last scale to be integrated (of course $h \rightarrow 1$ at the critical temperature $t = t_c$). The iterative procedure allows to write the effective potential $V^{(h)}$ as sum of monomials in the Grassmann variables, with coefficients which are (convergent) perturbative expansions in terms of a few running coupling constant, h , the effective coupling of the interaction between fermions, h , which takes into account the renormalization of the value of the critical temperature, h related to the renormalization of the fermionic velocity and other couplings $g_{ij}; h; \tilde{g}_{ij}; h$ which are the coefficients of quadratic terms in the Grassmann fields with a derivated field. Despite there are many similarities between our Grassmann integrals and the one describing relativistic fermions, there is a crucial difference; the interaction term in the action is not invariant under gauge transformation, hence terms which were absent in the free action can be generated in the RG iterations. One can check that indeed peculiar symmetries of the model (1.2) ensure that there is only one relevant term quadratic in the fermions (if there were more one gets in troubles, as there is only one free parameter in the hamiltonian). On the other hand we cannot exclude the generation of marginal quadratic terms which were absent in the free action; they are the terms $g_{ij}; h; \tilde{g}_{ij}; h$. At the end the result of this iterative integration is an expansion for the partition function in terms of the running coupling constants which is proved in x4 to be convergent provided that the running coupling constants are small for any h .

In x5 we prove indeed that it is possible to choose the counterterm as a function of j so that the running coupling constants are indeed small; the condition on the temperature $j \propto t_c j \propto e^{\frac{1}{a_3 - 2}}$ is used to control the flow of the running coupling constants, and some cancellations at the lower order in our expansion are also used to be as close as possible to the critical temperature.

Finally in x6 we define an expansion for the correlation functions and the specific heat; it is similar to the one for the partition function, with the main difference that one has to introduce new fields associated to the external fields. There are additional marginal terms in RG expansion to which other renormalization constants, with a non trivial behaviour in h , are associated, i.e. $Z_h^{(1)} \propto j^{-1h}$ and $Z_h^{(2)} \propto j^{-3h}$. By such expansion the statements in the theorem are derived.

Among interesting open problems there is the reaching of the critical temperature; surely one has to exploit suitable cancellations at every order of the expansion for $h; h$, as in the theory of $d = 1$ interacting Fermi systems, see for instance [BGPS] and [BM], but at the moment the main difficulty is in the flow of the couplings $g_{ij}; h$ and $\tilde{g}_{ij}; h$. Another interesting problem would be the analysis of the

spin-spin correlation function whose expression in terms of Grassmann variables is unfortunately quite complicated and not easy to manage. A similar problem appears in considering two Ising models coupled by a weak interaction bilinear in the spins; in such a case the interaction in terms of Grassmann variables has a quite complex expression and it seems difficult to study. On the other hand, for large coupling, such bilinear interaction should be irrelevant and universality should hold. Finally it should be interesting to study the case of two coupled Ising models at different temperatures or the case of four coupled Ising models; in this last case interacting spinning $d = 1$ fermions appear in the fermionic description, which are known to have a behaviour quite different from the spinless one (like in the $d = 1$ Hubbard model).

2. Fermionic representation

2.1 Grassmann integrals. If $\beta = 0$ the hamiltonian (1.1) is given by the sum of two independent Ising model hamiltonians, and the partition function is given by $Z_I^{(1)} Z_I^{(2)}$ where

$$Z_I^{(\beta)} = \sum_{\{s_i\}} e^{-\beta H_I(\{s_i\})} \quad (2.1)$$

It is well known that such partition function can be written in terms of Grassmann integrals, so we recall first their definition. Grassmann variables $\psi_i, \bar{\psi}_i, i = 1, 2, \dots, 2n$, n even, are anticommuting variables satisfying

$$\psi_i^2 = 0, \quad \bar{\psi}_i^2 = 0 \quad (2.2)$$

The Grassmann integration $\int d\psi_i d\bar{\psi}_i$ is a linear operation defined as

$$\int d\psi_i d\bar{\psi}_i = 0, \quad \int d\psi_i d\bar{\psi}_i \psi_i \bar{\psi}_i = 1 \quad (2.3)$$

It holds that

$$\int d\psi_i d\bar{\psi}_i e^{\frac{1}{2} \sum_{i,j} A_{ij} \psi_i \bar{\psi}_j} = Pf A \quad (2.4)$$

where A is an even antisymmetric $2n \times 2n$ matrix and $Pf A$ denotes the Pfaffian. It holds

$$\int d\psi_1 \dots d\psi_{2n} \exp \frac{1}{2} \sum_{i,j} A_{ij} \psi_i \bar{\psi}_j = Pf A \quad (2.5)$$

$$\int d\psi_{2n} \dots d\psi_1 (1 + \sum_{i,j} A_{ij} \psi_i \bar{\psi}_j) = \frac{1}{2^n n!} \sum_p (-1)^p A_{p_1 p_2} A_{p_3 p_4} \dots A_{p_{2n-1} p_{2n}} Pf A$$

where the sum is over all the permutations. We can consider another set of Grassmann variables $\psi_i, i = 1; \dots; 2n$, and

$$\int \prod_{i=1}^{2n} d\psi_i \exp \left[\sum_{i,j=1}^{2n} B_{ij} \psi_i \psi_j \right] = \text{Det} B \quad (2.6)$$

The well known relation $(\text{Pf} A)^2 = \text{det} A$ can be quite easily deduced by the above Grassmann integrals; it can be written as

$$\int \prod_{i=1}^{2n} d\psi_i e^{-\sum_{i,j=1}^{2n} A_{ij} \psi_i \psi_j} = \int \prod_{i=1}^{2n} d\psi_i^{(1)} e^{-\frac{1}{2} \sum_{i,j=1}^{2n} A_{ij}^{(1)} \psi_i^{(1)} \psi_j^{(1)}} \int \prod_{i=1}^{2n} d\psi_i^{(2)} e^{-\frac{1}{2} \sum_{i,j=1}^{2n} A_{ij}^{(2)} \psi_i^{(2)} \psi_j^{(2)}} \quad (2.7)$$

which can be proved by the change of variables

$$\psi_i = \frac{1}{\sqrt{2}} (\psi_i^{(1)} + i \psi_i^{(2)}) = \frac{1}{\sqrt{2}} (\psi_i^{(1)} - i \psi_i^{(2)}) \quad (2.8)$$

in $\prod_{i=1}^{2n} d\psi_i e^{-\sum_{i,j=1}^{2n} A_{ij} \psi_i \psi_j}$. Then $d\psi_i d\psi_j = i d\psi_i^{(1)} d\psi_j^{(2)}$ and

$$A_{ij} = \frac{1}{2} A_{ij}^{(1)} + \frac{1}{2} A_{ij}^{(2)} \quad (2.9)$$

as

$$A_{ij}^{(1)} A_{ij}^{(2)} = A_{ij}^{(1)} A_{ij}^{(2)} = 0 \quad (2.10)$$

2.2 The fermionic representation of the Ising model. It is well known that the partition function can be written as a Grassmann integral. It is proved in [K], [H], [MW], [S] that

$$Z_I = (\cosh J_r)^B 2^S \int \prod_{x \in \Lambda} dH_x dV_x \exp \left[\sum_{(x,y) \in E} J_{xy} H_x H_y + \sum_{x \in \Lambda} h_x H_x + \sum_{x \in \Lambda} v_x V_x \right] \quad (2.11)$$

where B is the total number of bonds and S is the total number of sites,

$$\begin{aligned} \sum_{x \in \Lambda} \tanh J_r [H_x H_{x+1} + V_x V_{x+1}] + \\ \sum_{x \in \Lambda} [h_x H_x + v_x V_x] \end{aligned} \quad (2.12)$$

and H_x, V_x are Grassmann variables such that

$$\begin{aligned} H_{x+M} &= H_x, \quad V_{x+M} = V_x \\ H_{x+M} &= H_x, \quad V_{x+M} = V_x \end{aligned} \quad (2.13)$$

and identical relations hold for the variables $V^{(\cdot)}; V^{(\cdot)}$. The Grassmann integration

$\int_{\mathbb{R}^Q} \prod_x dH_x^{(\cdot)} dH_x^{(\cdot)}$ is defined as the linear functional on the Grassmannian algebra, such that, given a monomial $Q(H^{(\cdot)}; H^{(\cdot)})$ in the variables $H_x^{(\cdot)}; H_x^{(\cdot)}$, $x \in \mathbb{Z}^2_M$, its value is 0, except in the case $Q(H^{(\cdot)}; H^{(\cdot)}) = \prod_x H_x^{(\cdot)} H_x^{(\cdot)}$, up to a permutation of the variables. In this case the value of the functional is determined, by using the anticommuting properties of the variables, by the condition

$$\int_{\mathbb{R}^Q} \prod_{x \in \mathbb{Z}^2_M} dH_x^{(\cdot)} dH_x^{(\cdot)} \prod_{x \in \mathbb{Z}^2_M} H_x^{(\cdot)} H_x^{(\cdot)} = 1 : \quad (2.14)$$

In a similar way is defined the Grassmann integration for $V^{(\cdot)}; V^{(\cdot)}$, just exchanging $H; H$ with $V; V$. In order to prove (2.11) the starting point is the high temperature expansion

$$Z_I^{(\cdot)} = (\cosh J_r)^B 2^S \sum_X (\tanh J_r)^L \quad (2.15)$$

where the sum is over all the closed polygons which may have points but not sides in common, see Fig.4. One then replaces the Ising lattice with another lattice, called second lattice, in which each site is replaced by four surrounding sites or terminals, see Fig.5. If we associate V_x with N_x , V_x with S_x , H_x with E_x and H_x with W_x , see Fig.5, it is easy to verify, by the rules of Grassmann integrations, that $e^{S_{\cdot}; \cdot}$ can be represented as a sum over polygon configurations over the second lattice. Such polygon configurations are defined such that each terminal must be coupled to another terminal, E_x may be coupled to $W_{x+1; x_0}$; $N_{x_0; x}$ may be coupled to $S_{x; x_0+1}$; E_x may be coupled to N_x ; W_x ; S_x ; N_x may be coupled to W_x or S_x ; W_x may be coupled to S_x , and no other types of couplings are allowed. An example of polygon configuration in the second lattice is in Fig.5. By a local identification of Ising configurations and configurations over the second lattice (see Fig.6), there is a correspondence between polygon configurations on the Ising lattice and polygon configurations on the second lattice. The correspondence is not one to one, as the configuration with zero lines in the Ising model corresponds to three configurations in the second lattice. However the signs add so that the sum over polygon configurations in the Ising lattice in (2.15) is exactly equal to the sum over polygon configurations in (2.11); the proof is based on a subtle combinatorial analysis first done in [K]. In order to ensure periodic boundary conditions, one has to sum over four terms as in (2.11), as explained in [MW]. We write $e^{S_{\cdot}; \cdot} = e^{S_{\cdot}; \cdot; 0} e^{S_{\cdot}; \cdot}$ where $S_{\cdot}; \cdot; 0$ is given by (2.12) with J replacing J_r and

$$S_{\cdot}; \cdot = \sum_{x \in \mathbb{Z}^2_M} [H_{x; x_0} H_{x+1; x_0} + V_{x; x_0} V_{x; x_0+1}] \quad (2.16)$$

If J is not constant but it depends on the bounds one obtains a similar formula in which $S_{x,x_0}^{() ; 0}$ is given by

$$S_{x,x_0}^{() ; 0} = \sum_x \left[\tanh J_{1;x;x_0;x+1;x_0} H_{x;x_0}^{()} H_{x+1;x_0}^{()} + \tanh J_{2;x;x_0;x;x_0+1} V_{x;x_0}^{()} V_{x;x_0+1}^{()} + \right. \\ \left. \sum_x \left[H_{x;x_0}^{()} H_{x;x_0}^{()} + V_{x;x_0}^{()} V_{x;x_0}^{()} + V_{x;x_0}^{()} H_{x;x_0}^{()} + V_{x;x_0}^{()} H_{x;x_0}^{()} + H_{x;x_0}^{()} V_{x;x_0}^{()} + V_{x;x_0}^{()} H_{x;x_0}^{()} \right] \right] \quad (2.17)$$

and the factor $(\cosh J)^B$ is replaced by $\prod_b \cosh J_b$, where the product is over all the possible nearest neighbor bounds. We will call $Z_I^{()}(J_{x;x_0})$ the Ising model partition function with non constant J .

2.3 The fermionic representation of coupled Ising models. The partition function of (1.1) is

$$Z_{2I} = \sum_{x=1}^X \sum_{x_0=1}^X e^{H_I^{(1)}} e^{H_I^{(2)}} e^{V^{(1);(2)}} \quad (2.18)$$

Let us consider first the essentially equivalent expression

$$\hat{Z}_{2I} = \sum_{x=1}^X \sum_{x_0=1}^X e^{H_I^{(1)}} e^{H_I^{(2)}} \\ \prod_x \left[1 + a_{x;x_0}^{(1)} a_{x+1;x_0}^{(1)} a_{x;x_0}^{(2)} a_{x+1;x_0}^{(2)} \right] \prod_x \left[1 + a_{x;x_0}^{(1)} a_{x;x_0+1}^{(1)} a_{x;x_0}^{(2)} a_{x;x_0+1}^{(2)} \right] \\ \prod_x \left[1 + b_{x;x_0}^{(1)} a_{x+1;x_0}^{(1)} a_{x;x_0}^{(2)} a_{x;x_0+1}^{(2)} \right] \prod_x \left[1 + b_{x;x_0}^{(1)} a_{x;x_0+1}^{(1)} a_{x+1;x_0+1}^{(2)} a_{x;x_0+1}^{(2)} \right] \quad (2.19) \\ \prod_x \left[1 + c_{x;x_0}^{()} a_{x+1;x_0}^{()} a_{x;x_0}^{()} a_{x;x_0+1}^{()} \right] \prod_x \left[1 + c_{x;x_0}^{()} a_{x;x_0}^{()} a_{x+1;x_0}^{()} a_{x+1;x_0+1}^{()} \right]$$

Noting that

$$a_{x;x_0}^{()} a_{x+1;x_0}^{()} e^{H_I^{()}} = \frac{\partial}{\partial J_{1;x;x_0;x+1;x_0}} Z_I^{()}(J_{x;x_0}) \Big|_{J_{x;x_0} g = f J^{()} g} \quad (2.20)$$

and from (2.17) this derivative gives an extra factor $\tanh J^{()} + \text{sech}^2 J^{()} H_{x;x_0}^{()} H_{x+1;x_0}^{()}$ in (2.11). In the same way

$$a_{x;x_0}^{()} a_{x;x_0+1}^{()} e^{H_I^{()}} = \frac{\partial}{\partial J_{2;x;x_0;x;x_0+1}} Z_I^{()}(J_{x;x_0}) \Big|_{J_{x;x_0} g = f J^{()} g} \quad (2.21)$$

and this derivative gives a factor $\tanh J^{(\cdot)} + \text{sech}^2 J^{(\cdot)} V_{x;x_0}^{(\cdot)} V_{x;x_0+1}^{(\cdot)}$. We can write than, if $_{+;+} = 1$ and $_{+; } = \text{ ; } = 2$

$$\hat{Z}_{2I}^{n^{(1)}; n^{(1)}; n^{(2)}; n^{(2)}} = \prod_{n^{(1)}; n^{(1)}}^X (1)^{n^{(1)}; n^{(1)}} \prod_{n^{(2)}; n^{(2)}}^X (1)^{n^{(2)}; n^{(2)}} \hat{Z}_{2I}^{n^{(1)}; n^{(1)}; n^{(2)}; n^{(2)}} \quad (2.22)$$

where

$$\begin{aligned} \hat{Z}_{2I}^{n^{(1)}; n^{(1)}; n^{(2)}; n^{(2)}} &= (\cosh J)^{2B} 2^{2S} \frac{1}{4} \\ &\int_{=1}^Z \int_{=1}^{Y^2} \int_{=1}^Y dH_x^{(\cdot)} dH_x^{(\cdot)} dV_x^{(\cdot)} dV_x^{(\cdot)} e^{S_{n^{(1)}; n^{(1)}}^{(1)}} e^{S_{n^{(2)}; n^{(2)}}^{(2)}} \\ &\int_{=1}^Y [1 + a(\tanh J + \text{sech}^2 J H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)}) (\tanh J + \text{sech}^2 J H_{x;x_0}^{(2)} H_{x+1;x_0}^{(2)})] \\ &\int_{=1}^x [1 + a(\tanh J + \text{sech}^2 J V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}) (\tanh J + \text{sech}^2 J V_{x;x_0}^{(2)} V_{x;x_0+1}^{(2)})] \\ &\int_{=1}^x [1 + b(\tanh J + \text{sech}^2 J H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)}) (\tanh J + \text{sech}^2 J V_{x;x_0}^{(2)} V_{x;x_0+1}^{(2)})] \\ &\int_{=1}^Y [1 + b(\tanh J + \text{sech}^2 J V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}) (\tanh J + \text{sech}^2 J H_{x-1;x_0+1}^{(2)} H_{x;x_0+1}^{(2)})] \\ &\int_{=1}^x \int_{=1}^Y [1 + c(\tanh J + \text{sech}^2 J H_{x;x_0}^{(\cdot)} H_{x+1;x_0}^{(\cdot)}) (\tanh J + \text{sech}^2 J V_{x;x_0}^{(\cdot)} V_{x;x_0+1}^{(\cdot)})] \\ &\int_{=1}^Y \int_{=1}^x [1 + c(\tanh J + \text{sech}^2 J H_{x;x_0}^{(\cdot)} H_{x+1;x_0}^{(\cdot)}) (\tanh J + \text{sech}^2 J V_{x+1;x_0-1}^{(\cdot)} V_{x+1;x_0}^{(\cdot)})] \end{aligned}$$

The above expression can be rewritten as

$$\hat{Z}_{2I}^{n^{(1)}; n^{(1)}; n^{(2)}; n^{(2)}} = (\cosh J)^{2B} 2^{2S} \frac{1}{4} \int_{=1}^Z \int_{=1}^{Y^2} \int_{=1}^Y dH_x^{(\cdot)} dH_x^{(\cdot)} dV_x^{(\cdot)} dV_x^{(\cdot)} e^{S_{n^{(1)}; n^{(1)}}^{(1)}} e^{S_{n^{(2)}; n^{(2)}}^{(2)}} \quad (2.24)$$

with

$$V = V_a + V_b + V_c \quad (2.25)$$

and, if $f_i = \log(1 + [i] \tanh^2 J)$ and $[i] = a; b; c$

$$\begin{aligned} V_a &= \prod_x [f_a + \tilde{a} [H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} + H_{x;x_0}^{(2)} H_{x+1;x_0}^{(2)}] + a H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} H_x^{(2)} H_{x+1;x_0}^{(2)}] \\ &+ \prod_x [f_a + \tilde{a} [V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)} + V_{x;x_0}^{(2)} V_{x;x_0+1}^{(2)}] + a V_x^{(1)} V_{x;x_0+1}^{(1)} V_{x;x_0}^{(2)} V_{x;x_0+1}^{(2)}] \quad (2.26) \\ V_b &= \prod_x [f_b + \tilde{b} [H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} + V_{x;x_0}^{(2)} V_{x;x_0+1}^{(2)}] + b H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} V_x^{(2)} V_{x;x_0+1}^{(2)}] \end{aligned}$$

$$\begin{aligned}
& + \sum_x [\tilde{f}_b + \tilde{b} [V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)} + H_{x-1;x_0+1}^{(2)} H_{x;x_0+1}^{(2)}] + \tilde{b} V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)} H_{x-1;x_0+1}^{(2)} H_{x;x_0+1}^{(2)}] \\
V_c = & \sum_x [\tilde{f}_c + \tilde{c} [H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} + V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}] + \tilde{c} H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}] \\
& + \sum_x [\tilde{f}_c + \tilde{c} [V_{x+1;x_0-1}^{(1)} V_{x+1;x_0}^{(1)} + H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)}] + \tilde{c} V_{x+1;x_0-1}^{(1)} V_{x+1;x_0}^{(1)} H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)}]
\end{aligned}$$

It is easy in fact to verify that

$$e^{\tilde{f}_i + \tilde{i} [H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} + V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}] + \tilde{i} H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}} = f(1 + [\tilde{i}] \tanh^2 J) \quad (2.27)$$

$$[1 + \tilde{i} [H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} + V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}] + (\tilde{i} + (\tilde{i})^2) H_{x;x_0}^{(1)} H_{x+1;x_0}^{(1)} V_{x;x_0}^{(1)} V_{x;x_0+1}^{(1)}] g$$

hence the equality between (2.23) and (2.24) holds with the identification

$$\begin{aligned}
\tilde{i} (1 + [\tilde{i}] \tanh^2 J) &= [\tilde{i}] \operatorname{sech}^2 J \tanh J \\
(1 + [\tilde{i}] \tanh^2 J) (\tilde{i} + (\tilde{i})^2) &= [\tilde{i}] \operatorname{sech}^4 J \quad (2.28)
\end{aligned}$$

An expression identical to (2.22), (2.23) holds for Z_{2I} , the only difference being that the relation with respect \tilde{i} ; \tilde{i} is slightly more complicated than (2.28), but for small J again

$$\tilde{i} = [\tilde{i}] (\tanh J \operatorname{sech}^2 J + O(J^3)) \quad \tilde{i} = [\tilde{i}] (\operatorname{sech}^4 J + O(J^3)) \quad (2.29)$$

2.4 Massive and massless fermions. We define

$$P_{\mathbf{n}^{(1)}; \mathbf{n}^{(0)}}^{(1)}(d\mathbf{H}; d\mathbf{V}^{(1)}) = \prod_x dH_x^{(1)} dH_x^{(1)} dV_x^{(1)} dV_x^{(1)} e^{S_{\mathbf{n}^{(1)}; \mathbf{n}^{(0)}}^{(1)}} \quad (2.30)$$

It is convenient to perform the following change of variables [D], [SP]

$$\begin{aligned}
H_x^{(1)} + iH_x^{(1)} &= e^{i\bar{4}} x^{(1)} e^{i\bar{4}} x^{(1)} \quad H_x^{(1)} - iH_x^{(1)} = e^{-i\bar{4}} x^{(1)} e^{-i\bar{4}} x^{(1)} \\
V_x^{(1)} + iV_x^{(1)} &= x^{(1)} + x^{(1)} \quad V_x^{(1)} - iV_x^{(1)} = x^{(1)} + x^{(1)} \quad (2.31)
\end{aligned}$$

It holds that

$$P_{\mathbf{n}^{(1)}; \mathbf{n}^{(0)}}^{(1)}(d\mathbf{H}; d\mathbf{V}^{(1)}) = P_{\mathbf{n}^{(1)}; \mathbf{n}^{(0)}}^{(1)}(d) P_{\mathbf{n}^{(1)}; \mathbf{n}^{(0)}}^{(1)}(d) e^{Q^{(1)}; \mathbf{n}^{(0)}} \quad (2.32)$$

where, if $t = \tanh J$

$$P_{\mathbf{n}^{(1)}; \mathbf{n}^{(0)}}^{(1)}(d^{(1)}) = \left[\prod_{x \in M} d_x^{(1)} d_x^{(1)} \right] \exp \frac{t}{4} \sum_{x \in M} x^{(1)} (\partial_1 - i\partial_0) x^{(1)} + x^{(1)} (\partial_1 + i\partial_0) x^{(1)}$$

$$\begin{aligned}
& + \frac{t}{4} \sum_{x \in \mathbb{Z}^M} [i \psi_x^{(1)} (\phi_{1,x}^{(1)} + \phi_{0,x}^{(1)}) + i \psi_x^{(2)} (\phi_{1,x}^{(2)} + \phi_{0,x}^{(2)})] \\
& + \sum_{x \in \mathbb{Z}^M} i \left(\frac{p}{2} - 1 + t \right) \psi_x^{(1)} \psi_x^{(2)} g
\end{aligned} \quad (2.33)$$

where

$$\phi_{1,x}^{(1)} = \psi_{x+1, x_0}^{(1)} \psi_x^{(1)} \quad \phi_{0,x}^{(1)} = \psi_{x, x_0+1}^{(1)} \psi_x^{(1)} \quad (2.34)$$

Moreover

$$\begin{aligned}
P_{n,n_0}^{(1)}(d^{(1)}) &= \left[\sum_{x \in \mathbb{Z}^M} d_x^{(1)} d_x^{(1)} \right] \exp \left[\frac{t}{4} \sum_{x \in \mathbb{Z}^M} \psi_x^{(1)} (\phi_{1,x}^{(1)} - i \phi_{0,x}^{(1)}) \psi_x^{(1)} + \psi_x^{(1)} (\phi_{1,x}^{(1)} + i \phi_{0,x}^{(1)}) \psi_x^{(1)} + \right. \\
& \left. \frac{t}{4} \sum_{x \in \mathbb{Z}^M} [i \psi_x^{(1)} (\phi_{1,x}^{(1)} + \phi_{0,x}^{(1)}) + i \psi_x^{(2)} (\phi_{1,x}^{(2)} + \phi_{0,x}^{(2)})] \sum_{x \in \mathbb{Z}^M} i \left(\frac{p}{2} - 1 + t \right) \psi_x^{(1)} \psi_x^{(2)} g \right]
\end{aligned} \quad (2.35)$$

and finally

$$\begin{aligned}
Q^{(1)}(\psi^{(1)}; \psi^{(2)}) &= \sum_x \frac{t}{4} f \left[\psi_x^{(1)} (\phi_{1,x}^{(1)} + i \phi_{0,x}^{(1)}) \psi_x^{(1)} (\phi_{1,x}^{(1)} - i \phi_{0,x}^{(1)}) \right. \\
& \left. \psi_x^{(1)} (\phi_{1,x}^{(1)} + i \phi_{0,x}^{(1)}) \psi_x^{(1)} (\phi_{1,x}^{(1)} - i \phi_{0,x}^{(1)}) + i \psi_x^{(1)} (\phi_{1,x}^{(1)} - \phi_{0,x}^{(1)}) \right. \\
& \left. + \psi_x^{(1)} (\phi_{1,x}^{(1)} - \phi_{0,x}^{(1)}) + \psi_x^{(1)} (\phi_{1,x}^{(1)} - \phi_{0,x}^{(1)}) + \psi_x^{(1)} (\phi_{1,x}^{(1)} + \phi_{0,x}^{(1)}) g \right]
\end{aligned} \quad (2.36)$$

We find convenient to rewrite the Grassmann variables in momentum space. We call D_{n,n_0} the set of k such that

$$k = \frac{2n_1}{M} + \frac{(n-1)}{2M} \quad k_0 = \frac{2n_0}{M} + \frac{(n^0-1)}{2M} \quad (2.37)$$

and $[M=2] \quad n_0 \in [(M-1)=2], \quad [M=2] \quad n_1 \in [(M-1)=2], \quad n_0, n_1 \in \mathbb{Z}$. We can write if $k = (k_0; k)$

$$\psi_x^{(1)} = \frac{1}{M^2} \sum_{k \in D_{n,n_0}} \psi_k^{(1)} e^{ikx} \quad \psi_x^{(2)} = \frac{1}{M^2} \sum_{k \in D_{n,n_0}} \psi_k^{(2)} e^{ikx} \quad (2.38)$$

Hence

$$\begin{aligned}
P_{n,n_0}^{(1)}(d) &= \left[\sum_{k \in D_{n,n_0}} d_k^{(1)} d_k^{(1)} \right] \exp \left[\frac{t}{4M^2} \sum_{k \in D_{n,n_0}} [\psi_k^{(1)} \psi_k^{(1)} (i \sin k + \sin k_0) \right. \\
& \left. + \psi_k^{(1)} \psi_k^{(1)} (i \sin k - \sin k_0) + i 2m(k) \psi_k^{(1)} \psi_k^{(2)}] \right]
\end{aligned} \quad (2.39)$$

where $m(k) = \frac{2}{t} \left(\frac{p}{2} - 1 - t \right) + (2 \cos k_0 + 2 \cos k - 4) = 2 \frac{(t - t_c)}{t} + O(k^2)$. In deriving the above formula we have used that

$$\sum_x \psi_x^{(0)} \psi_x^{(0)} = \sum_{k \in 2D_{n,n_0}} (e^{ik_0} - 1) \psi_k^{(0)} \psi_k^{(0)} = \sum_{k \in 2D_{n,n_0}} i \sin k_0 \psi_k^{(0)} \psi_k^{(0)} \quad (2.40)$$

as

$$\begin{aligned} \sum_{k \in 2D_{n,n_0}} (\cos k_0 - 1) \psi_k^{(0)} \psi_k^{(0)} &= \sum_{k \in 2D_{n,n_0}} (\cos k_0 - 1) \psi_k^{(0)} \psi_k^{(0)} \\ &= \sum_{k \in 2D_{n,n_0}} (\cos k_0 - 1) \psi_k^{(0)} \psi_k^{(0)} = 0 \end{aligned} \quad (2.41)$$

2.5 Majorana and Dirac fermions. We write $\psi_x^{(0)}$ in (2.38) as

$$\psi_x^{(0)} = e^{ip_{n,n_0} x} \frac{1}{M^{1/2}} \sum_{k \in 2D_{n,n_0}} \psi_{k^0 + p_{n,n_0}}^{(0)} e^{-ik^0 x} = e^{ip_{n,n_0} x} \psi_x^{(0)} \quad (2.42)$$

where $p_n = \left(-\frac{(n+1)}{2M}; -\frac{(n^0+1)}{2M} \right)$, and $\psi_{k^0}^{(0)} = \psi_{k^0 + p_{n,n_0}}^{(0)}$; moreover

$$P_{n,n_0}^{(0)}(d) = P^{(0)}(d^0) e^{\mathcal{Q}_{n,n_0}^{(0)}} \quad (2.43)$$

where

$$\begin{aligned} P^{(0)}(d^0) &= \left[\sum_{k^0 \in 2D_{n,n_0}} d_{k^0}^{(0)} d_{k^0}^{(0)} \right] \exp \left[\frac{t}{4M^{1/2}} \sum_{k^0 \in 2D_{n,n_0}} \left[d_{k^0}^{(0)} d_{k^0}^{(0)} (i \sin k^0 + \sin k_0^0) \right. \right. \\ &\quad \left. \left. + d_{k^0}^{(0)} d_{k^0}^{(0)} (i \sin k^0 - \sin k_0^0) + i 2m(k^0) d_{k^0}^{(0)} d_{k^0}^{(0)} \right] \right] \end{aligned} \quad (2.44)$$

and $\mathcal{Q}_{n,n_0}^{(0)}$ is defined by (2.43) and is formally vanishing in the limit $M \rightarrow 1$. Proceeding in the same way for $P(d)$ and Q we find

$$\begin{aligned} \sum_{n=1}^Z \sum_{n^0=1}^{Y^2} & \left[P_{n(n),n^0(n)}^{(0)}(d) P_{n(n),n^0(n)}^{(0)}(d) \right] e^{\mathcal{Q}_{n(n),n^0(n)}^{(0)}} e^{V(n;n)} = \\ & \sum_{n=1}^Z \sum_{n^0=1}^{Y^2} \left[P^{(0)}(d^0) P^{(0)}(d^0) \right] e^{\mathcal{Q}_{n(n),n^0(n)}^{(0)}} e^{\mathcal{Q}_{n(n),n^0(n)}^{(0)}} e^{V(n;n)} \end{aligned} \quad (2.45)$$

where

$$\mathcal{Q}_{n(n),n^0(n)}^{(0)} = \sum_{n=1}^{X^2} \left[\mathcal{Q}_{n(n),n^0(n)}^{(0)} + \mathcal{Q}_{n(n),n^0(n)}^{(0)} + \mathcal{Q}_{n(n),n^0(n)}^{(0)} \right] \quad (2.46);$$

of course $\mathcal{Q}^{(0)}_{\pm} = 0$. If

$$D_{\pm}^{(0)} = \sqrt{k^0} D_{\pm}; \quad k^0 > 0 \quad (2.47)$$

we can write $P^{(0)}(d^{(0)})$ in the following way

$$P^{(0)}(d^{(0)}) = \int_{k^0 D^+}^Y d_{k^0}^{(0)} d_{k^0}^{(0)} d_{k^0}^{(0)} d_{k^0}^{(0)} \exp\left[\frac{t}{4M^2} X^{(0)T} A(k^0) X^{(0)}\right] \quad (2.48)$$

where

$$X^{(0)T} = (d_{k^0}^{(0)}; d_{k^0}^{(0)}; d_{k^0}^{(0)}; d_{k^0}^{(0)}) \quad (2.49)$$

and $A(k^0)$ is

$$A(k^0) = \begin{pmatrix} \sin k^0_0 + i \sin k^0_0 & 0 & 0 & 1 \\ \sin k^0_0 & i \sin k^0_0 & 0 & \sin k^0_0 \\ 0 & 0 & \sin k^0_0 + i \sin k^0_0 & 0 \\ 0 & 0 & \sin k^0_0 & i \sin k^0_0 \end{pmatrix} \quad (2.50)$$

Hence if we perform the change of variables

$$d_{k^0}^{(1)} = \frac{1}{\sqrt{2}} (d_{k^0}^{(0)} + i d_{k^0}^{(2)}) \quad d_{k^0}^{(2)} = \frac{1}{\sqrt{2}} (d_{k^0}^{(0)} - i d_{k^0}^{(1)}) \quad (2.51)$$

$$d_{k^0}^{(1)} = \frac{1}{\sqrt{2}} (d_{k^0}^{(0)} + i d_{k^0}^{(2)}) \quad d_{k^0}^{(2)} = \frac{1}{\sqrt{2}} (d_{k^0}^{(0)} - i d_{k^0}^{(1)})$$

we find by (2.7)

$$P^{(1)}(d^{(0)}) P^{(2)}(d^{(0)}) = \int_{k^0 D^+}^Y d_{k^0}^{(1)} d_{k^0}^{(1)} d_{k^0}^{(2)} d_{k^0}^{(2)} \exp\left[\frac{t}{4M^2} X^{(1)T} A(k^0) X^{(2)}\right] \quad (2.52)$$

where

$$X^{(1)T} = (d_{k^0}^{(1)}; d_{k^0}^{(1)}; d_{k^0}^{(2)}; d_{k^0}^{(2)}) \quad (2.53)$$

$$X^{(2)T} = (d_{k^0}^{(1)}; d_{k^0}^{(1)}; d_{k^0}^{(2)}; d_{k^0}^{(2)})$$

We can make another change of variables $d_{k^0}^{(1)} \rightarrow d_{k^0}^{(1)}$ so that at the end, if

$$P^{(1)}(d^{(0)}) P^{(2)}(d^{(0)}) = P(d^{(0)}) \quad (2.54)$$

$$P(d^{(0)}) = \int_{k^0 D^+}^Y d_{k^0}^{(1)} d_{k^0}^{(1)} d_{k^0}^{(2)} d_{k^0}^{(2)} \exp\left[\frac{t}{2M^2} X^{(1)T} A(k^0) X^{(2)}\right]$$

where

$$X^{(1)T} = (d_{k^0}^{(1)}; d_{k^0}^{(1)}; d_{k^0}^{(2)}; d_{k^0}^{(2)})$$

$$\hat{\mathbf{k}}^+;T = \begin{pmatrix} + & + \\ k;1 & k;1 \end{pmatrix} \quad (2.55)$$

and in terms of the original variables

$$_{1;k^0} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ k^0 & k^0 \end{pmatrix} \quad +_{1;k^0} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ k^0 & k^0 \end{pmatrix} \quad (2.56)$$

$$_{1;k} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ k^0 & k^0 \end{pmatrix} \quad +_{1;k} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ k^0 & k^0 \end{pmatrix}$$

In the following we will call k^0 simply k . If we write

$$\mathbf{x} = \frac{1}{M^2} \sum_{k \in D^+} e^{ikx} \quad (2.57)$$

we can rewrite (2.56) in the coordinate space as

$$_{1;x} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ x & x \end{pmatrix} \quad +_{1;x} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ x & x \end{pmatrix} \quad (2.58)$$

$$_{1;x} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ x & x \end{pmatrix} \quad +_{1;x} = \frac{1}{2} \begin{pmatrix} {}^0(1) & {}^0(2) \\ x & x \end{pmatrix}$$

or in terms of the original variables

$$\begin{aligned} _{1;x} &= \frac{1}{2} (e^{ip_n(1);n^0(1)x} {}^{(1)}_x + ie^{ip_n(2);n^0(2)x} {}^{(2)}_x) \\ +_{1;x} &= \frac{1}{2} (e^{ip_n(1);n^0(1)x} {}^{(1)}_x - ie^{ip_n(2);n^0(2)x} {}^{(2)}_x) \\ _{1;x} &= \frac{1}{2} (e^{ip_n(1);n^0(1)x} {}^{(1)}_x + ie^{ip_n(2);n^0(2)x} {}^{(2)}_x) \\ +_{1;x} &= \frac{1}{2} (e^{ip_n(1);n^0(1)x} {}^{(1)}_x - ie^{ip_n(2);n^0(2)x} {}^{(2)}_x) \end{aligned} \quad (2.59)$$

If $k \in D^+$, then $k \in D$; with $D^+ \subset D$; $D^+ \cup D^+ = D$; so that

$$P(d) = \quad (2.60)$$

$$\sum_{k \in D^+} d_{k;1}^+ d_{k;1} \exp \left[\frac{t}{2M^2} \sum_{k \in D^+} \tilde{A}^T(k) {}^{(+)}_k \right]$$

$$\tilde{A}(k) = \frac{i \sin k + \sin k_0}{i \sin k - \sin k_0} \quad \text{im} \quad (k)$$

$$\tilde{\mathbf{k}}^T = \begin{pmatrix} + & + \\ k;1 & k;1 \end{pmatrix} \quad \tilde{\mathbf{k}}^+;T = \begin{pmatrix} + & + \\ k;1 & k;1 \end{pmatrix} \quad (2.61)$$

Remark. In the physical language, the change of variables (2.59) means that one is describing the system in terms of Dirac fermions instead in terms of Majorana fermions.

A similar computation can be done for $P(d)$; proceeding exactly as above we find

$$P^{(1)}(d^{(1)})P^{(2)}(d^{(2)}) = P(d) \quad (2.62)$$

where

$$P(d) = \prod_{k \in D, i=1}^Y d_{k,i}^+ d_{k,i} \exp \left[\frac{t}{2M^2} \sum_{k \in D, i=1}^X \text{tr} B(k) \text{tr} B^{(+)}(k) \right] \quad (2.63)$$

$$B(k) = \frac{i \sin k + \sin k_0}{i \sin k} \text{tr} m(k) \quad \text{tr} m(k) = \frac{i \sin k + \sin k_0}{i \sin k} \quad (2.64)$$

$$\text{tr} m(k) = \frac{i \sin k + \sin k_0}{i \sin k} \quad \text{tr} m(k) = \frac{i \sin k + \sin k_0}{i \sin k}$$

Note that $\text{tr} m(k) = 2 \left(\frac{1}{2} + 1 + t \right) + 2t(2 \cos k_0 \cos k_1)$, and the mass of the fields never vanishes. It holds that

$$P(d)_{x,i}^+ y,i = g_{i,i}^{(+)}(x-y); \quad (2.65)$$

where

$$g_{i,i}^{(+)}(x-y) = \frac{1}{2tM^2} \sum_k e^{ik(x-y)} B^{-1}(k)_{i,i}^{(+)}; \quad (2.66)$$

and $B^{-1}(k)$ is the inverse of the $B(k)$ defined in (2.63).

If we set

$$A(k) = \det B(k) = \sin^2 k_0 \sin^2 k [\text{tr} m(k)]^2; \quad (2.67)$$

then

$$B^{-1}(k) = \frac{1}{A(k)} \frac{\sin k_0 + i \sin k}{i \sin k} \text{tr} m(k) : \quad (2.68)$$

Similar formulas hold for $g^{(-)}(x-y)$ and, if $i = j$, the following bounds holds, for any $N > 1$

$$|g_{i,i}^{(i)}(x-y)| \leq \frac{1}{1 + |d_M(x-y)| + |d_M(x-y)|^N} C_N \quad (2.69)$$

$$|g_{i,i}^{(i)}(x-y)| \leq \frac{|d_M(x-y)|^N C_N}{1 + |d_M(x-y)|^N} \quad (2.70)$$

where

$$d_M(x-y) = \left(\frac{M}{2} \sin \left(\frac{(x-y)}{M} \right) \right)^2; \quad \left(\frac{M}{2} \sin \left(\frac{(x_0-y_0)}{M} \right) \right)^2 \quad (2.71)$$

The same transformations are done for the term $e^{S_{\pi^0}}$ which can be written as the product of three terms similar to (2.33), (2.35) and (2.36) with t replaced by $\frac{p}{2} - 1$ in (2.33) and $\frac{p}{2} + 1$ in (2.35) replaced by 0; then (2.33), (2.35) can be written in terms of Dirac fermions as (2.61) and (2.64) with t replaced by $\frac{p}{2} - 1$ or $\frac{p}{2} + 1$ replaced by 0.

We have then written the partition function as Grassmann integral over the ψ -elds, which are massive with a mass $O(1)$, and the $\bar{\psi}$ -elds with a small mass $O(t - t_c)$ close to the critical temperature; we integrate out first the heavy fermions to get an effective theory in terms of the light fermions only.

3. Integration of heavy fermions

3.1 Local interactions. By the change of variables in the preceding section we can write (2.23) as

$$Z_{2I}^{\pi^{(1)}; \pi^{(1)}; \pi^{(2)}; \pi^{(2)}} = e^{M^2 N} \int P(d\psi) \int P(d\bar{\psi}) e^{Q(\psi; \bar{\psi})} e^{Q(\bar{\psi}; \psi)} e^{V(\psi; \bar{\psi})} \quad (3.1)$$

where N is a normalization constant, $P(\psi)$ and $P(d\bar{\psi})$ are given by (2.60), (2.63), V, Q, \bar{Q} are obtained respectively from (2.36) and (2.46) by the change of variables (2.59). We write

$$V(\psi; \bar{\psi}) = V_2(\psi) + V_4(\psi) + V(\psi; \bar{\psi}) \quad (3.2)$$

where $V_2(\psi)$ is a sum of monomials bilinear in the ψ variables, $V_4(\psi)$ is a sum of monomials quartic in the ψ and $V(\psi; \bar{\psi})$ is sum of monomials bilinear or quartic each one containing at least a $\bar{\psi}$ eld. It holds that, from (2.31),

$$H_x^{(1)} H_x^{(1)} = \frac{i}{2} \psi_x^{(1)} \psi_x^{(1)} + R_1^{(1)} \quad V_x^{(1)} V_x^{(1)} = \frac{i}{2} \psi_x^{(1)} \psi_x^{(1)} + R_2^{(1)} \quad (3.3)$$

where $R_1^{(1)}$ is sum of monomials bilinear in the elds $\psi; \bar{\psi}; \psi$ and containing at least one eld $\bar{\psi}$ or ψ . Moreover, from (2.58)

$$\psi_x^{(1)} \psi_x^{(1)} = \frac{1}{2} [\psi_{1;x} \psi_{1;x} + \psi_{1;x}^+ \psi_{1;x}^+ + \psi_{1;x}^+ \psi_{1;x} + \psi_{1;x} \psi_{1;x}^+] \quad (3.4)$$

$$\psi_x^{(2)} \psi_x^{(2)} = \frac{1}{2} [\psi_{1;x} \psi_{1;x} + \psi_{1;x}^+ \psi_{1;x}^+ + \psi_{1;x}^+ \psi_{1;x} + \psi_{1;x} \psi_{1;x}^+]$$

hence

$$\begin{aligned} \psi_x^{(1)} \psi_x^{(1)} \psi_x^{(2)} \psi_x^{(2)} &= 0 \\ \psi_x^{(1)} \psi_x^{(1)} \psi_x^{(2)} \psi_x^{(2)} &= e^{2i(p_{\pi^{(1)}}; \pi^{(1)} + p_{\pi^{(1)}}; \pi^{(1)})x} \psi_{1;x}^+ \psi_{1;x}^+ \psi_{1;x} \psi_{1;x} \end{aligned} \quad (3.5)$$

and from (2.26)

$$V_4(\vec{r}) = 2e^{2i(p_{n(1)}^{(1)} + p_{n(1)}^{(1)}) \cdot \vec{r}} (\tilde{a} + \tilde{b})^X \prod_{\vec{k}}^{+} \prod_{\vec{k}}^{+} \prod_{\vec{k}}^{+} \prod_{\vec{k}}^{+} + V_4^R(\vec{r}) \quad (3.6)$$

and V_4^R is a sum of quartic monomials with coupling $O(\epsilon)$ in which at least a ∂ eld. In the same way it is easy to check that

$$V_2 = \frac{1}{2M^2} \sum_{\vec{k}}^X f[\vec{r} + \vec{f}_1(\vec{r})][i! + \cos k_0 + \cos k_1] \prod_{\vec{k}}^{+} \prod_{\vec{k}}^{+} + [\vec{r} + \vec{f}_2(\vec{r})][i! \sin k_0 + i \sin k] \prod_{\vec{k}}^{+} \prod_{\vec{k}}^{+} g \quad (3.7)$$

where $\vec{f}_1, \vec{f}_2 \in C^j$.

3.2 Integration of the heavy elds We now integrate the heavy elds

$$Z P(d) e^{M^{2N(1)} + V^{(1)}(\vec{r})} = \int P(d) \int P(d) e^{Q(\vec{r})} e^{G(\vec{r})} e^{V(\vec{r})} \quad (3.8)$$

where $N^{(1)}$ is a constant. Let us consider a set of coordinates x_1, \dots, x_{2n} which are not all different one to the other and we will denote by $\sum_{x_1, \dots, x_{2n}}$ the sum over all the distinct $2n$ variables. We prove the following result.

Theorem 2. There exists an ϵ such that, for $j \geq j_0$

$$V^{(1)} = \sum_{n=1}^X \sum_{\vec{r}}^X \sum_{\vec{r}}^X W_n(x_1, \dots, x_{2n}) \partial^{-1} \prod_{x_1, \dots, x_{2n}}^{n_1} \dots \partial^{-2n} \prod_{x_{2n}, \dots, x_{2n}}^{n_{2n}} \quad (3.9)$$

and, for $n > 2$

$$\sum_{x_1, \dots, x_{2n}}^X |W_n(x_1, \dots, x_{2n})| \leq M^{2C} n^j j^{n=2} \quad (3.10)$$

For $n = 2$

$$V_2(x_1, \dots, x_{2n}) \partial^{-1} \prod_{x_1, \dots, x_{2n}}^{n_1} \dots \partial^{-2n} \prod_{x_{2n}, \dots, x_{2n}}^{n_{2n}} = \sum_{x_1, \dots, x_4}^X \sum_{\vec{r}}^X W_2(x_1, \dots, x_4) \partial^{-1} \prod_{x_1, \dots, x_4}^{n_1} \dots \partial^{-2n} \prod_{x_{2n}, \dots, x_{2n}}^{n_{2n}} \quad (3.11)$$

with $\sum_{x_1, \dots, x_4}^P |W_2(x_1, \dots, x_4)| \leq M^{2C} j$ and $\hat{W}_2(0, \dots, 0) = O(\epsilon^2)$; for $n = 1$

$$\sum_{x_1, x_2}^X \sum_{\vec{r}}^X W_1(x_1, x_2) \partial^{-1} \prod_{x_1}^{n_1} \dots \partial^{-2} \prod_{x_2}^{n_2} = \sum_{x_1, x_2}^X f(\vec{r}) \prod_{\vec{r}} g$$

$$V_2(\varphi) + \sum_{x_1, x_2 \in \mathbb{Z}^D} \sum_{f \in \mathcal{F}} W_1(x_1; x_2) @^{(1)}_{x_1; f} @^{(2)}_{x_2; f} \quad (3.12)$$

with $\sum_{x_1, x_2} W_1(x_1; x_2) j = M^2 C j$; V_2 and V_4 are given by (3.6) and (3.7). Finally

$$P(\varphi) = \sum_{k \in \mathbb{Z}^D} \sum_{f \in \mathcal{F}} d_{k; f}^+ d_{k; f} \exp \left[\frac{t}{M^2} \sum_{k \in \mathbb{Z}^D} \sum_{f \in \mathcal{F}} d_{k; f}^+ \tilde{A}_{f; 0}(k) d_{k; f} \right] \quad (3.13)$$

with

$$\tilde{A}(k) = \frac{1}{C_0 + f_{0;0}(k)} \left[Z(i \sin k + \sin k_0) + f_{1;1}(k) i(t - t_c) C_0 + f_{1;2}(k) Z(i \sin k - \sin k_0) + f_{2;2}(k) \right]$$

with $C_0 = (t + 1 + \frac{P}{2})^2$ and $Z = \frac{t}{2} [(2t + 2\frac{P}{2}t) + (2\frac{P}{2} + 3 + t^2)]$ and $f_{0;0}(k)$, $f_{i;j}(k)$, $i, j = 1, 2$ analytic $O(k^2)$ functions.

Remark 1. The fact that $\hat{W}_2(0; \dots; 0) = O(k^2)$ can be checked by an explicit computations of all the contributions with coupling $O(k)$ to W_2 , noting that they can be only obtained contracting a term s quartic in the fields with one of the addend of (2.36); each of such term s carries a derivative in the coordinate space, hence the Fourier transform of such term s is vanishing at zero momentum.

Remark 2. The fermionic integration $P(\varphi)$ reduces, neglecting the functions $f_{0;0}(k)$, $f_{i;j}(k)$ which are $O(k^2)$, to the integration of a system of Dirac fermions on a lattice with mass $O(t - t_c)$. Hence, apart from the functions $f_{0;0}(k)$, $f_{i;j}(k)$, it coincides with the free action of the Thirring model. Note however that the interaction in the Thirring model is quartic and invariant under the transformation $\psi_i \rightarrow e^{i\theta} \psi_i$ in the massless case; both this properties are not true for (3.9).

Proof. We start from the definition of truncated expectation:

$$E^T(X; n) = \frac{\partial^n}{\partial^n} \log \sum_{\varphi} P(\varphi) e^{X(\varphi)} \Big|_{j=0} \quad (3.14)$$

so that, calling

$$V(\varphi) = Q(\varphi) + \tilde{Q}(\varphi) + V(\varphi) \quad (3.15)$$

we obtain

$$M^2 N^{(1)} + V^{(1)}(\varphi) = \log \sum_{\varphi} P(\varphi) e^{V(\varphi)} = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} E^T(V(\varphi); n) \quad (3.16)$$

We label each one of the monomials (whose number will be called C_0) in V by an index v_i , so that each monomial can be written as

$$\sum_{x_{v_i}} \prod_{f \in \mathcal{F}_{v_i}} v(x_{v_i}) @^{(f)}_{x_{v_i}; f} \prod_{f \in \mathcal{F}_{v_i}} @^{(f)}_{x_{v_i}; f} \quad (3.17)$$

where x_{v_i} is the total set of coordinates associated to v_i and P_{v_i} and \mathcal{P}_{v_i} are set of indices labeling the or - elds. We can write

$$V^{(1)}(\cdot) = \sum_{\mathcal{P}_{v_0} \in 0} V^{(1)}(\mathcal{P}_{v_0}); \quad (3.18)$$

$$V^{(1)}(\mathcal{P}_{v_0}) = \sum_{x_{v_0} \in \mathcal{P}_{v_0}} \sum_{\substack{Y \\ \text{at } (f) \\ \text{! } (f); x(f)}} \sum_{\substack{X \\ \text{! } (f); x(f)}} K_{\mathcal{P}_{v_0}}^{(0)}(x_{v_0}); \quad (3.19)$$

$$K_{\mathcal{P}_{v_0}}^{(0)}(x_{v_0}) = \sum_{n=1}^X \frac{1}{n!} \sum_{v_1, \dots, v_n} E^T[\sim(P_{v_1}); \dots; \sim(P_{v_n})] \sum_{i=1}^Y v_i(x_{v_i}); \quad (3.20)$$

where $\mathcal{P}_{v_1, \dots, v_n} = \mathcal{C}_0^n, \mathcal{P}_{v_0} = \sum_i \mathcal{P}_{v_i}$ and $x_{v_0} = \sum_i x_{v_i}$. We use now the wellknown expression for E^T (see for instance [Le])

$$E^T(\sim(P_1); \dots; \sim(P_s)) = \sum_{T \in \mathcal{T}} \sum_{\substack{Y \\ \text{at } (f) \\ \text{! } (f); x(f)}} \sum_{\substack{X \\ \text{! } (f); x(f)}} g_{\substack{! \\ +}}(x_1 \dots x_Y) dP_T(t) \det G^T(t) \quad (3.21)$$

where:

→ \mathcal{P} is a set of indices, and

$$\sim(P) = \sum_{f \in \mathcal{P}} \sum_{\substack{Y \\ \text{at } (f) \\ \text{! } (f); x(f)}} \sum_{\substack{X \\ \text{! } (f); x(f)}} \quad (3.22)$$

→ \mathcal{T} is a set of lines forming an anchored tree between the cluster of points $P_1; \dots; P_s$ i.e. \mathcal{T} is a set of lines which becomes a tree if one identifies all the points in the same clusters.

→ $t = ft_{i,i^0} \in [0;1]; 1 \leq i, i^0 \leq s$, $dP_T(t)$ is a probability measure with support on a set of t such that $t_{i,i^0} = u_i \cdot \psi$ for some family of vectors $u_i \in \mathbb{R}^s$ of unit norm.

→ $G^T(t)$ is a $(n \times s + 1) \times (n \times s + 1)$ matrix, whose elements are given by $G_{ij,i^0j^0}^T = t_{i,i^0} g_{\substack{! \\ +}}(x_{ij} \dots y_{i^0j^0})$ with $(f_{ij}; f_{i^0j^0}^+)$ not belonging to \mathcal{T} .

If $s = 1$ the sum over \mathcal{T} is empty, but we can still use the above equation by interpreting the r.h.s. as 1 if \mathcal{P}_1 is empty, and $\det G(\mathcal{P}_1)$ otherwise.

We bound the determinant using the well known Gram-Hadamard inequality, stating that, if M is a square matrix with elements M_{ij} of the form $M_{ij} = \langle A_i; B_j \rangle$, where A_i, B_j are vectors in a Hilbert space with scalar product $\langle \cdot; \cdot \rangle$, then

$$|\det M| \leq \sum_i \|A_i\| \|B_i\| \quad (3.23)$$

where $\| \cdot \|$ is the norm induced by the scalar product.

Let $H = \mathbb{R}^s \otimes H_0$, where H_0 is the Hilbert space of complex four dimensional vectors $F(k) = (F_1(k); \dots; F_4(k))$, $F_i(k)$ being a function on the set D , with scalar product

$$\langle F; G \rangle = \sum_{i=1}^4 \frac{1}{M^2} \sum_k F_i(k) G_i(k) \quad (3.24)$$

and it is easy to verify that

$$G_{ij;i^0j^0}^T = t_{i;i^0} g_{!_1; !_1}^{(+)}(\mathbf{x}_{ij} \quad \mathbf{y}_{i^0j^0}) = \langle u_i \quad A_{\mathbf{x}(\mathbf{f}_{ij});!(\mathbf{f}_{ij})}; u_{i^0} \quad B_{\mathbf{x}(\mathbf{f}_{i^0j^0});!(\mathbf{f}_{i^0j^0})}^+ \rangle; \quad (3.25)$$

where $u_i \in \mathbb{R}^s$, $i = 1; \dots; s$, are the vectors such that $t_{i;i^0} = u_i \cdot \psi$, and

$$\begin{aligned} A_{\mathbf{x};!}(\mathbf{k}) &= e^{ik^0x} \mathcal{P} \frac{1}{A(\mathbf{k})} \begin{cases} (\sin k_0 + i \sin \mathbf{k}; 0; \sin(\mathbf{k}); 0); & \text{if } ! = +1, \\ (0; \sin(\mathbf{k}); 0; \sin(\mathbf{k})); & \text{if } ! = -1, \end{cases} \\ B_{\mathbf{x};!} &= e^{ik^0y} \mathcal{P} \frac{1}{A(\mathbf{k})} \begin{cases} (1; 1; 0; 0); & \text{if } ! = +1, \\ (0; 0; 1; (\sin k_0 + i \sin \mathbf{k}) = \sin(\mathbf{k})); & \text{if } ! = -1. \end{cases} \end{aligned} \quad (3.26)$$

Hence from (3.23) we immediately find

$$\mathcal{G}_{ij;i^0j^0}^T \in C_1^n \quad (3.27)$$

Note that C_1 is an $O(1)$ constant; this follows from the fact that the Grassmann variable ψ has an $O(1)$ mass. Finally we get

$$\sum_{\mathbf{x}_{v_0}} \mathcal{K}_{\mathcal{P}_{v_0}}^{(1)}(\mathbf{x}_{v_0}) \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{v_1; \dots; v_n} \sum_{\mathbf{x}_{v_1}; \dots; \mathbf{x}_{v_n}} C_1^n \left[\sum_{T \in \mathcal{T}} \mathcal{G}(\mathbf{x}_1 \quad \mathbf{y}_1) \right] \prod_{i=1}^n \mathcal{J}_i(\mathbf{x}_{v_i}) \quad (3.28)$$

where we have used that $\int d\mathcal{P}_T(t) = 1$. Regarding the sum over T , it is empty if $n = 1$. If $n > 1$ the number of anchored trees with d_i lines branching from the vertex v_i can be bounded, by using Cayley's formula, by

$$\frac{(s_v - 2)!}{(d_1 - 1)! \dots (d_{s_v} - 1)!} \mathcal{P}_{v_1}^{d_1} \dots \mathcal{P}_{v_n}^{d_n}; \quad (3.29)$$

hence the number of addenda in \sum_T is bounded by $n! C_2^n$. Finally T and the \mathbf{x}_{v_i} form a tree connecting all points, so that using that the propagator is massive and that the interactions are short ranged $\sum_{\mathbf{x}_{v_1}; \dots; \mathbf{x}_{v_n}} \sum_{T \in \mathcal{T}} \left[\sum_{i=1}^n \mathcal{J}_i(\mathbf{x}_{v_i}) \right] C_3^n \mathcal{J}^M$, where n is the number of coupling $O(\cdot)$.

Let us consider the case $\mathcal{P}_{v_0} \in \mathcal{Q}$. Note that if to v_i are associated only terms from $V(\cdot; \cdot)$, then $n = n$. The same bound holds for M large enough, if there are v_i associated with \mathcal{Q} ; in fact such terms are vanishing as $M \rightarrow \infty$ (one has in the bounds an extra M^{-1} for any of such v_i , for dimensional reasons). Let us consider now the case in which there are end-points associated to $Q(\cdot; \cdot)$, which have $O(1)$ coupling; there are at most \mathcal{P}_{v_0} end-points associated with $Q(\cdot; \cdot)$. In fact in $Q(\cdot; \cdot)$ there are only terms of the form $\mathcal{J}^{\frac{n}{2}}$, so at most the number of them is equal to the number of fields. If we call n the number of vertices quartic in the fields it is clear that $n \leq \max\{1; \mathcal{P}_{v_0} \} \leq 2$; hence

$$\sum_{\mathbf{x}_{v_0}} \mathcal{K}_{\mathcal{P}_{v_0}}^{(1)}(\mathbf{x}_{v_0}) \mathcal{J}^M \sum_{n=1}^{\infty} C^{n+\mathcal{P}_{v_0}} \mathcal{J}^{\frac{n}{2}} \mathcal{J}^{\max\{\frac{1}{2}; \mathcal{P}_{v_0} \} - 1} \mathcal{J}^{2g} \quad (3.30)$$

and (3.10) holds for $\mathcal{P}_{v_0} j = 4$ (in the r.h.s. of (3.30) the sum over the number of v_i to which are associated quadratic monomials with coupling $O(\epsilon)$ is already done).

Consider now the case $\mathcal{P}_{v_0} j = 2$; in this case there are terms independent, obtained when to all the v_i are associated with elements of $Q(\epsilon; \epsilon)$. It is convenient to include all such terms in the free measure, as they cannot be considered as perturbations (they are not $O(\epsilon)$). Instead of computing all such terms, we can proceed in a more rapid way by noting that

$$\sum_{\mathbf{d}} P(\mathbf{d}) = \sum_{\mathbf{d}} P(\mathbf{d}) \sum_{\mathbf{d}'} P(\mathbf{d}') e^{Q(\mathbf{d}, \mathbf{d}')} \quad (3.31)$$

and if, if $\langle X \rangle_0 = \sum_{\mathbf{d}} P^0(\mathbf{d}) X$, it holds

$$\langle x_{i,1} + y_{i,1} \rangle_0 = \frac{1}{2} \langle (x^{(1)} + i x^{(2)}) (y^{(1)} - i y^{(2)}) \rangle_0 = \langle x^{(1)} y^{(1)} \rangle_0 \quad (3.32)$$

as $\langle x^{(1)} y^{(2)} \rangle_0 = 0$ if $\epsilon = 0$ (the two systems are decoupled). In the same way

$$\langle x_{i,1} + y_{i,1} \rangle_0 = \frac{1}{2} \langle (x^{(1)} + i x^{(2)}) (y^{(1)} - i y^{(2)}) \rangle_0 = \langle x^{(1)} y^{(1)} \rangle_0 \quad (3.33)$$

and finally

$$\langle x_{i,1} + y_{i,1} \rangle_0 = \frac{1}{2} \langle (x^{(1)} + i x^{(2)}) (y^{(1)} - i y^{(2)}) \rangle_0 = \langle x^{(1)} y^{(1)} \rangle_0 \quad (3.34)$$

Note that $\langle x y \rangle_0$ are the Ising correlation computed in [MPW], [SP] where is found that

$$\langle x^{(1)} y^{(1)} \rangle_0 = \frac{1}{M^2} \sum_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x} - \mathbf{y})}}{c_{1,1}(\mathbf{k})} \quad (3.35)$$

$$\langle x^{(1)} y^{(2)} \rangle_0 = \frac{1}{M^2} \sum_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x} - \mathbf{y})}}{c_{2,2}(\mathbf{k})} \quad (3.36)$$

$$\langle x^{(2)} y^{(1)} \rangle_0 = \frac{1}{M^2} \sum_{\mathbf{k}} \frac{e^{i\mathbf{k}(\mathbf{x} - \mathbf{y})}}{c_{1,2}(\mathbf{k})} \quad (3.37)$$

where

$$c_{1,1}(\mathbf{k}) = 2t(1 - t^2)(2 - \cos k_0 - \cos k) + (t + 1 + \frac{P}{2})^2 (t - \frac{P}{2} + 1)^2 \quad (3.38)$$

and

$$c_{1,1}(\mathbf{k}) = \frac{t}{2} [(2t + 2\frac{P}{2} - 2t)(\sin k_0 \cos k - i \sin k \cos k_0) + (2\frac{P}{2} + 3 + t^2)(\sin k_0 - i \sin k)] \quad (3.39)$$

$$c_{2;2}(k) = \frac{t}{2} [(2t + \frac{p}{2}) (\sin k_0 \cos k - i \sin k \cos k_0) + (\frac{p}{2} + 3 + t^2) (-\sin k_0 - i \sin k)] \quad (3.40)$$

$$c_{1;2}(k) = c_{1;2}(k) = \frac{i}{2} [(2\frac{p}{2}t^2 + 4t^2) \cos k \cos k_0 + (t^3 + 2t\frac{p}{2} + t) (\cos k + \cos k_0) - 2\frac{p}{2} + 2t^2] \quad (3.41)$$

Note that

$$\frac{c_{1;1}(k)c_{2;2}(k) - c_{1;2}(k)c_{2;1}(k)}{(k)} (t + 1 + \frac{p}{2})^2 = f(k)$$

where $f(k) = O(k^2)$ and with bounded derivatives.

4. Renormalization Group for light fermions

4.1 Multiscale analysis. We start considering $Z_{2I}^{j,j}$ in the sum (2.22), written as in (3.8). We introduce a scaling parameter $\lambda > 1$ and a positive function $(k) \in C^1$ such that

$$(k) = (k) = \begin{cases} 1 & \text{if } |k| < t_0 \\ 0 & \text{if } |k| > a_0 \end{cases}; \quad (4.1)$$

where

$$|k| = \frac{q}{\sin k_0^2 + \sin k^2}; \quad (4.2)$$

We define also, for any integer $h \geq 0$,

$$f_h(k) = (|k|^{-h}) - (|k|^{-h+1}); \quad (4.3)$$

we have, for any $h_M < 0$,

$$(k) = \sum_{h=h_M+1}^{X^0} f_h(k) + (|k|^{-h_M}); \quad (4.4)$$

Note that, if $h \geq 0$, $f_h(k) = 0$ for $|k| < t_0^{-h-1}$ or $|k| > t_0^{-h+1}$, and $f_h(k) = 1$, if $|k| = t_0^{-h}$. Therefore

$$f_h(k) = 0 \quad \forall h < h_M = \min \{h : t_0^{-h+1} > \frac{p}{2} (M^{-1})^2 g\}; \quad (4.5)$$

and

$$1 = \sum_{h=h_M}^{X^1} f_h(k) \quad f_1 = 1 \quad (k); \quad (4.6)$$

We define a sequence of effective potentials $V^{(h)}(\cdot)$ defined iteratively in the following way; assuming that we have integrated the scales $h = 1; 0; -1; -2; \dots; -h+1$

$$e^{-M^{-2}E_M} = \int_{\mathbb{Z}} P_{Z_h, m_h; C_h} (d^{(-h)}) e^{-V^{(h)}(\overline{Z_h^{(-h)}})} e^{-M^{-2}E_h}; \quad V^{(h)}(0) = 0; \quad (4.7)$$

where

$$P_{Z_h, m_h; C_h} (d^{(-h)}) = \prod_{k \in \mathbb{Z}_h^{-1}(k) > 0} \frac{d^{(-h)+}_{k;!} d^{(-h)}_{k^0;!}}{N(k)} \exp \left\{ \frac{1}{M^2} \sum_{k \in \mathbb{Z}_h^{-1}(k) > 0} C_h(k) Z_h \sum_{l=0}^{\infty} \frac{d^{(-h)+}_{k;!} T^{(h+1)}_{l;!^0} d^{(-h)}_{k;!^0}}{l!^0} \right\}; \quad (4.8)$$

$$C_h(k)^{-1} = \sum_{j=h_M}^{X^h} f_j(k^0); \quad (4.9)$$

and the 2×2 matrix $T_h(k^0)$ is given by

$$\frac{1}{C_0 + f_{0;0}(k)} \begin{pmatrix} Z(isink + \sin k_0) + f_{1;1}(k) Z_h^{-1} & \sin_{h-1}(k) + f_{1;2}(k) Z_h^{-1} \\ \sin_{h-1}(k) + f_{2;1}(k) Z_h^{-1} & Z(isink - \sin k_0) + f_{2;2}(k) Z_h^{-1} \end{pmatrix} \quad (4.10)$$

with $m_1 = C_0(t - t_c)$ and $C_0; Z$ defined in Theorem 2.

Finally $V^{(h)}$ is given by

$$V^{(h)}(\cdot^{(-h)}) = \sum_{n=1}^X \sum_{\substack{x_1, \dots, x_{2n} \\ \underline{j} \in \mathbb{Z}}} \sum_{i=1}^{2^n} \mathcal{C}_{\substack{i \\ x_i; !^i}}^{(-h)} W_{2n; \underline{j} \in \mathbb{Z}}^{(h)}(x_1; \dots; x_{2n}) \quad (4.11)$$

4.2 The localization operator. We define an L operation, for $h \geq 0$, in the following way:

1) If $2n = 4$, then

$$L \hat{W}_{4; \underline{j} \in \mathbb{Z}}^{(h)}(k_1; k_2; k_3) = \hat{W}_{4; \underline{j} \in \mathbb{Z}}^{(h)}(k_{++}; k_{++}; k_{++}); \quad (4.12)$$

where

$$k_{++} = \frac{1}{M}; \quad \frac{0}{M} : \quad (4.13)$$

2) If $2n = 2$ then

$$L \hat{W}_{2; \underline{j} \in \mathbb{Z}}^{(h)}(k) = \frac{1}{4} \sum_{i=0}^X \hat{W}_{2; \underline{j} \in \mathbb{Z}}^{(h)}(k_{++}^i)$$

$$+\hat{W}_{2;-;!}^{(h)}(k_0) \left(\frac{L}{2} \sin k + \frac{0}{2} - \sin k_0 \right) \quad (4.14)$$

3) In all the other cases

$$L\hat{W}_{2n;-;!}^{(h)}(k_1; \dots; k_{2n-1}) = 0 : \quad (4.15)$$

By (4.12) the operator L satisfies the relation $RL = 0$.

Remark. First note that in the limit $M \rightarrow 1$ (4.14) becomes simply

$$L\hat{W}_{2;-;!}^{(h)}(k) = [\hat{W}_{2;-;!}^{(h)}(0) + k_0 \partial_{k_0} \hat{W}_{2;-;!}^{(h)}(0) + k \partial_k \hat{W}_{2;-;!}^{(h)}(0)]; \quad (4.16)$$

hence $L\hat{W}_{2;-;!}^{(h)}(k)$ has to be understood as a discrete version of the Taylor expansion up to order 1. Moreover the localization operator acting on the effective potential in the x -space representation can be seen as an operator on monomials of Grassmann variables; for instance (4.12) implies in the x -space representation, in the $M \rightarrow 1$ limit

$$L \prod_{x_1; \dots; x_4}^{Y^4} \frac{(h)^{n_i}}{x_i!} W_4^{(h)}(x_1; \dots; x_4) = \hat{W}_4(0; 0; 0) \prod_{i=1}^{Y^4} \frac{(h)^{n_i}}{x_i!} \quad (4.17)$$

where the r.h.s. of the above equation is always vanishing unless $\prod_{i=1}^4 \frac{(h)^{n_i}}{x_i!}$ is a permutation of $\frac{(h)^{+}}{x_1!} \frac{(h)^{+}}{x_1!} \frac{(h)^{+}}{x_1!} \frac{(h)^{+}}{x_1!}$; hence L acts on a quartic monomial producing a local expression. Analogous considerations can be done for $n = 1$.

4.3 We have, before continuing, to exploit the consequences of some symmetries. There are no local terms of the form $\frac{+}{x_1!} \frac{+}{x_1!}$; such local terms can be written as $\frac{(1)}{x} \frac{(2)}{x}$, but the model is invariant under the transformation

$$\begin{aligned} & \frac{(1)}{x}; \frac{(1)}{x}; \frac{(1)}{x}; \frac{(1)}{x}! \quad \frac{(1)}{x}; \frac{(1)}{x}; \frac{(1)}{x}; \frac{(1)}{x} \\ & \frac{(2)}{x}; \frac{(2)}{x}; \frac{(2)}{x}; \frac{(2)}{x}! \quad \frac{(2)}{x}; \frac{(2)}{x}; \frac{(2)}{x}; \frac{(2)}{x}; \end{aligned} \quad (4.18)$$

hence such terms cannot be present as they violate such symmetry.

There are no local terms of the form $\frac{+}{1;x} \frac{+}{1;x}$ or $\frac{+}{1;x} \frac{+}{1;x}$ (or $\frac{+}{1;x} \otimes \frac{+}{1;x}$, $\frac{+}{1;x} \otimes \frac{+}{1;x}$); in fact,

$$\frac{1}{1;x} \frac{1}{1;x} = \frac{1}{2} \left[\frac{(1)}{x} \frac{(1)}{x} \frac{(2)}{x} \frac{(2)}{x} + i \frac{(1)}{x} \frac{(2)}{x} + i \frac{(2)}{x} \frac{(1)}{x} \right] \quad (4.19)$$

and the last two terms violate the symmetry (4.18); moreover the first two terms are odd in the exchange $\frac{(1)}{x}; \frac{(2)}{x}! \rightarrow \frac{(2)}{x}; \frac{(1)}{x}$ and the model is invariant in the exchange $\frac{(1)}{x}; \frac{(2)}{x}! \rightarrow \frac{(2)}{x}; \frac{(1)}{x}$. Moreover the model is invariant under complex conjugation and the exchange

$$\frac{(\)}{x}; \frac{(\)}{x}! \rightarrow \frac{(\)}{x}; \frac{(\)}{x} \quad \frac{(\)}{x}; \frac{(\)}{x}! \rightarrow \frac{(\)}{x}; \frac{(\)}{x}; \quad (4.20)$$

this follows from the fact that, from (2.31), $H^{(1)}; H^{(2)}; V^{(1)}; V^{(2)}$, written in terms of $(1); (2); (3); (4)$, are invariant under such transformation. Hence the coefficient of the local part of the quartic (non vanishing) terms is real; in fact $\hat{w}(0;0;0)_{1;x}^{(1)}_{1;x}^{(1)}_{1;x}^{(2)}_{1;x} = \hat{w}(0;0;0)_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}^{(1)}_{x}$ must be equal, by the above invariance, to $\hat{w}(0;0;0)_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}^{(1)}_{x}$, hence $\hat{w}(0;0;0) = \hat{w}(0;0;0)$. Finally the combination of local terms $\hat{w}(0;0;0)_{x;1}^{(1)}_{x;1}^{(1)}_{x;1}^{(2)}_{x;1} + \hat{w}(0;0;0)_{x;1}^{(1)}_{x;1}^{(1)}_{x;1}^{(2)}_{x;1}$ is equal to $\frac{1}{2} [\hat{w}(0;0;0)_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}^{(1)}_{x} + \hat{w}(0;0;0)_{x}^{(2)}_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}]$ so it cannot be present as it violates the symmetry (4.18). On the other hand $\hat{w}(0;0;0)_{x;1}^{(1)}_{x;1}^{(1)}_{x;1}^{(2)}_{x;1} + \hat{w}(0;0;0)_{x;1}^{(1)}_{x;1}^{(1)}_{x;1}^{(2)}_{x;1}$ is equal to $\frac{1}{2} [\hat{w}(0;0;0)_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}^{(1)}_{x} + \hat{w}(0;0;0)_{x}^{(2)}_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}]$; hence the coefficient of the local part is imaginary; in fact $\hat{w}(0) [\hat{w}(0;0;0)_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}^{(1)}_{x} + \hat{w}(0;0;0)_{x}^{(2)}_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}]$ must be equal to $\hat{w}(0) [\hat{w}(0;0;0)_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}^{(1)}_{x} + \hat{w}(0;0;0)_{x}^{(2)}_{x}^{(1)}_{x}^{(1)}_{x}^{(2)}_{x}]$, by the invariance under complex conjugation and (4.20), hence $\hat{w}(0) = \hat{w}(0)$.

4.4 By (4.12), (4.14), (4.15) and the symmetry relations in x4.3, we can write $LV^{(h)}$ in the following way:

$$LV^{(h)}(\vec{h}) = (s_h + i n_h) F^{(h)} - l_h F^{(h)} + z_h F^{(h)} + i F^{(h)} + \sum_{i=1}^X \frac{X^2}{2} \frac{X}{z_{i;1}} F_i^{(h)} + \sum_{i=1}^X \frac{X^2}{2} \frac{X}{z_{i;1}} F_i^{(h)} \quad (4.21)$$

where, if $j \neq j'$, $z_{j;1} z_{j';1} = 0$, $s_1 = 0$, $l_1 = 2 \operatorname{sech}^2 J(a+b) + 0$, $z_{1;1} = 1 + 0$ and

$$\begin{aligned} F_m^{(h)} &= \sum_{k=1}^X \frac{i!}{M^2} \frac{X}{k!} \hat{F}_m^{(h)}(\vec{k}); \\ F^{(h)} &= \frac{1}{(M)^4} \sum_{k_1, \dots, k_4 \in \mathbb{Z}^{D_M}} \hat{F}^{(h)}(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4); \\ F^{(h)} &= \frac{1}{M} \sum_{k \in \mathbb{Z}^{D_M}} i \sin k \hat{F}^{(h)}(\vec{k}); \quad F^{(h)} = \frac{1}{M} \sum_{k \in \mathbb{Z}^{D_M}} ! \sin k_0 \hat{F}^{(h)}(\vec{k}); \\ F_i^{(h)} &= \frac{1}{M} \sum_{k \in \mathbb{Z}^{D_M}} f_i(k) \hat{F}_i^{(h)}(\vec{k}); \quad F_i^{(h)} = \sum_{k \in \mathbb{Z}^{D_M}} f_i \hat{F}_i^{(h)}(\vec{k}); \end{aligned} \quad (4.22)$$

with $f_1 = \sin k$ and $f_2 = \sin k_0$. The constants $n_h; s_h; l_h; z_h; i_h$ are real. At the end of our iterative construction it will appear that it is possible to write (see Remark 1 in x4.9)

$$\hat{W}_{2; i; 1}^{(h)} = \hat{W}_{2; i; 1}^{a(h)} + \hat{W}_{2; i; 1}^{b(h)} \quad (4.23)$$

with $\hat{W}_{2; i; 1}^{a(h)}$ vanishing if at least one $m_k = 0$, for $1 \leq k \leq h+1$, and $\hat{W}_{2; i; 1}^{b(h)}$ is the rest; we define

$$s_h = i; ! \left[\frac{1}{4} \sum_{k=0}^X \hat{W}_{2; i; 1}^{a(h)}(k_0) \right]; \quad n_h = i; ! \left[\frac{1}{4} \sum_{k=0}^X \hat{W}_{2; i; 1}^{b(h)}(k_0) \right]; \quad (4.24)$$

In the same way we include in $\prod_{i=1}^P z_i; !; h F_i^{(h)}$ all the terms not in $z_h F^{(h)} + z_h F^{(h)}$ with $z_h; h$ real.

We renormalize the free integration $P_{Z_h m_h; C_h}(d^{(h)})$ by adding to it part of the r.h.s. of (4.21). We get

$$\begin{aligned} & \int P_{Z_h m_h; C_h}(d^{(h)}) e^{-V^{(h)}(\overline{Z_h}^{(h)})} = \\ & = e^{-L t_h} \int P_{Z_{h-1} m_{h-1}; C_h}(d^{(h)}) e^{-V^{(h)}(\overline{Z_h}^{(h)})}; \end{aligned} \quad (4.25)$$

where $P_{Z_{h-1} m_{h-1}; C_h}(d^{(h)})$ is obtained from $P_{Z_h m_h; C_h}(d^{(h)})$ by substituting Z_h with

$$\tilde{Z}_{h-1}(k) = Z_h [1 + C_h^{-1}(k) \tilde{Z}^{-1}(C_0 + f_{0;0}(k)) Z_h] \quad (4.26)$$

and $m_h(k)$ with

$$m_{h-1}(k) = \frac{Z_h}{\tilde{Z}_{h-1}(k)} [m_h(k) + C_h^{-1}(k) (C_0 + f_{0;0}(k)) s_h]; \quad (4.27)$$

moreover $V^{(h)} = V^{(h)} - Z_h s_h F^{(h)} - Z_h z_h (F^{(h)} + F^{(h)})$. We will call $m_h(0)$ m_h . The r.h.s of (4.25) can be written as

$$e^{-M^2 t_h} \int P_{Z_{h-1} m_{h-1}; C_{h-1}}(d^{(h-1)}) \int P_{Z_{h-1} m_{h-1}; f_h^{-1}}(d^{(h)}) e^{-V^{(h)}(\overline{Z_h}^{(h)})}; \quad (4.28)$$

where the factor $\exp(-M^2 t_h)$ in (4.25) takes into account the different normalization of the two integrations and

$$\tilde{Z}_{h-1} = Z_h (1 + C_0 \tilde{Z}^{-1} Z_h); \quad \tilde{f}_h(k) = Z_{h-1} \left[\frac{C_h^{-1}(k)}{\tilde{Z}_{h-1}(k)} - \frac{C_{h-1}^{-1}(k)}{Z_{h-1}} \right]; \quad (4.29)$$

Note that $\tilde{f}_h(k)$ has the same support of $f_h(k)$. The single scale propagator is

$$\int P_{Z_{h-1} m_{h-1}; f_h^{-1}}(d^{(h)}) \frac{g_{!; !^0}^{(h)}(x-y)}{x; !^0 y; !^0} = \frac{g_{!; !^0}^{(h)}(x-y)}{Z_{h-1}}; \quad (4.30)$$

where

$$g_{!; !^0}^{(h)}(x-y) = \frac{1}{M^2} \sum_k e^{-ik(x-y)} \tilde{f}_h(k) [T_h^{-1}(k)]_{!; !^0}; \quad (4.31)$$

and $T_h^{-1}(k)$ is the inverse of the $T_h(k)$ defined in (4.10). The large distance behaviour of $g_{!; !^0}^{(h)}(x-y)$ is given by, if $|x-y| \geq \frac{1}{2}$ and $\sup_k |x-y| \leq \frac{1}{2}$ $e^{|x-y|}$, given the positive integers $N; n_0; n_1$ and putting $n = n_0 + n_1$, and the constant $C_{N; n}$

$$|g_{x_0}^{n_0} g_x^{n_1} g_{!; !^0}^{(h)}(x-y)| \leq C_{N; n} \frac{e^{h+n}}{1 + (|x-y|)^N} \quad (4.32)$$

$$\| \mathcal{D}_{x_0}^{n_0} \mathcal{D}_x^{n_1} g_{!; !}^{(h)}(x-y) \| \leq C_{N,m} \frac{j_h^m}{1 + (j_h(x-y))^N} : \quad (4.33)$$

where \mathcal{D}_x denotes the discrete derivative. It will be useful to write

$$g_{!; !}^{(h)}(x-y) = g_{L; !}^{(h)}(x-y) + g_{!; !}^{(h)}(x-y) + g_{!; !}^{(h)}(x-y) \quad (4.34)$$

with

$$g_{L; !}^{(h)}(x-y) = \frac{1}{M^2} \sum_k e^{ik(x-y)} \frac{C_0 f_h(k)}{Z^2 \sin k_0 + iZ \sin k} ; \quad (4.35)$$

obeying to the bound (4.32) while

$$\| \mathcal{D}_{x_0}^{n_0} \mathcal{D}_x^{n_1} g_{!; !}^{(h)}(x-y) \| \leq C_{N,m} \frac{j_h^{\frac{3}{2}h+n}}{1 + (j_h(x-y))^N} \quad (4.36) :$$

$$\| \mathcal{D}_{x_0}^{n_0} \mathcal{D}_x^{n_1} g_{!; !}^{(h)}(x-y) \| \leq C_{N,m} \frac{j_h^m}{1 + (j_h(x-y))^N} : \quad (4.37)$$

Moreover

$$g_{!; !}^{(h)}(x-y) = g_{!; !}^{(h)}(x-y) + g_{!; !}^{(h)}(x-y) \quad (4.38)$$

with

$$g_{!; !}^{(h)}(x-y) = \frac{1}{M^2} \sum_k e^{ik(x-y)} f_h(k) \frac{iC_0 m_h(k)}{Z^2 \sin^2 k_0 + Z^2 \sin^2 k + m_h^2(k)} \quad (4.39)$$

verifying (4.33) and $g_{!; !}^{(h)}(x-y)$ verifying (4.36).

We now rescale the field so that

$$\hat{V}^{(h)} \left(\frac{\cdot}{Z_h} \right) = \hat{V}^{(h)} \left(\frac{\cdot}{Z_{h-1}} \right) ; \quad (4.40)$$

it follows that

$$L \hat{V}^{(h)}(\cdot) = \sum_{i=1}^h h F^{(h)}(\cdot) + \sum_{i=1}^h F^{(h)}(\cdot) + \sum_{i=1}^h F^{(h)}(\cdot) + \sum_{i=1}^h \sum_{j=1}^h F_i^{(h)}(\cdot) + \sum_{i=1}^h \sum_{j=1}^h \tilde{v}_{i; jh} F_i^{(h)}(\cdot) ; \quad (4.41)$$

where

$$h = \frac{Z_h}{Z_{h-1}} n_h ; \quad h = \frac{Z_h}{Z_{h-1}} (a_h - z_h) ; \quad h = \left(\frac{Z_h}{Z_{h-1}} \right)^2 l_h ; \quad (4.42)$$

and $\tilde{v}_{i; jh} = \frac{Z_h}{Z_{h-1}} z_{i; jh}$, $v_{i; jh} = \frac{Z_h}{Z_{h-1}} z_{i; jh}$. We call $v_h = (h; h; \sum_{i; jh} \tilde{v}_{i; jh}; \sum_{i; jh} v_{i; jh})$ and the set of $(v_h; m_h; n_h)$ are called running coupling constants.

If we now define

$$e^{V^{(h-1)} \left(\frac{\cdot}{Z_{h-1}} \right)} L E_h = \sum_{i=1}^Z P_{Z_{h-1} m_{h-1} i f_{h-1}}(d^{(h)}) e^{\hat{V}^{(h)} \left(\frac{\cdot}{Z_{h-1}} \right)} ; \quad (4.43)$$

it is easy to see that $V^{(h-1)}(\frac{p}{Z_{h-1}}(\cdot^{(h-1)}))$ is of the form (4.11) and that

$$E_{h-1} = E_h + t_h + \tilde{E}_h : \quad (4.44)$$

It is sufficient to use the well known identity

$$M^{-2} \tilde{E}_h + V^{(h-1)}(\frac{p}{Z_{h-1}}(\cdot^{(h-1)})) = \sum_{n=1}^{\infty} \frac{1}{n!} (1)^{n+1} E_h^{T,n} (\hat{V}^{(h)}(\frac{p}{Z_{h-1}}(\cdot^{(h)}))) ; \quad (4.45)$$

where $E_h^{T,n}$ denotes the truncated expectation of order n with propagator $Z_{h-1}^{-1} g_{!}^{(h)}$, see (4.30), and observe that $(\cdot^{(h)}) = (\cdot^{(h-1)}) + (\cdot^{(h)})$.

Let us define

$$h = \inf \{ h : 0 \leq h \leq h_M ; a_0^{h-1} \leq 4j_{n_h} ; 8h \leq 0 \leq h \leq h_g : \quad (4.46)$$

Of course this definition is meaningful only for m_0 such that $j_{n_0} \leq \frac{a_0}{4}$.

The integration of the scales from h to h_M will be performed "in a single step". This follows from the following result

4.5 Lemma. Assume that h is finite uniformly in M , so that $j_{n_{h-1}} \leq h$, for a suitable constant and define

$$\frac{g_{!}^{(h)}(x, y)}{Z_{h-1}} = \int_{Z_{h-1}^{-1} m_{h-1} C_h} (d(\cdot^{(h)})) \frac{(\cdot^{(h)})}{x!} \frac{(\cdot^{(h)})^+}{y!} : \quad (4.47)$$

Then, given the positive integers N, n_0, n_1 and putting $n = n_0 + n_1$, there exist a constant $C_{N,n}$ such that

$$| \int_{x_0}^{n_0} @_{x_1}^{n_1} g_{!}^{(h)}(x; y) | \leq C_{N,n} \frac{h+n}{1 + (\int (x-y))^{N-1}} : \quad (4.48)$$

4.6 Remark. Let us now explain the main motivations of the integration procedure discussed above. In a Renormalization Group framework one has to identify the relevant, marginal and irrelevant effective interactions. By a power counting argument one sees that the terms bilinear in the fields are relevant, and the terms quartic in the fields or quadratic with a derivated (in the x space) field are marginal. As it was shown in 4.3, there is only one kind of relevant term, and this is a rather crucial point. The Renormalization Group flow of the relevant terms can be controlled by the introduction of a counterterm; as we have at our disposal only one counterterm, it is important to have only one kind of relevant effective interactions. The unique relevant effective interaction has the form $\int_{x;1}^+ x; 1 \int_{x;1}^+ x; 1 x; 1$, which can be interpreted as a mass term in a fermionic

quantum field theory like the Thirring model. There is however an important difference: in fermionic models coming from QFT if there is no mass term in the formal free action then no mass terms are generated by Renormalization Group iterations, in absence of spontaneous symmetry breaking; the reason is that such models are invariant under the local gauge transformation $\psi(x) \rightarrow e^{i\theta(x)} \psi(x)$ if there is no mass term in the formal free action. In our model this is not true, as the interaction is not invariant under this symmetry; hence even if $t = t_c$ (or $m_1 = 0$) a mass term can be generated in the RG iterations. Hence we collect all the relevant terms which are vanishing if $m_k = 0, 1 \leq k \leq h+1$, in S_h , which we include in the fermionic free integration; the "mass" has a non trivial flow producing at the end the critical index of the correlation length. The remaining terms are left in the effective interaction; they are constituting the running coupling constant g_h (in a gauge invariant theory $g_h = 0$) whose flow is controlled by a suitable choice of the counterterm; the running coupling constant g_h takes into account the renormalization of the critical temperature. The running coupling constant v_h are dimensionally marginal; again the $\tilde{g}_{i,j,h}; g_{i,j,h}$ correspond to effective interactions which are absent from the free action, but which could be possibly generated by the RG iterations, and they would be absent in a QFT fermionic model. Finally, due to the mass gap, the propagator of the integration of all the scales between h and h_M has the same bound as the propagator of the integration of a single scale greater than h ; this property is used to perform the integration of all the scales h in a single step.

4.7 Theorem 3. Let $h > h_0$ and, for some constants c_1 , if

$$\max_{k \leq h} [j_k + j_k] \leq g_h; \sup_{h^0 > h} \frac{m_{h^0}}{m_{h^0-1}} \leq e^{c_1 g_h}; \sup_{h^0 > h} \frac{Z_{h^0}}{Z_{h^0-1}} \leq e^{c_1 g_h^2} \quad (4.49)$$

there exists a constant ϵ (depending on c_1) such that, if $g_h \leq \epsilon$, then, for a suitable constant c_0 , independent of c_1 , as well as of M , the kernels in (4.11) verify

$$\sum_{x_1, \dots, x_{2n}} |W_n^{(h)}(x_1, \dots, x_{2n})| \leq M^{2-hD_k(P_{v_0})} (c_0 g_h)^{\max(1, n-1)} \quad (4.50)$$

where

$$D_k(P_{v_0}) = 2 + n + k : \quad (4.51)$$

and $k = \sum_{i=1}^P 2n_i$.

4.8 Proof. We write $V^{(h)}$ in terms of a tree expansion, similar to that described, for example, in [BM].




Fig. 1

We need some definitions and notations.

1) Let us consider the family of all trees which can be constructed by joining a point r , the root, with an ordered set of $n - 1$ points, the endpoints of the unlabeled tree, so that r is not a branching point. n will be called the order of the unlabeled tree and the branching points will be called the non trivial vertices. The unlabeled trees are partially ordered from the root to the endpoints in the natural way; we shall use the symbol $<$ to denote the partial order. Two unlabeled trees are identified if they can be superposed by a suitable continuous deformation, so that the endpoints with the same index coincide. It is then easy to see that the number of unlabeled trees with n end-points is bounded by 4^n . We shall consider also the labeled trees (to be called simply trees in the following); they are defined by associating some labels with the unlabeled trees, as explained in the following items.

2) We associate a label $h \geq 0$ with the root and we denote $T_{h,n}$ the corresponding set of labeled trees with n endpoints. Moreover, we introduce a family of vertical lines, labeled by an integer taking values in $[h; 2]$, and we represent any tree

$T_{h,n}$ so that, if v is an endpoint or a non trivial vertex, it is contained in a vertical line with index $h_v > h$, to be called the scale of v , while the root is on the line with index h . There is the constraint that, if v is an endpoint, $h_v > h + 1$; if there is only one endpoint its scale must be equal to $h + 2$, for $h \geq 0$.

The tree will intersect in general the vertical lines in set of points different from the root, the endpoints and the non trivial vertices; these points will be called trivial vertices. The set of the vertices of T will be the union of the endpoints, the trivial vertices and the non trivial vertices. Note that, if v_1 and v_2 are two vertices and $v_1 < v_2$, then $h_{v_1} < h_{v_2}$.

Moreover, there is only one vertex immediately following the root, which will be denoted v_0 and can not be an endpoint; its scale is $h + 1$.

3) With each endpoint v of scale $h_v = +2$ we associate one of the contributions to $V^{(1)}$ given by (3.9); with each endpoint v of scale $h_v = 1$ one of the terms in

$LV^{(h_v-1)}$ defined in (4.41). Moreover, we impose the constraint that, if v is an endpoint and $h_v = 1$, $h_v = h_{v^0} + 1$, if v^0 is the non trivial vertex immediately preceding v .

4) If v is not an endpoint, the cluster L_v with frequency h_v is the set of endpoints following the vertex v ; if v is an endpoint, it is itself a (trivial) cluster. The tree provides an organization of endpoints into a hierarchy of clusters.

5) We introduce a field label f to distinguish the field variables appearing in the terms associated with the endpoints as in item 3); the set of field labels associated with the endpoint v will be called I_v . Analogously, if v is not an endpoint, we shall call I_v the set of field labels associated with the endpoints following the vertex v ; $x(f)$, $"(f)$ and $!(f)$ will denote the space-time point, the $"$ index and the $!$ index, respectively, of the field variable with label f .

If $h = 0$, the effective potential can be written in the following way:

$$V^{(h)}\left(\overline{Z_h}^{(h)}\right) + M^2 E_{h+1} = \sum_{n=1}^X \sum_{2T_{0,n}} V^{(h)}\left(\overline{Z_h}^{(h)}\right); \quad (4.52)$$

where, if v_0 is the first vertex of \mathcal{T} and \mathcal{T}_s ($s = s_{v_0}$) are the subtrees of \mathcal{T} with root v_0 , $\overline{Z_h}^{(h)}$ is defined inductively by the relation

$$V^{(h)}\left(\overline{Z_h}^{(h)}\right) = \frac{(1)^{s+1}}{s!} E_{h+1}^T [V^{(h+1)}(\mathcal{T}_1; \overline{Z_h}^{(h+1)}); \dots; V^{(h+1)}(\mathcal{T}_s; \overline{Z_h}^{(h+1)})]; \quad (4.53)$$

and $V^{(h+1)}(\mathcal{T}_i; \overline{Z_h}^{(h+1)})$

- a) is equal to $R V^{(h+1)}(\mathcal{T}_i; \overline{Z_h}^{(h+1)})$ if the subtree \mathcal{T}_i is not trivial;
- b) if \mathcal{T}_i is trivial and $h = 1$, it is equal to one of the terms in $LV^{(h+1)}(\overline{Z_h}^{(h+1)})$ or, if $h = 0$, to one of the terms contributing to $V^{(1)}(1)$ (3.9).

In (4.53) E_{h+1}^T denotes the truncated expectation with respect to the measure $Q_{\mathcal{T}}(d\mathcal{T}^{(h+1)})$. The r.h.s. of (4.53) can be written more explicitly in the following way. Given $2T_{0,n}$, there are n endpoints of scale 2 and only another one vertex, v_0 , of scale 1; let us call $v_1; \dots; v_n$ the endpoints. We choose, in any set I_{v_i} , a subset Q_{v_i} and we define $P_{v_0} = [Q_{v_i}]$. We have

$$V^{(0)}(\mathcal{T}; (0)) = \sum_{P_{v_0}} V^{(0)}(\mathcal{T}; P_{v_0}); \quad (4.54)$$

$$V^{(0)}(\mathcal{T}; P_{v_0}) = \int dx_{v_0} \sim^0(P_{v_0}) K_{P_{v_0}}^{(1)}(x_{v_0}); \quad (4.55)$$

$$K_{P_{v_0}}^{(1)}(x_{v_0}) = \frac{1}{n!} E_1^T [\sim^{(1)}(P_{v_1} n Q_{v_1}); \dots; \sim^{(1)}(P_{v_n} n Q_{v_n})] \prod_{i=1}^n K_{v_i}^{(2)}(x_{v_i}); \quad (4.56)$$

where we use the definitions $\sim^{(h)}(P_v) = \bigotimes_{f \in P_v} \mathcal{Q}^{m(f)}_{x(f);!(f)} \sim^{(h)}(f)$, and $K_{v_1}^{(2)}(x_{v_1})$ are the kernels in (3.9). We now write $V^{(0)}$ as $LV^{(0)} + RV^{(0)}$, with $LV^{(0)}$ defined as in (4.21), and we write for $RV^{(0)}$ a decomposition similar to the previous one, with $RV^{(0)}(\cdot; P_{v_0})$ in place of $V^{(0)}(\cdot; P_{v_0})$. By iterating the previous procedure, one gets for $V^{(h)}(\cdot; \sim^{(h)})$, for any $h \geq 2$, the representation described below (see [GM] and [BM] for details).

We associate with any vertex v of the tree a subset P_v of I_v , the external fields of v . These subsets must satisfy various constraints. First of all, if v is not an endpoint and $v_1; \dots; v_{s_v}$ are the vertices immediately following it, then $P_v \cap P_{v_i} = \emptyset$; if v is an endpoint, $P_v = I_v$. We shall denote Q_{v_i} the intersection of P_v and P_{v_i} ; this definition implies that $P_v = \bigcup_i Q_{v_i}$. The subsets $P_{v_i} \cap Q_{v_i}$, whose union I_v will be made, by definition, of the internal fields of v , have to be non empty, if $s_v > 1$. Given $h \geq 2$, there are many possible choices of the subsets P_v , $v \in \mathcal{T}_h$, compatible with all the constraints; we shall denote \mathcal{P} the family of all these choices and \mathcal{P} the elements of \mathcal{P} . We can write

$$V^{(h)}(\cdot; \sim^{(h)}) = \sum_{P \in \mathcal{P}} V^{(h)}(\cdot; P); \quad (4.57)$$

$V^{(h)}(\cdot; P)$ can be represented as

$$V^{(h)}(\cdot; P) = \int_{\mathcal{P}_{v_0}} \frac{d\mathbf{x}_{v_0}}{Z_h} \sim^{(h)}(P_{v_0}) K_{\mathcal{P}}^{(h+1)}(x_{v_0}); \quad (4.58)$$

with $K_{\mathcal{P}}^{(h+1)}(x_{v_0})$ defined inductively (recall that $h_{v_0} = h + 1$) by the equation, valid for any $v \in \mathcal{T}_h$ which is not an endpoint,

$$K_{\mathcal{P}}^{(h_v)}(x_v) = \frac{1}{s_v!} \left(\frac{Z_{h_v}}{Z_{h_v-1}} \right)^{\frac{\mathcal{P}_{v_j}}{2}} \prod_{i=1}^{s_v} [\mathcal{K}_{v_i}^{(h_v+1)}(x_{v_i})] E_{h_v}^T[\sim^{(h_v)}(P_{v_1} \cap Q_{v_1}); \dots; \sim^{(h_v)}(P_{v_{s_v}} \cap Q_{v_{s_v}})]; \quad (4.59)$$

Moreover, if v is an endpoint, $K_v^{(2)}(x_v)$ is defined as in (3.9) if $h_v = 2$, otherwise

$$K_v^{(h_v)}(x_v) = \begin{cases} \int_{\mathcal{P}_{v_1}} \int_{\mathcal{P}_{v_2}} \dots \int_{\mathcal{P}_{v_8}} \mathcal{K}_{v_1}^{(h_v+1)}(x_{v_1}) \mathcal{K}_{v_2}^{(h_v+1)}(x_{v_2}) \dots \mathcal{K}_{v_8}^{(h_v+1)}(x_{v_8}) & \text{if } v \text{ is of type } \text{---}, \\ \int_{\mathcal{P}_{v_1}} \int_{\mathcal{P}_{v_2}} \dots \int_{\mathcal{P}_{v_{h_v-1}}} \mathcal{K}_{v_1}^{(h_v+1)}(x_{v_1}) \mathcal{K}_{v_2}^{(h_v+1)}(x_{v_2}) \dots \mathcal{K}_{v_{h_v-1}}^{(h_v+1)}(x_{v_{h_v-1}}) & \text{if } v \text{ is of type } \text{---}, \\ \int_{\mathcal{P}_{v_1}} \int_{\mathcal{P}_{v_2}} \dots \int_{\mathcal{P}_{v_{h_v-1}}} \mathcal{K}_{v_1}^{(h_v+1)}(x_{v_1}) \mathcal{K}_{v_2}^{(h_v+1)}(x_{v_2}) \dots \mathcal{K}_{v_{h_v-1}}^{(h_v+1)}(x_{v_{h_v-1}}) & \text{if } v \text{ is of type } \text{---}, \\ \int_{\mathcal{P}_{v_1}} \int_{\mathcal{P}_{v_2}} \dots \int_{\mathcal{P}_{v_{h_v-1}}} \mathcal{K}_{v_1}^{(h_v+1)}(x_{v_1}) \mathcal{K}_{v_2}^{(h_v+1)}(x_{v_2}) \dots \mathcal{K}_{v_{h_v-1}}^{(h_v+1)}(x_{v_{h_v-1}}) & \text{if } v \text{ is of type } \text{---}, \end{cases} \quad (4.60)$$

If v_i is not an endpoint,

$$K_{v_i}^{(h_v+1)}(x_{v_i}) = R K_{\mathcal{P}_i}^{(h_v+1)}(x_{v_i}); \quad (4.61)$$

where \mathcal{P}_i is the subtree of \mathcal{T}_h starting from v and passing through v_i (hence with root the vertex immediately preceding v), \mathcal{P}_i are the restrictions to \mathcal{P}_i of \mathcal{P} .

(4.57) is not the normal form of our expansion, since we further decompose $V^{(h)}(\cdot; P)$, by using the representation of the truncated expectation (3.21). If we apply the expansion (3.21) in each non trivial vertex of Γ , we get an expression of the form

$$V^{(h)}(\cdot; P) = \sum_{T \in \mathcal{T}} \frac{P}{Z_h} \mathcal{P}_{v_0} \int^X \int^Z dx_{v_0} \sim^{(h)} (P_{v_0}) W_{\cdot; P; T}^{(h)}(x_{v_0}); \quad (4.62)$$

where \mathcal{T} is a special family of graphs on the set of points x_{v_0} , obtained by putting together an anchored tree graph T_v for each non trivial vertex v . Note that any graph $T \in \mathcal{T}$ becomes a tree graph on x_{v_0} , if one identifies all the points in the sets x_v , for any vertex v which is also an endpoint.

Taking into account the effect of the R operation we obtain (see [BM], x3 for a detailed proof of a similar formula)

$$W_{\cdot; P; T}(x_{v_0}) = \sum_{A_T} \prod_{i=1}^n \int^{Y_i} \int^{Z_i} \frac{1}{s_v!} dP_{T_v}(t_v) \det G^{h_v; T_v}(t_v) \prod_{v \text{ not ep.}} \left[\int^{Y_v} \left(\frac{Z_{h_v}}{Z_{h_v-1}} \right)^{\frac{P_{v,j}}{2}} \int^{h_{Y_v}} \int^{q(f_1)} \int^{q(f_1^+)} [(x_1 \dots y_1)^{b(l)} \int^{m_1} \int^{(h_v)} g_{i_1; i_1^+}^{(h_v)}(x_1 \dots y_1)]^{i_1} \right]; \quad (4.63)$$

where "\ep." is an abbreviation of "endpoint" and, together with the definitions used before, we are using the following ones:

- 1) A_T is a set of indices which allows to distinguish the different terms produced by the non trivial R operations and the iterative decomposition of the zeros;
- 2) v_1, \dots, v_n are the endpoints of Γ and $h_i = h_{v_i}$;
- 3) $b(l)$, $q(f_1)$ and $q(f_1^+)$ are positive integers ≥ 3 ; $\hat{e}^0 = I$;
- 4) if v is a non trivial vertex (so that $s_v > 1$), the elements $G_{i,j;i^0,j^0}^{h_v; T_v}$ of $G^{h_v; T_v}(t_v)$ are of the form

$$G_{i,j;i^0,j^0}^{h_v; T_v} = t_{i,i^0} \hat{e}^{q(f_{ij})} \hat{e}^{q(f_{i^0,j^0}^+)} \int^{m(f_1)} \int^{m(f_1^+)} g_{i_1; i_1^+}^{(h_v)}(x_{ij} \dots y_{i^0,j^0}); \quad (4.64)$$

if v is trivial, T_v is empty and $\int^R dP_{T_v}(t_v) \det G^{h_v; T_v}(t_v)$ has to be interpreted as 1, if $j_{I_v} = 0$ (I_v is the set of internal eds of v), otherwise it is the determinant of a matrix of the form (4.64) with $t_{i,i^0} = 1$.

It would be very difficult to give a precise description of the various contributions to the sum over A_T , but fortunately we only need to know some very general properties, in particular that $\# A_T \leq C^n$ for some constant C and that for any $\sum_{T \in \mathcal{T}} \dots$, the following inequality is satisfied

$$\sum_{f \in I_{v_0}} \int^{h_{Y_v}} \int^{h(f)q(f)} \int^{h(l)b(l)} \int^{i_1} \int^{Y_v} \dots \int^{z(P_v)}; \quad (4.65)$$

$\sum_{f \in I_{v_0}} \quad \sum_{T \in \mathcal{T}} \quad \sum_{v \text{ not ep.}}$

where $h(f) = h_{v_0} - 1$ if $f \in P_{v_0}$, otherwise it is the scale of the vertex where the edge with label f is contracted; $h(l) = h_v$, if $l \in T_v$ and

$$z(P_v) = \begin{cases} 1 & \text{if } |P_v| = 4, \\ 2 & \text{if } |P_v| = 2, \\ 0 & \text{otherwise.} \end{cases} \quad (4.66)$$

By a standard computation (see for instance [BM], x3) by bounding the determinant by the Gram-Hadamard inequality (see (3.23)) we obtain

$$\begin{aligned} & \sum_{v \text{ not ep.}} \int_{\mathbb{R}^{d_{v_0}}} \int_{\mathbb{R}^{d_T}} (x_{v_0}) \int_{\mathbb{R}^{d_n}} C^n M^{2n} h_{D_k(P_{v_0})} \\ & \leq \sum_{v \text{ not ep.}} \frac{1}{s_v!} C^{s_v} \prod_{i=1}^{s_v} |P_{v_i}| |P_v| \left(\frac{Z_{h_v}}{Z_{h_v-1}} \right)^{\frac{|P_v|}{2}} [2 + \frac{|P_v|}{2} + z(P_v)]^{\frac{|P_v|}{2}}; \end{aligned} \quad (4.67)$$

with $2 + \frac{|P_v|}{2} + z(P_v) > 0$. In order to perform the sums note that the number of unlabeled trees is 4^n ; fixed an unlabeled tree, the number of terms in the sum over the various labels of the tree is bounded by C^n , except the sums over the scale labels. In order to bound the sums over the scale labels and P we first use the inequality

$$\begin{aligned} & \sum_{v \text{ not ep.}} [2 + \frac{|P_v|}{2} + z(P_v)]^{\frac{|P_v|}{2}} \\ & \leq \sum_{v \text{ not ep.}} \prod_{\mathfrak{v}} [2 + (h_{\mathfrak{v}} - h_{\mathfrak{v}^0})] [2 + |P_v|]; \end{aligned} \quad (4.68)$$

where \mathfrak{v} are the non trivial vertices, and \mathfrak{v}^0 is the non trivial vertex immediately preceding \mathfrak{v} or the root. The factors $2 + (h_{\mathfrak{v}} - h_{\mathfrak{v}^0})$ in the r.h.s. of (4.68) allow to bound the sums over the scale labels by C^n ; $\frac{1}{40}$ is a suitable constant (one finds $\frac{1}{40}$).

Finally the sum over P can be bounded by using the following combinatorial inequality, trivial for n large enough. Let $f_{p_v}; v \in \mathbb{N}$ a set of integers such that $p_v = \sum_{i=1}^{s_v} p_{v_i}$ for all $v \in \mathbb{N}$ which are not endpoints; then

$$\sum_{v \text{ not ep.}} \prod_{p_v} C^{\frac{p_v}{40}} \leq C^n; \quad (4.69)$$

It follows that

$$\begin{aligned} & \sum_{v \text{ not ep.}} \prod_{p_v} C^{\frac{p_v}{40}} \leq C^n; \\ & \sum_{v \text{ not ep.}} \prod_{p_v} C^{\frac{p_v}{40}} \leq C^n; \end{aligned} \quad (4.70)$$

4.9 Remark 1. The decomposition in (4.23) respects the determinant structure of the truncated expectations, as we can decompose the propagators as in

(4.34), (4.38), obtaining, for any tree with n end-points, a family of C^n different contributions to $\hat{W}_2^{(h)}$, which can be bounded as before. Hence we can include in $\hat{W}^{(a)}$ the terms with at least a propagator $\hat{g}_1^{(k)}$, defined in (4.39).

Remark 2. If the tree has an end-point on scale k , the bound (4.50) can be improved by a factor h^{-k} , as it follows immediately from (4.68). In particular if to an end-point is associated a term $R V^{(1)}$ there is an extra factor h ; this is a short memory property.

4.10 The above results were proved for $Z_{2I}^{i, i, i}$; a similar analysis can be repeated for $Z_{2I}^{n^{(1)}, n^{0(1)}, n^{(2)}, n^{0(2)}}$, for any value of $n^{(1)}, n^{0(1)}, n^{(2)}, n^{0(2)}$. The only difference is that one has in addition the function Q in the interaction and the oscillating functions $e^{ip_M x}$. One can split V^1 in a part identical to the one for $Z_{2I}^{i, i, i}$, called $V^{(1)}$, and the rest; hence we repeat the multiscale analysis by writing $V^{(h)} = V^{(h)} + V^{0(h)}$, with $V^{0(h)}$ given by a sum of trees with at least an end-point associated to $V^{(1)}$; we define the localization operators acting non trivially only on $V^{(h)}$ (and defined as above). It is easy to see that the terms from trees with at least one end-point associated to $V^{(1)}$ are vanishing in the limit $M \rightarrow 1$; in fact the bounds for such terms is improved by the factor $\frac{h}{M}$, by simply dimensional considerations, and we will see in the following section that h is a finite number independent from M , hence such terms are vanishing in the limit $M \rightarrow 1$.

5. The flow of the running coupling constants

5.1 By the analysis of the preceding section it follows that the running coupling constants $v_k; k; m_k, 1 \leq k \leq h$, verify a set of recursive equations called Beta function equations, of the form

$$\begin{aligned} v_{h-1} &= v_h + h (v_h; v_h; \dots; v_1; 1) \\ v_{h-1} &= v_h + \frac{h}{v} (v_h; v_h; \dots; v_1; 1) \\ \frac{m_{h-1}}{m_h} &= 1 + \frac{h}{m} (v_h; v_h; \dots; v_1; 1) \\ \frac{Z_{h-1}}{Z_h} &= 1 + \frac{h}{z} (v_h; v_h; \dots; v_1; 1) \end{aligned} \tag{5.1}$$

By repeating the analysis for proving Theorem 3 to the functions $\frac{h}{i}$ we have that, if (4.49) holds, then the $\frac{h}{i}$ are expressed by convergent series. We want to show that there exists a positive constant a_3 such that, if $\|t_c\| \leq \frac{1}{a_3^2}$, it is possible to find a function ϕ_1 (hence a function ϕ) so that (4.49) holds (if ϕ is small enough).

Iterating the first of (5.1) we find

$$\varphi_h = \varphi_{h+1} + \sum_{k=h+1}^{\infty} X^1_{k-2-k} (\varphi_k; \dots; \varphi_0) ; \quad (5.2)$$

where now the functions φ_k are thought as functions of $\varphi_k; \dots; \varphi_1$ only.

If we put $h = h$ in (5.2), we get the following identity:

$$\varphi_h = \varphi_{h+1} + \sum_{k=h+1}^{\infty} X^1_{k-2-k} (\varphi_k; \dots; \varphi_0) ; \quad (5.3)$$

and we look for a φ_1 verifying

$$\varphi_1 = \sum_{k=h+1}^{\infty} X^1_{k-2-k} (\varphi_k; \dots; \varphi_1) ; \quad (5.4)$$

so that

$$\varphi_h = \varphi_h + \sum_{k=h+1}^{\infty} X^h_{k-1-k} (\varphi_k; \dots; \varphi_1) ; \quad h < h-1 : \quad (5.5)$$

Let be $\sim = f_{\gamma_k; h} \quad h-1$ and $\|\varphi\| = \sup_{h \leq h-1} \|\varphi_h\|$; we call M the set of \sim with bounded norm. We decouple the beta function equations (5.1) imagining that in the last three equations of (5.1) φ_k is replaced by γ_k acting as a parameter; we call $m_h(\sim); v_k(\sim)$ the solution of the second and third of (5.1) as functions of the parameter \sim . We shall prove the following lemma.

5.2 Lemma. There exists ϵ and a_3 such that, for $\|\varphi\| \leq e^{\frac{1}{a_3-1}}$, $\|\varphi\| \leq \epsilon$ and any \sim such that $\|\varphi\|, \|\varphi^0\| \leq C \|\varphi\|$, it holds, if $m_0 = \|\varphi\|$ and C, C are constants

$$\|\varphi_h(\sim)\| \leq \|\varphi\| + \frac{\|\varphi\|}{4} \quad \|\varphi_h(\sim)\| \leq \|\varphi_h(\sim^0)\| + C \|\varphi\| \max_{k>h} \|\varphi_k\| \quad \|\varphi_k^0\| \quad (5.6)$$

$$\|\varphi_h(\sim)\| \leq \|\varphi\| + \frac{\|\varphi\|}{4} \quad \|\varphi_h(\sim)\| \leq \|\varphi_h(\sim^0)\| + C \|\varphi\| \max_{k>h} \|\varphi_k\| \quad \|\varphi_k^0\|$$

$$\|\varphi_{i;h}(\sim)\| \leq \|\varphi_{0;h}\| + \|\varphi\| \quad \|\varphi_{i;h}(\sim)\| \leq \|\varphi_{i;h}(\sim^0)\| + C \|\varphi\| \max_{k>h} \|\varphi_k\| \quad \|\varphi_k^0\|$$

$$\|\varphi_{i;h}(\sim)\| \leq \|\varphi_{0;h}\| + \|\varphi\| \quad \|\varphi_{i;h}(\sim)\| \leq \|\varphi_{i;h}(\sim^0)\| + C \|\varphi\| \max_{k>h} \|\varphi_k\| \quad \|\varphi_k^0\|$$

$$\|\varphi_0\|^{c_1 a_2 - 1} \|\varphi_h(\sim)\| \leq \|\varphi_0\|^{c_2 a_2 - 1} \|\varphi\| \frac{\|\varphi_h(\sim)\|}{\|\varphi_h(\sim^0)\|} \leq \|\varphi\| \max_{k>h} \|\varphi_k\| \quad \|\varphi_k^0\| \|\varphi\|^{-1}$$

$$c_3 a_4 \|\varphi\|^{-2} \|\varphi_h(\sim)\| \leq c_4 a_4 \|\varphi\|^{-2} \|\varphi\| \frac{\|\varphi_{h-1}(\sim)\|}{\|\varphi_{h-1}(\sim^0)\|} \leq \|\varphi\| \max_{j>h} \|\varphi_j\| \quad \|\varphi_k^0\|$$

with $a_2; a_4; c_1; c_2; c_3; c_4$ positive constants.

Remark. An obvious corollary of the above statement is that there exists a finite (uniformly in M) h such that $m_{h-1} \leq h^{-1}$ for a suitable constant ϵ .

5.3 Proof. The proof is done by induction. By iterating the second equation of (5.1) we find

$$j_{h-1}(\sim) = \sum_{k=h}^{X^1} j_k(v_h(\sim); \gamma_h; \dots; v_1(\sim); \gamma_1) \quad (5.7)$$

and by induction

$$j_k(v_h(\sim); \gamma_h; \dots; v_1(\sim); \gamma_1) \leq C_1 [j_1]^3 + \frac{2}{1} \frac{j_{h-1}}{h} + \frac{2}{1} h^{-1} \quad (5.8)$$

The first addend of the r.h.s. of (5.8) is a bound on the terms containing only propagators $g_{L;1}^{(h)}(x, y)$, see (4.35), and trees with end-points v with $h_v = 1$, and the bound follows from the fact that the second order terms cancel out. The second addend is a bound on the terms containing at least a propagator $g_{1;1;2}^{(h)}(x, y)$ (4.39), (4.37) (the bound follows from the short memory property and (4.33), (4.37)). The third addend is a bound on the sum of terms with at least a propagator $g_{1;1;2}^{(h)}(x, y)$ or from trees with at least an end-point v with $h_v = 2$, and we have used (4.36).

Inserting (5.8) in (5.7) we get

$$j_{h-1}(\sim) \leq C_1 [j_1]^3 + \sum_{k=h}^{X^0} \left[\frac{j_k}{k} + j_k \right] j_1 \frac{j_1}{4} \quad (5.9)$$

for a_3 large enough and ϵ small enough, where we have used that $\sum_{k=h}^P \frac{j_k}{k}$ is bounded (by induction and the definition of h).

Moreover the expansion for $j_{h-1}(\sim) = j_{h-1}(\sim^0)$ is given by a sum of terms similar to the ones for $j_{h-1}(\sim)$ in which a v_k is replaced by $v_k(\sim) = v_k(\sim^0)$ bounded by $C^{-2} \max_{j>k} j_j \sim_j^0 j_{\text{orm } k}$ or Z_k replaced by the relative difference; one finds, for a_3 large and ϵ small enough

$$j_{h-1}(\sim) = j_{h-1}(\sim^0) \leq C_2 [j_1]^3 + \sum_{k=h}^{X^0} \left[\frac{j_k}{k} + j_k \right] \max_{k>h} j_k \sim_k^0 j_1 \leq C j_1 \max_{k>h} j_k \sim_k^0 j_1 \quad (5.10)$$

where the first term is a bound on the terms containing only propagators $g_{L;1}^{(h)}(x, y)$, and trees with end-points v with $h_v = 1$, containing at least a $v_k(\sim) = v_k(\sim^0)$ or a $\gamma_k \sim_k^0$ (note also that the first term with end-point \sim is $\frac{2}{h} \frac{2}{h}$). A similar analysis can be repeated for h .

The recursive equation for $\tilde{m}_{i;!,h}$ is

$$\tilde{m}_{i;!,h-1}(\sim) = \tilde{m}_{i;!,h}(\sim) + \tilde{m}_{i;!,h} \left(\frac{2}{h} a_z + z \right) + R_z^{(h)} \quad (5.11)$$

with $j_z^h j = C_3 j^3$ and $\mathcal{R}_z^h j = C_4 \left[h + \frac{j_m h j}{h} \right]^2$. In $\frac{2}{h} (a_2 + z)$ in (5.11) are the contributions from terms with only diagonal propagators $g_{L;!,}^{(h)}(x)$ and from trees with endpoints v with $h_v = 1$; there is necessarily at least a vertex \sim . In fact assume that a contribution has n end-points of type \sim and n end-points of type $!$; there are then $n \frac{1}{2} + \frac{n}{2}$ propagators $! = 1$ and $n \frac{1}{2} + \frac{n}{2}$ propagators $! = 1$, so that n must be odd; the total number of propagators $2n = 1 + n$ is an even number; the derivated integrand is odd under the exchange $k \leftrightarrow k$ and so it is vanishing. Moreover there is no contribution of order $h z_h$ (they corresponds to tadpole graphs, whose derivative is zero). In $R_z^{(h)}$ are the trees with at least an end point v with $h_v = 2$, or with a propagator $\hat{g}_{!_1;!_2}^{(k)}$, or with a propagator $\hat{g}_{!_1;!_2}^{(k)}$. It holds by iterating, for a_3 large and \sim small enough, $\tilde{m}_{i;!,h} j = \mathcal{R}_{i;!,0} j + j$. Moreover

$$\begin{aligned} \tilde{m}_{i;!,h-1}(\sim) - \tilde{m}_{i;!,h-1}(\sim^0) j &= C_3 \max_{j>h} j_k - \sim_k^0 j \left[j^3 h j + j j \sum_{k=h}^{X^0} \left[\frac{j_m k j}{k} + k \right] \right] \\ &= C j \max_{j>h} j_k - \sim_k^0 j \end{aligned} \quad (5.12)$$

Similar computations can be repeated for $\tilde{m}_{i;!,h}$.

By definition (see (4.23)) $m_{h-1}(\sim)$ is given by a sum over terms with at least a non diagonal propagator $\hat{g}_{!;!,}^{(h)}(x-y)$, hence

$$\frac{m_{h-1}(\sim)}{m_h(\sim)} = 1 + h (a_2 + h_m) \quad (5.13)$$

where $a_2 > 0$ is a constant and $j_m j = C j_1 j$; then (assuming $j_1 > 0$; similar computations can be repeated for $j_1 < 0$), for \sim small enough

$$\left(1 - \frac{5}{4} a_2^{-1} \right) \frac{j_{h-1}(\sim) j}{j_h(\sim) j} \left(1 - \frac{3}{4} a_2^{-1} \right) \quad (5.14)$$

Then

$$j_{h-1}(\sim) j = j_0 j^{2a_2-1(h-1)} \frac{\left(1 - \frac{5}{4} a_2^{-1} \right)}{2a_2-1} j_0 j^{2a_2-1(h-1)} \quad (5.15)$$

and

$$j_{h-1}(\sim) j = j_0 j^{\frac{a_2}{2}-1(h-1)} \frac{\left(1 - \frac{3}{4} a_2^{-1} \right)}{1-\frac{a_2}{2}} j_0 j^{\frac{a_2}{2}-1(h-1)} \quad (5.16)$$

$$C_4 \log \max_{0 \leq k \leq h} j_k(\mathbf{v}) - v_k(\mathbf{v}) j_{\max_{j \geq h} j_k} \sim_k^0 j j^{-1} \quad (5.17)$$

5.4 Lemma. There exists ϵ, a_3 and C such that, for $j \leq t_c j \leq \frac{1}{e^{a_3 - 2}}$ and $j \leq \epsilon$, there exists a $\delta_1(\epsilon)$ so that $\max_{k \in [h, j_k] j} C \leq j$.

$$\tilde{z}_h^0 = \sum_{k=h+1}^{k-1} X^h_k (v_k(\sim); \gamma_k; \dots; v_0(\sim); \gamma_0); \quad h < h-1: \quad (5.18)$$

By (5.18) and Theorem 3

where C_1 is a constant depending on the first order contribution (i.e. the tadpole). So, for a proper C , $\|\tilde{j}^0\| \leq C \|j\|$.

$$\tilde{\sim}_{1,h}^0 \quad \tilde{\sim}_{2,h}^0 = \sum_{k=h+1}^h X^h \quad k=1$$

and

$$\tilde{J}_{1,h}^0 \sim_{2,h}^0 J \quad C_1 J \quad J \quad \begin{matrix} X^h \\ k=h+1 \end{matrix} \quad \begin{matrix} X \\ j=h \end{matrix} \quad (h-j) \max_{i,j} \tilde{J}_{1,i} \sim_{2,i} \tilde{J} \quad C_2 J \quad \tilde{J}_{1,i} \sim_{2,i} \tilde{J}$$

with C_1, C_2 constants, so that T is a contraction.

5.5 If $a = b = 0$ one has the case treated in [SP], corresponding to two independent Ising models with a quartic or a next to nearest neighbor interaction. In such a case the local part of the quartic terms is vanishing by Pauli principle, so that if $\mathcal{P}_v = 4$ one can apply "freely" a first order renormalization obtaining an additional $(h_v - h_{v0})$, for any v such that $\mathcal{P}_v = 4$, in the bounds. At each step one can include all the quadratic running coupling constants in the free integrations; their beta function is $\tilde{g}_{i,h+1} = \tilde{g}_{i,h} + O(h^{-1})$ and $g_{i,h+1} = g_{i,h} + O(h^{-1})$, as the beta function is sum over all the trees with end-points necessarily at scale $h_v = 2$ (contrary to our case, as the local part of the quartic terms is vanishing, and there are no end-points associated to the quadratic terms as they are included in the free integration). Hence all the quadratic couplings are $O(h^{-1})$ and they do not change the scaling properties of the propagator on a single scale; the qualitative behaviour of C_v for $v = 0$ or $v \neq 0$ are then the same for temperatures up to t_c .

6. Correlation functions

6.1 Flow of observables. We consider the following functional integral

$$e^{S(\varphi)} = \int P(d\varphi) e^{V^{(1)}(\varphi) + B(\varphi; \varphi)} \quad (6.1)$$

where $V^{(1)}$ given by (3.8) and $B = B^1 + B^2$ with

$$B^{(1)}(\varphi; \varphi) = \int dx^{(1)}(x) \left[\varphi_{x;1}^+ \varphi_{x;1} + \varphi_{x;1}^+ \varphi_{x;1}^+ \right] \quad (6.2)$$

$$B^{(2)}(\varphi; \varphi) = \int dx^{(2)}(x) \left[\varphi_{x;1}^+ \varphi_{x;1}^+ + \varphi_{x;1} \varphi_{x;1} \right]; \quad (6.3)$$

After integrating the fields $\varphi^{(1)}; \dots; \varphi^{(h+1)}, 0 \leq h \leq h$, we find

$$e^{S(\varphi)} = e^{L - E_h + S^{(h+1)}(\varphi)} \int_{\mathcal{P}_{Z_h} \mathcal{M}_h; \mathcal{C}_h} (d\varphi^{(h)}) e^{V^{(h)}(\varphi^{(h)}) + B^{(h)}(\varphi^{(h)}; \varphi^{(h)})}; \quad (6.4)$$

where $\mathcal{P}_{Z_h} \mathcal{M}_h; \mathcal{C}_h}(d\varphi^{(h)})$ and $V^{(h)}$ are given by (4.8) and (4.11), respectively, while $S^{(h+1)}(\varphi)$, which denotes the sum over all the terms dependent on φ but independent of the field, and $B^{(h)}(\varphi^{(h)}; \varphi^{(h)})$, which denotes the sum over all the terms containing at least one field and two fields, can be represented in the form

$$S^{(h+1)}(\varphi) = \sum_{m=1}^{\infty} \int dx_1 \dots dx_m \varphi_m^{(h+1)}(x_1; \dots; x_m) \sum_{i=1}^h \varphi_i^{(i)}(x_i) \quad (6.5)$$

with $j_i \in \mathbb{C}^2, i = 1, 2$; proceeding as in the proof of Lemma 5.2

$$c_4 h < \frac{Z_h^{(1)}}{Z_h} < c_3 h \quad c_1 h < \frac{Z_h^{(2)}}{Z_h} < c_2 h : \quad (6.14)$$

The fields of scale between h and h_L , are integrated in a single step, and it follows that

$$S(\cdot) = L E_L; + S^{(h)}(\cdot) = L E_L; + \sum_{h=h}^{X^1} S^{(h)}(\cdot); \quad (6.15)$$

As a fields is dimensionally analogous to two external fields, we get, proceeding as in the proof of Theorem 3 (see also [BM] for a detailed proof of similar bounds in a related case)

$$j \frac{\partial^2}{\partial^{(1)}(x) \partial^{(1)}(y)} S^{(h)}(\cdot) j = 0 \quad 2h \left[\frac{Z_h^{(1)}}{Z_h} \right]^2 \frac{C_N}{1 + (h j(x-y) j)^N} \quad (6.16)$$

$$j \frac{\partial^2}{\partial^{(2)}(x) \partial^{(2)}(y)} S^{(h)}(\cdot) j = 0 \quad 2h \left[\frac{Z_h^{(2)}}{Z_h} \right]^2 \frac{C_N}{1 + (h j(x-y) j)^N} \quad (6.17)$$

and if $x = x; y$ and $\cdot = 1; 2$

$$j \frac{\partial^3 S^{(h)}(\cdot)}{\partial^{(1)}(x_1) \partial^{(2)}(x_2) \partial^{(3)}(x_3)} j = 0 \quad h^4 \left[\prod_{i=1}^3 \frac{Z_h^{(i)}}{Z_h} \right] \frac{C_N}{1 + (h j(x-y) j)^N} \quad (6.18)$$

$$j \frac{\partial^4 S^{(h)}(\cdot)}{\partial^{(1)}(x_1) \partial^{(2)}(x_2) \partial^{(3)}(x_3) \partial^{(4)}(x_4)} j = 0 \quad 2h^6 \left[\prod_{i=1}^4 \frac{Z_h^{(i)}}{Z_h} \right] \frac{C_N}{1 + (h j(x-y) j)^N} \quad (6.19)$$

6.2 Correlation functions. The correlation function $\langle \cdot_x^{()} \cdot_{x^0}^{()} \cdot_y^{()} \cdot_{y^0}^{()} \rangle_T$, where $x^0 = (x+1; x_0)$ or $(x; x_0+1)$ is given by

$$\langle \cdot_x^{()} \cdot_{x^0}^{()} \cdot_y^{()} \cdot_{y^0}^{()} \rangle_T = \frac{\partial}{\partial J_{x;x^0}} \frac{\partial}{\partial J_{y;y^0}} \log Z_{2I} (f J_{x;x^0} g) j_{J_{x;x^0} g = f J g} \quad (6.20)$$

If $O^{()}(x) = \cdot_{x;x_0}^{()} \cdot_{x+1;x_0}^{()}$ each derivative produces a factor

$$\text{sech}^2 J H_x H_{x+1;x_0}^{()} + \tanh J + \frac{\partial}{\partial J_{x;x+1;x_0}} V \quad (6.21)$$

with V given by (2.25). We define, using (2.30), (2.25)

$$\begin{aligned} & \langle O^{(\cdot)}(x) O^{(\cdot)}(y) \rangle_{;n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}} = \\ & \frac{\int_{\mathbb{R}^Q} \prod_{i=1}^2 P_{n^{(i)};n^{0(i)}}^{(\cdot)}(dH^{(\cdot)};dV^{(\cdot)}) e^V O^{(\cdot)}(x) O^{(\cdot)}(y)}{\int_{\mathbb{R}^Q} \prod_{i=1}^2 P_{n^{(i)};n^{0(i)}}^{(\cdot)}(dH^{(\cdot)};dV^{(\cdot)}) e^V} \end{aligned} \quad (6.22)$$

Moreover we call

$$\begin{aligned} & \langle O^{(\cdot)}(x) O^{(\cdot)}(y) \rangle_{;T;n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}} = \langle O^{(\cdot)}(x) O^{(\cdot)}(y) \rangle_{;n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}} \\ & \langle O^{(\cdot)}(x) \rangle_{;n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}} < O^{(\cdot)}(x) \rangle_{;n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}} \end{aligned} \quad (6.23)$$

so that

$$\begin{aligned} & \langle O^{(\cdot)}(x) O^{(\cdot)}(x) \rangle_{;T} = \sum_{n^{(1)};n^{0(1)}}^X (1)^{n^{(1)};n^{0(1)}} \sum_{n^{(2)};n^{0(2)}}^X (1)^{n^{(2)};n^{0(2)}} \\ & \frac{\sum_{n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}}^X \langle O^{(\cdot)}(x) O^{(\cdot)}(y) \rangle_{;T;n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}}}{\sum_{n^{(1)};n^{0(1)}}^X (1)^{n^{(1)};n^{0(1)}} \sum_{n^{(2)};n^{0(2)}}^X (1)^{n^{(2)};n^{0(2)}} \sum_{n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}}^X} \end{aligned} \quad (6.24)$$

Suppose that x and y are fixed to an M -independent value; then

$$\lim_{M \rightarrow 1} \langle O^{(\cdot)}(x) O^{(\cdot)}(y) \rangle_{;T;n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}} < O^{(\cdot)}(x) O^{(\cdot)}(y) \rangle_{;T; ; ; ;} = 0: \quad (6.25)$$

In fact, as explained in x4.10, the l.h.s. of (6.25) can be written as a sum of trees, and in each of them there is a factor $e^{ip^{(1)};n^{0(1)}} \cdot 1$ or an end-point associated to $Q^{n^{(1)};n^{0(1)};n^{(2)};n^{0(2)}}$. With respect to the previous bounds, simply dimensional analysis says there is now an extra factor $\frac{h}{M}$ in the bounds so it is vanishing in the limit $M \rightarrow 1$, if x, y and $t \rightarrow t_c$ are fixed to an M -independent value. We can then simply study $\langle O^{(\cdot)}(x) O^{(\cdot)}(y) \rangle_{;T; ; ; ;}$ which is given by the Grassmann integral (6.22) with $O^{(\cdot)}$ given by (6.21). By performing the change of variables (2.31) and (2.59), we get a sum of averages of monomials in the ϕ and ψ elds; we integrate the ψ -elds, as discussed in x3, and we obtain a sum of Grassmann integrals of monomials in the ϕ -elds; remembering that

$$\begin{aligned} & H_x H_x^{(\cdot)} H_y H_y^{(\cdot)} = \frac{1}{16} [\phi_{1;x} \phi_{1;x} + \phi_{1;y} \phi_{1;y} + \phi_{1;y} \phi_{1;y} + \phi_{1;x} \phi_{1;x} + \\ & \phi_{1;x} \phi_{1;x} + \phi_{1;y} \phi_{1;y} + \phi_{1;y} \phi_{1;y} + \phi_{1;x} \phi_{1;x} + \phi_{1;x} \phi_{1;y} + \phi_{1;y} \phi_{1;x} + \phi_{1;y} \phi_{1;y}] \end{aligned} \quad (6.26)$$

we find the first two of (1.13) and (1.15) by (6.1), (6.13), (6.15) and (6.16), (6.17). In $(\cdot)^c$ we include to contributions to the correlation function corresponding to

monomials with six or more fields, and from (6.18), (6.19) the last of (1.13) follows.

Finally the specific heat is obtained in a similar way, noting that we have to sum over x, y , and this produces an extra 2^h in the r.h.s. of (6.16), (6.17), (6.18), (6.19).

Acknowledgments. This paper was partly written in the stimulating atmosphere of the Institute for Advanced Studies, in Princeton. I thank Prof. Spencer for his invitation and for many clarifying discussions about his work [PS].

References

- [B] R. Baxter, Exactly solved models in statistical mechanics, Academic Press (1982)
- [BG] G. Benfatto, G. Gallavotti: Renormalization group. Physics notes 1, Princeton University Press (1995).
- [BGPS] G. Benfatto, G. Gallavotti, A. Procacci, B. Scoppola: Beta Functions and Schwinger Functions for a Many Fermions System in One Dimension. Comm. Math. Phys. 160, 93{171 (1994).
- [BM] G. Benfatto, V. Mastropietro. Renormalization group, hidden symmetries and approximate Ward identities in the XYZ model. Rev. Math. Phys. 13 (2001), no. 11, 1323{143; and to appear in Comm. Math. Phys.
- [GM] Gentile, G., Mastropietro, V. Renormalization group for one-dimensional fermions. A review on mathematical results. Phys. Rep. 352 (2001), no. 4-6, 273{43
- [K] P. W. Kasteleyn, Dimer Statistics and phase transitions, J. Math. Phys. 4, 287 (1963)
- [F] C. Fan, On critical properties of the Ashkin-Teller model, Phys. Rev. B, 6, 136-136 (1972)
- [H] C. Hurst, New approach to the Ising problem, J. Math. Phys. 7, 2, 305-310 (1966)
- [ID] C. Itzykson, J. Drouot, "Statistical field theory: 1," Cambridge Univ. Press, 1989.
- [Le] A. Lesniewski: Effective action for the Yukawa 2 quantum field Theory. Comm. Math. Phys. 108, 437-467 (1987).
- [Li] H. Lieb, Exact solution of the problem of entropy of two-dimensional ice, Phys. Rev. Lett., 18, 692-694, (1967)
- [LP] A. Luther, I. Peschel. Calculations of critical exponents in two dimension from quantum field theory in one dimension. Phys. Rev. B 12, 3908-3917 (1975)
- [MW] B. McCoy, T. Wu, "The two-dimensional Ising model," Harvard Univ. Press, 1973.
- [MPW] J. M. Montroll, R. Potts, J. Ward. Correlation and spontaneous magnetization of the two dimensional Ising model. J. Math. Phys. 4, 308 (1963)

- [N] M. P. M. den Nijs. Derivation of extended scaling relations between critical exponents in two dimensional models from the one dimensional Luttinger model, *Phys. Rev. B*, 23, 11 (1981) 6111-6125
- [PB] A. M. M. P. Ruiskens, A. C. Brown. Universality for the critical lines of the eight vertex, Ashkin-Teller and Gaussian models, *Phys. Rev. B*, 23, 3 (1981) 1459-1468
- [PS] H. P. Pinson, T. Spencer. Universality in 2D critical Ising model. To appear in *Comm. Math. Phys.*
- [S] S. Samuel, The use of anticommuting variable integrals in statistical mechanics", *J. Math. Phys.* 21 (1980) 2806
- [Spe] T. Spencer: A mathematical approach to universality in two dimensions. *Physica A* 279, 250-259 (2000).
- [SML] T. Schultz, D. Mattis, E. Lieb, Two-dimensional Ising model as a soluble problem of many Fermions, *Rev. Mod. Phys.* 36 (1964) 856.
- [W] F. W. Wu The Ising model with four spin interaction *Phys. Rev. B* 4, 2312-2314 (1971)




Fig.4: Closed polygon in the high temperature expansion of the Ising model partition function.




Fig.3. The Eight-vertex model in the rotated frame, with spins relabelled; the solid lines represent the quartic interaction.




Fig.2. The spin representation of the Eight-vertex model;
the solid points have x_0+x even and the void points,
odd; the solid lines represent the quartic interaction.




Fig.1: The spin representation of the Ashkin Teller model ;
the dotted lines represent the Ising interactions, the solid
lines the quartic interaction.




Fig.5. The second lattice and its polygon configurations.




Fig.6. Local identification of Ising configurations and configurations over the second lattice.

