

# Thermodynamics of the one-dimensional $s=1$ $XXZ$ Heisenberg model: analytical results

Onofre Rojas,<sup>1</sup> E.V. Corrêa Silva,<sup>2</sup> Winder A. Moura-Melo,<sup>3</sup> S.M. de Souza,<sup>4</sup> and M.T. Thomaz<sup>2,\*</sup>

<sup>1</sup>*Departamento de Física, Universidade Federal de São Carlos, CEP 13565-905, São Carlos - SP, Brazil*

<sup>2</sup>*Instituto de Física, Universidade Federal Fluminense,*

*Av. Gal. Milton Tavares de Souza s/n°, CEP 24210-340, Niterói - RJ, Brazil*

<sup>3</sup>*Faculdades Federais Integradas de Diamantina, Rua da Glória nº 187, CEP 39100-000, Diamantina-MG, Brazil*

<sup>4</sup>*Departamento de Ciências Exatas, Universidade Federal de Lavras,*

*Caixa Postal 37, CEP 37200-000, Lavras-MG, Brazil*

(Dated: January 10, 2022)

We apply the results derived by Rojas *et al.*[1] to derive the  $\beta$ -expansion of the Helmholtz free energy of the spin-1  $XXZ$  Heisenberg model up to order  $\beta^5$ . The analytical expansion obtained is valid for all phases of this model. Our curves of the specific heat fit well Blöte's numerical results[2] in the high temperature regime.

PACS numbers: 02.50.-r, 05.50.+q, 05.30.Fk

Keywords: Chain Models; Mathematical Physics; Finite Temperature; Statistical Mechanics

The spin-1  $XXZ$  Heisenberg model is non-integrable and therefore cannot be solved by the algebraic Bethe *ansatz* method[3]. Since the 70's its thermodynamics has been studied numerically[2, 4, 5, 6, 7]. More recently, Yamamoto and Miyashita[5, 6, 7] applied a Monte Carlo method to study the specific heat, static magnetic susceptibility and magnetization of rings and chains of different sizes as a function of temperature and extended those results to the thermodynamic limit. Bao *et al.*[8] derived a set of self-consistent equations to approximately obtain the thermodynamics of this model for all range of temperature, and solved those equations numerically.

The cummulant series[9] yields a  $\beta$ -expansion of the Helmholtz free energy for exact integrable as well as for non-integrable models. Recently Rojas *et al.*[1] showed for any one-dimensional chain model with periodic boundary condition, invariance under spatial translation and interaction between nearest neighbours that, in the thermodynamic limit, the coefficient of the high temperature expansion of arbitrary order  $\beta^n$  can be derived from an auxiliary function  $\varphi$ .

Our aim in this work is to apply the results of Ref. 1 to derive an analytical  $\beta$ -expansion of the Helmholtz free energy per site of the anisotropic spin-1  $XXZ$  Heisenberg model with single-ion anisotropy, up to order  $\beta^5$ .

The hamiltonian of the spin-1  $XXZ$  Heisenberg model with anisotropy[10] is

$$H = \sum_{i=1}^N J (S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+) + \Delta S_i^z S_{i+1}^z - h S_i^z + D (S_i^z)^2, \quad (1)$$

where  $N$  is the number of sites in the periodic chain,  $\Delta$  the anisotropy constant,  $h$  the external magnetic field in the  $z$ -axis and  $D$  the single-ion anisotropy parameter. We have  $S_i^\pm \equiv \frac{1}{\sqrt{2}} (S_i^x \pm i S_i^y)$ . We do not fix the sign of

any constant in eq.(1).

Applying the results of Ref. 1 to the spin-1  $XXZ$  Heisenberg model with single-ion anisotropy, we obtain its Helmholtz free energy  $\mathcal{W}(\beta)$  up to order  $\beta^5$ :

$$\begin{aligned} \mathcal{W}(\beta) = & -\frac{\ln(3)}{\beta} + \frac{2}{3} D + \\ & + \left( -\frac{1}{9} D^2 - \frac{4}{9} J^2 - \frac{1}{3} h^2 - \frac{2}{9} \Delta^2 \right) \beta + \\ & + \left( -\frac{1}{81} D^3 + \frac{4}{9} \Delta h^2 + \frac{4}{27} \Delta^2 D - \frac{4}{27} J^2 D + \right. \\ & \quad \left. + \frac{1}{9} h^2 D - \frac{1}{9} J^2 \Delta \right) \beta^2 + \\ & + \left( \frac{1}{36} h^4 - \frac{1}{54} \Delta^4 + \frac{1}{324} D^4 + \frac{13}{81} J^2 \Delta^2 + \right. \\ & \quad \left. + \frac{1}{54} h^2 D^2 + \frac{7}{162} J^4 - \frac{8}{27} \Delta h^2 D + \frac{5}{27} J^2 h^2 + \right. \\ & \quad \left. - \frac{10}{27} \Delta^2 h^2 + \frac{1}{27} J^2 D^2 \right) \beta^3 + \\ & + \left( \frac{7}{243} J^2 D^3 + \frac{1}{18} J^4 \Delta + \frac{1}{972} D^5 + \right. \\ & \quad \left. - \frac{1}{27} D J^2 h^2 + \frac{1}{81} J^2 \Delta D^2 - \frac{1}{81} D \Delta^4 + \right. \\ & \quad \left. - \frac{11}{162} D J^2 \Delta^2 - \frac{4}{243} \Delta^2 D^3 + \frac{22}{81} h^2 \Delta^3 + \right. \\ & \quad \left. - \frac{10}{27} J^2 \Delta h^2 - \frac{1}{36} D h^4 + \frac{8}{27} \Delta^2 h^2 D + \right. \\ & \quad \left. - \frac{1}{162} D^3 h^2 + \frac{1}{36} J^2 \Delta^3 + \frac{13}{162} D J^4 + \right. \\ & \quad \left. - \frac{4}{27} h^4 \Delta \right) \beta^4 + \end{aligned}$$

$$\begin{aligned}
& + \left( -\frac{1}{18} D^2 h^2 J^2 + \frac{16}{81} D h^4 \Delta - \frac{2}{81} D^2 h^2 \Delta^2 + \right. \\
& + \frac{5}{243} D^2 \Delta^4 - \frac{77}{486} h^2 \Delta^4 + \frac{13}{1620} D^2 J^4 + \\
& - \frac{31}{324} J^4 h^2 - \frac{5}{1944} D^4 h^2 - \frac{7}{14580} D^4 J^2 + \\
& - \frac{53}{4860} J^2 \Delta^4 + \frac{29}{81} h^4 \Delta^2 - \frac{1241}{14580} \Delta^2 J^4 + \\
& - \frac{19}{324} J^2 h^4 + \frac{1}{648} D^2 h^4 - \frac{1}{729} D^4 \Delta^2 + \\
& + \frac{8}{243} h^2 D^3 \Delta - \frac{52}{243} h^2 D \Delta^3 + \frac{26}{81} J^2 h^2 \Delta^2 + \\
& - \frac{76}{1215} J^2 D^2 \Delta^2 + \frac{2}{1215} \Delta D^3 J^2 - \frac{2}{81} D J^2 \Delta^3 + \\
& + \frac{2}{81} D \Delta J^4 - \frac{7}{87480} D^6 - \frac{13}{3240} h^6 + \\
& \left. + \frac{173}{43740} \Delta^6 - \frac{131}{43740} J^6 + \frac{14}{81} h^2 D \Delta J^2 \right) \beta^5 + \\
& + \mathcal{O}(\beta^6). \tag{2}
\end{aligned}$$

We point out that this analytic expansion is valid for *any* arbitrary set of values of the parameters  $J$ ,  $\Delta$ ,  $D$  and  $h$  in the high temperature regime. Therefore it applies equally well to any phase of the one-dimensional spin-1  $XXZ$  Heisenberg model[11, 12, 13]. The coefficients for each order in  $\beta$  in the expansion (2) are exact.

In addition, from eq.(2) we see that the thermodynamic properties of this model are insensitive to the sign of the constants  $J$  and  $h$ . From now on we take  $J \geq 0$ . By redefining all the parameters of the hamiltonian (1), eq.(2) becomes an expansion in powers of  $(J\beta)^n$ .

A beautiful numerical work by Blöte[2] tabulates numerical values for the specific heat at varied temperatures (including the high temperature region) for the  $s = 1/2$  and  $s = 1$  Heisenberg models. In all his calculations Blöte has set  $h = 0$ . To verify the correctness of our expansion (2) we compare the analytical expression for the specific heat derived from it with the numerical results of Ref. 2. Comparison of the hamiltonian (1) in Ref. 2 and ours, given by eq.(1), yields the following correspondence of parameters:  $J_{\parallel} = -\Delta/2$  and  $J_{\perp} = -J/2$ . In Fig. 1 we compare our analytical expression for the specific heat derived from eq.(2) for anisotropic Heisenberg models in the absence of single-ion anisotropy ( $D = 0$ ) with different values of  $J$  in the antiferromagnetic phase. In the high temperature region our curves fit well the numerical data.

It is certainly important to verify that eq.(2) applies equally well to the ferromagnetic phase of the spin-1 Heisenberg model. To show this we have Fig. 2 where we plot the specific heat, in the high temperature region, of the isotropic Heisenberg model with different values for the single-ion parameter  $D$  in this phase. As in Fig. 1, our curves in Fig. 2 fit pretty well Blöte's numerical results in the high temperature region.

Our analytical results for the specific heat, obtainable

from (2), fully agrees in its isotropic limit with that of section 5.2 of Ref. 2 in the high temperature limit. We also recover the high temperature static magnetic susceptibility for the isotropic Heisenberg model [eq.(4.5a) of Ref. 6]. However, we do not agree with the high temperature limit of the correlation function for the z-component of spin between nearest neighbors  $\langle S_0^z S_1^z \rangle$  of Ref. 8. The correct limit is  $\langle S_0^z S_1^z \rangle \approx -\frac{4}{9} \Delta \beta$ .

In summary, in this letter we applied the method of Ref. 1 to a non-integrable spin-1  $XXZ$  Heisenberg model with single-ion anisotropy in the presence of an external magnetic field in the  $z$ -direction. We obtained the analytical  $\beta$ -expansion of the Helmholtz free energy per site of the model up to order  $\beta^5$ . Each coefficient in expansion (2) is exact and valid for *any* phase of the model.

Our curves in the high temperature region fit well Blöte's numerical results for the spin-1  $XXZ$  Heisenberg model. We recover the few analytical results known in the literature[2, 6] about the isotropic spin-1 Heisenberg model. We correct the high temperature limit of  $\langle S_0^z S_1^z \rangle$  of Ref. 8. For the first time in the literature we have analytical results easily handled for the anisotropic spin-1 Heisenberg model with single-ion anisotropy and in the presence of an external magnetic field.

Finally we should mention that our calculation procedures have been implemented in the symbolic computational language Maple[14]. Currently, refinements are being made so that the expansion of the Helmholtz free energy (2) for the  $s = 1$  Heisenberg model can be extended to higher orders in  $\beta$ .

An enlarged version of this letter will be published elsewhere.

O. R. thanks CLAF for financial support. E.V.C.S. thanks CNPq for financial support. S.M. de S. thanks FAPEMIG for partial financial support. M.T.T. thanks CNPq and FAPERJ for partial financial support. WAMM acknowledges FAPEMIG and CNPq for partial financial support.

---

\* Corresponding author: mtt@if.uff.br

- [1] Onofre Rojas, S. M. de Souza and M. T. Thomaz, J. Math. Phys. **43** (2002) 1390.
- [2] H.W.J. Blöte, Physica **79B** (1975) 427.
- [3] M. Takahashi, *Thermodynamics of One-Dimensional Solvable Models*, Cambridge Univ. Press. (1999).
- [4] T. De Neef, Phys. Lett. **47A** (1974) 51.
- [5] S. Yamamoto and S. Miyashita, Phys. Rev. **B48** (1993-I) 9528.
- [6] S. Yamamoto and S. Miyashita, Phys. Rev. **B50** (1994-I) 6277.
- [7] S. Yamamoto and S. Miyashita, Phys. Rev. **B51** (1995-II) 3649.
- [8] S.-Q. Bao, H. Zhao, J.-L. Shen and G.-Z. Yang, Phys. Rev. **B53** (1996-II) 735.

- [9] G.S. Rushbrooke, G.A. Baker Jr. and P.J. Wood, *Phase Transitions and Critical Phenomena, vol. 3. Series Expansions for Lattice Models*, ed. C. Domb and M.S. Green, page 246, Academic Press (London, 1974).
- [10] H.-t. Wang, J.-l. Shen, K.-r. Li and Z.-b. Su, Phys. Rev. **B49** (1994-II) 12805.
- [11] S.T. Chui and K.B. Ma, Phys. Rev. **B29** (1984) 1287.
- [12] U. Glaus and T. Schneider, Phys. Rev. **B30** (1984) 215.
- [13] J. Sólyom and T.A.L. Ziman, Phys. Rev. **B30** (1984) 3980.
- [14] E.V.Corrêa Silva, *Maple V procedures for analytical high-temperature expansion of the Helmholtz free energy in Heisenberg chain models*. In preparation.




FIG. 1: Comparison of our analytic expression for the specific heat and Blöte's numerical result[2] for the anisotropic Heisenberg model with different anisotropies and  $h = 0$ . Solid lines stand for the specific heat of our analytical result and dotted lines correspond to data from Ref. 2.




FIG. 2: Comparison of our analytic expression for the specific heat and Blöte's numerical results[2] for the isotropic Heisenberg model with different single-ion parameters and  $h = 0$ . Solid lines stand for our results and dotted lines correspond to data from [2].