

Quantal Density Functional Theory of Degenerate States

Viraht Sahni and Xiao-Yin Pan

*Department of Physics, Brooklyn College and the Graduate School
of the City University of New York, New York, New York 10016.*

(Dated: March 22, 2022)

Abstract

The treatment of degenerate states within Kohn-Sham density functional theory (KS-DFT) is a problem of longstanding and current interest. We propose a solution to this mapping from the interacting degenerate system to that of the noninteracting fermion model whereby the equivalent density and energy are obtained via the unifying physical framework of quantal density functional theory (Q-DFT). We describe the Q-DFT of *both ground* and *excited* degenerate states, and for the cases of *both pure* state and *ensemble* v-representable densities. The Q-DFT description further provides a rigorous physical interpretation of the corresponding KS-DFT energy functionals of the density, ensemble density, bidensity and ensemble bidensity, and of their respective functional derivatives. We conclude with examples of the mappings within Q-DFT.

The treatment of degenerate states within the context of Kohn-Sham density functional theory KS-DFT [1] is a problem of longstanding [2] and continued recent [3, 4] interest. The basic idea underlying the KS-DFT methodology is the mapping from the interacting electronic system as described by Schrödinger theory to that of a system of *noninteracting fermions* such that the equivalent density and total energy are obtained. The existence of the model system is an assumption. Further, there is the distinction between the S system of noninteracting fermions whereby the equivalent density is obtained from a *single* Slater determinant, and the noninteracting system whose orbitals could be degenerate so that the density is obtained from a weighted sum of the Slater determinants constructed from the orbitals. The following cases have been considered. The mapping from: (a) a *pure* degenerate ground state[5]; (b) a *pure* degenerate excited state[2]. In addition, maps to obtain the density and energy constructed from (c) an *ensemble* of pure degenerate ground states[4], and (d) an *ensemble* of pure degenerate excited states[3], have been developed. The interest in the ensemble cases stems from a *ground* state theorem due to Levy [6] and Lieb [7]. According to the theorem, most ensemble densities constructed from pure degenerate ground states are not *interacting* v-representable. In other words, no *single* ground state wave function of the Schrödinger Hamiltonian will yield this ensemble density. Such ensemble densities are said to be *ensemble* v-representable. The translation of the theorem to the S system [6] means that there is no *single* Slater determinant that leads to this ensemble density. At this time, the question of interacting v-representability of the ensemble density of degenerate excited states is still unanswered [8]. In this paper we propose a solution to the problem of mapping from the interacting degenerate system to that of the equivalent noninteracting fermion model within the unifying physical framework of quantal density functional theory (Q-DFT).

In contrast to the present work, the KS-DFT description of the noninteracting system is in terms of an energy functional of the density, and its functional derivative. For case (a), the energy is a functional of the degenerate ground state density; for (b), the energy is a *bidensity* functional of the ground and excited state densities, with the functional derivative taken at the excited state density; for (c), the energy, which is a functional of the ground state ensemble density, is constructed by the ensemble generalization of the coupling constant scheme; and for (d) the energy is also a *bidensity* functional, in this instance of

the ground state and excited state ensemble densities, with the functional derivative taken at the ensemble density.

We describe here the Q-DFT of *both ground* and *excited* degenerate states, and for the cases of *both* the *pure* state and *ensemble* v-representable densities. Q-DFT is an alternate description [9, 10] of the mapping from the interacting to the noninteracting system. A key point to note is that within Q-DFT the densities concerned are always obtained from solution of the Schrödinger equation. Thus, the pure state density, and *each* component of the ensemble density, are interacting v-representable. The assumption of existence of an S system in Q-DFT therefore means that the pure state density and *each* component of the ensemble density are also noninteracting v-representable. (By an S system we mean a noninteracting fermion system whose wavefunction is a single Slater determinant, and which maybe in a ground or excited state). In Q-DFT, the local electron interaction potential energy $v_{ee}(\mathbf{r})$ of the S system, and the expression for the total energy are in terms of ‘classical’ fields whose quantal sources are quantum-mechanical expectations of Hermitian operators. We begin by (i) describing the Q-DFT of the *individual* degenerate pure state. For the mapping from a degenerate *ground* state of the interacting system, the corresponding S system is in its *ground* state. For the mapping from a degenerate *excited* state, the state of the S system is *arbitrary* in that it may be in a *ground* or *excited* state. In either case, the highest occupied eigenvalue is the negative of the ionization potential. (The mapping from a nondegenerate excited state [10] is similar.) For the ground *and* excited state ensemble cases, we describe *two* different schemes within Q-DFT. Thus, (ii) in the first, the corresponding noninteracting system ensemble density is obtained by constructing g S systems, where g is the degeneracy of the state. Once again for excited states, the g S systems may either be in a ground or excited state or a combination of the two. Next, (iii) we describe the Q-DFT whereby the ensemble density is obtained from a *single* noninteracting fermion system whose orbitals could be degenerate. The construction of this model system is a consequence of the linearity of the differential virial theorem. Here the highest occupied eigenvalue is degenerate, and the ensemble density is obtained from the resulting Slater determinants as described by Ullrich and Kohn [4] whose work in turn is based on that of Chayes et al [11]. Again for the mapping from an excited state, the noninteracting system may be in a ground or excited state. The Q-DFT description

then (*iv*) provides the physics underlying *all* the various KS-DFT degenerate state energy density and bidensity functionals, and of their functional derivatives. Finally, (*v*) we present examples that demonstrate the above mappings within Q-DFT.

A. The Q-DFT of the bound *individual* degenerate pure state is as follows. The Schrodinger equation for a degenerate state whether ground or excited is

$$\hat{H}\Psi_{n,\eta}(\mathbf{X}) = [\hat{T} + \hat{V} + \hat{U}]\Psi_{n,\eta}(\mathbf{X}) = E_n \Psi_{n,\eta}(\mathbf{X}), \quad (1)$$

$\hat{T} = -\frac{1}{2} \sum_i \nabla_i^2$, $\hat{V} = \sum_i v(\mathbf{r}_i)$, $\hat{U} = \frac{1}{2} \sum_{i,j}' \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$, where $\Psi_{n,\eta}(\mathbf{X})$ and E_n are a bound degenerate state wave function and energy, n corresponds to the state, and $\eta = 1, \dots, g_n$ the degeneracy, $\mathbf{X} = \mathbf{x}_1, \dots, \mathbf{x}_N$, $\mathbf{x} = \mathbf{r}\sigma$, with σ the spin coordinate. As the equations to follow are valid for arbitrary states, we drop the subscript n . The degenerate pure state density $\rho_\eta(\mathbf{r}) = \langle \Psi_\eta | \hat{\rho} | \Psi_\eta \rangle$, where $\hat{\rho} = \sum_i \delta(\mathbf{r} - \mathbf{r}_i)$, and the energy $E_\eta = \langle \Psi_\eta | \hat{H} | \Psi_\eta \rangle$.

The corresponding differential equation for the S system of noninteracting fermions with the same density is

$$\left[-\frac{1}{2} \nabla^2 + v(\mathbf{r}) + v_{ee,\eta}(\mathbf{r})\right] \phi_i(\mathbf{x}) = \varepsilon_i \phi_i(\mathbf{x}); \quad i = 1, \dots, N, \quad (2)$$

with

$$\rho_\eta(\mathbf{r}) = \langle \Phi_\eta \{ \phi_i \} | \hat{\rho} | \Phi_\eta \{ \phi_i \} \rangle = \sum_{i,\sigma} |\phi_i(\mathbf{x})|^2, \quad (3)$$

and $\Phi_\eta \{ \phi_i \}$ is the *single* Slater determinant of the orbitals $\phi_i(\mathbf{x})$. This is the S system wave function. The electron-interaction potential energy $v_{ee,\eta}(\mathbf{r})$ is representative of electron correlations due to the Pauli exclusion principle, Coulomb repulsion, and Correlation-Kinetic effects. Correlation-Kinetic contributions to the potential energy are a consequence of the difference in kinetic energy between the interacting and noninteracting systems. The potential energy $v_{ee,\eta}(\mathbf{r})$ is the work done to move a model Fermion in the force of a conservative field $\mathcal{F}_\eta(\mathbf{r})$:

$$v_{ee,\eta}(\mathbf{r}) = - \int_{\infty}^{\mathbf{r}} \mathcal{F}_\eta(\mathbf{r}') \cdot d\mathbf{l}', \quad (4)$$

where $\mathcal{F}_\eta(\mathbf{r}) = \mathcal{E}_{ee,\eta}(\mathbf{r}) + \mathcal{Z}_{tc,\eta}(\mathbf{r})$. The fields $\mathcal{E}_{ee,\eta}(\mathbf{r})$ and $\mathcal{Z}_{tc,\eta}(\mathbf{r})$ are not necessarily conservative. Their sum always is. The electron-interaction field $\mathcal{E}_{ee,\eta}(\mathbf{r})$ is representative of Pauli and Coulomb correlation: $\mathcal{E}_{ee,\eta}(\mathbf{r}) = \mathbf{e}_{ee,\eta}(\mathbf{r})/\rho_\eta(\mathbf{r})$, where the electron-interaction ‘force’

$\mathbf{e}_{ee,\eta}(\mathbf{r})$ is obtained via Coulomb's law as $\mathbf{e}_{ee,\eta}(\mathbf{r}) = \int d\mathbf{r}' P_\eta(\mathbf{r}\mathbf{r}')(\mathbf{r} - \mathbf{r}')/|\mathbf{r} - \mathbf{r}'|^3$, where $P_\eta(\mathbf{r}\mathbf{r}') = \langle \Psi_\eta | \hat{P}(\mathbf{r}\mathbf{r}') | \Psi_\eta \rangle$, with $\hat{P}(\mathbf{r}\mathbf{r}') = \sum_{i,j}' \delta(\mathbf{r} - \mathbf{r}_i) \delta(\mathbf{r}' - \mathbf{r}_j)$. Equivalently, the field $\mathcal{E}_{ee,\eta}(\mathbf{r})$ may be thought of as being due to its quantal source, the pair-correlation density $g_\eta(\mathbf{r}\mathbf{r}') = P_\eta(\mathbf{r}\mathbf{r}')/\rho_\eta(\mathbf{r})$. The Correlation-Kinetic field $\mathcal{Z}_{tc,\eta}(\mathbf{r}) = \mathcal{Z}_{s,\eta}(\mathbf{r}) - \mathcal{Z}_\eta(\mathbf{r})$, $\mathcal{Z}_\eta(\mathbf{r}) = \mathbf{z}_\eta(\mathbf{r}; [\gamma_\eta])/\rho_\eta(\mathbf{r})$, $\mathcal{Z}_{s,\eta}(\mathbf{r}) = \mathbf{z}_{s,\eta}(\mathbf{r}; [\gamma_{s,\eta}])/\rho_\eta(\mathbf{r})$, and where $\mathcal{Z}_\eta(\mathbf{r})$ and $\mathcal{Z}_{s,\eta}(\mathbf{r})$ are the interacting and S system kinetic fields, respectively. The kinetic 'force' $\mathbf{z}_\eta(\mathbf{r})$ is defined by its component $z_{\eta,\alpha} = 2 \sum_\beta \partial t_{\alpha,\beta} / \partial r_\beta$, with $t_{\alpha,\beta}(\mathbf{r}, [\gamma_\eta]) = \frac{1}{4} [\partial^2 / \partial r'_\alpha \partial r''_\beta + \partial^2 / \partial r'_\beta \partial r''_\alpha] \gamma_\eta(\mathbf{r}', \mathbf{r}'')|_{\mathbf{r}'=\mathbf{r}''=\mathbf{r}}$ the kinetic energy density tensor. The source of the kinetic field $\mathcal{Z}_\eta(\mathbf{r})$ is the spinless single particle density matrix $\gamma_\eta(\mathbf{r}, \mathbf{r}') = \langle \Psi_\eta | \hat{X} | \Psi_\eta \rangle$, $\hat{X} = \hat{A} + i\hat{B}$, $\hat{A} = \frac{1}{2} \sum_j [\delta(\mathbf{r}_j - \mathbf{r}) T_j(\mathbf{a}) + \delta(\mathbf{r}_j - \mathbf{r}') T_j(-\mathbf{a})]$, $\hat{B} = -\frac{i}{2} \sum_j [\delta(\mathbf{r}_j - \mathbf{r}) T_j(\mathbf{a}) - \delta(\mathbf{r}_j - \mathbf{r}') T_j(-\mathbf{a})]$, $T_j(\mathbf{a})$ is a translation operator, and $\mathbf{a} = \mathbf{r}' - \mathbf{r}$. The field $\mathcal{Z}_s(\mathbf{r})$ is defined in a similar manner in terms of the S system Dirac density matrix $\gamma_{s,\eta}(\mathbf{r}, \mathbf{r}') = \langle \Phi_\eta \{ \phi_i \} | \hat{X} | \Phi_\eta \{ \phi_i \} \rangle = \sum_{i,\sigma} \phi_i^*(\mathbf{r}\sigma) \phi_i(\mathbf{r}'\sigma)$.

The proof of Eq.(4) follows by equating the differential virial theorems [10, 12] for the interacting and S systems which are, respectively

$$\nabla v(\mathbf{r}) = -\mathbf{F}_\eta(\mathbf{r}) \text{ and } \nabla v(\mathbf{r}) = -\mathbf{F}_{s,\eta}(\mathbf{r}), \quad (5)$$

where $\mathbf{F}_\eta(\mathbf{r}) = -\mathcal{E}_{ee,\eta}(\mathbf{r}) + \mathcal{D}_\eta(\mathbf{r}) + \mathcal{Z}_\eta(\mathbf{r})$, $\mathbf{F}_{s,\eta}(\mathbf{r}) = \nabla v_{ee,\eta}(\mathbf{r}) + \mathcal{D}_\eta(\mathbf{r}) + \mathcal{Z}_{s,\eta}(\mathbf{r})$, the differential density field $\mathcal{D}_\eta(\mathbf{r}) = \mathbf{d}_\eta(\mathbf{r})/\rho_\eta(\mathbf{r})$, $\mathbf{d}_\eta(\mathbf{r}) = -\frac{1}{4} \nabla \nabla^2 \rho_\eta(\mathbf{r})$. Thus, one obtains

$$\nabla v_{ee,\eta}(\mathbf{r}) = -\mathcal{F}_\eta(\mathbf{r}), \quad (6)$$

from which the interpretation of Eq. (4) follows.

The total energy of the degenerate state η is then

$$E_\eta = T_{s,\eta} + \int \rho_\eta(\mathbf{r}) v(\mathbf{r}) d\mathbf{r} + E_{ee,\eta} + T_{c,\eta}, \quad (7)$$

where $T_{s,\eta} = \langle \Phi_\eta \{ \phi_i \} | \hat{T} | \Phi_\eta \{ \phi_i \} \rangle$ is the S system kinetic energy, and the electron-interaction $E_{ee,\eta}$ and Correlation-Kinetic $T_{c,\eta}$ energies in terms of the respective fields are

$$E_{ee,\eta} = \int d\mathbf{r} \rho_\eta(\mathbf{r}) \mathbf{r} \cdot \mathcal{E}_{ee,\eta}(\mathbf{r}), \text{ and} \quad (8)$$

$$T_{c,\eta} = \frac{1}{2} \int d\mathbf{r} \rho_\eta(\mathbf{r}) \mathbf{r} \cdot \mathcal{Z}_{tc,\eta}(\mathbf{r}). \quad (9)$$

These expressions are independent of whether the fields $\mathcal{E}_{ee,\eta}(\mathbf{r})$ and $\mathcal{Z}_{tc,\eta}(\mathbf{r})$ are conservative or not.

The S system whereby the density and total energy equivalent to that of the interacting system degenerate state η is defined by Eqs.(2)-(4) and (7). If the degenerate state is excited, the S system may be constructed to be either in a ground or excited state. Since the electron-interaction field remains unchanged, the difference between the corresponding potential energies is independent of the Pauli principle and Coulomb repulsion and due entirely to the corresponding Correlation-Kinetic fields $\mathcal{Z}_{tc,\eta}(\mathbf{r})$. Hence, the potential energy $v_{ee,\eta}(\mathbf{r})$ is different depending on whether the S system is in a ground or excited state. In either case, the highest occupied eigenvalue of the S system differential equation is the negative of the ionization potential. This follows by equating the asymptotic structure of the density for the interacting and S systems.

In the transformation from an excited pure degenerate state to an S system in its ground state, the fact that the interacting system wave function has nodes is of no relevance. By construction, the S and interacting system density $\rho(\mathbf{r})$ are equivalent, and the density $\rho(\mathbf{r}) \geq 0$. Such a mapping for an excited pure nondegenerate state has been demonstrated in Ref. [10].

B. We next describe the first of *two* ways of obtaining the *ensemble* density and energy of the degenerate states via Q-DFT. The interacting system ensemble density matrix operator $\hat{D}(\mathbf{X}\mathbf{X}')$ is defined as

$$\hat{D}(\mathbf{X}\mathbf{X}') = \sum_{\eta=1}^g \omega_{\eta} \Psi_{\eta}^*(\mathbf{X}) \Psi_{\eta}(\mathbf{X}'); \quad \sum_{\eta=1}^g \omega_{\eta} = 1; \quad 0 \leq \omega_{\eta} \leq 1, \quad (10)$$

so that the ensemble density $\rho_{ens}(\mathbf{r})$ and energy E_{ens} are respectively

$$\rho_{ens}(\mathbf{r}) = tr(\hat{D}\hat{\rho}) = \sum_{\eta=1}^g \omega_{\eta} \rho_{\eta}(\mathbf{r}), \quad (11)$$

and

$$E_{ens} = tr(\hat{D}\hat{H}) = \sum_{\eta=1}^g \omega_{\eta} E_{\eta}, \quad (12)$$

with $\rho_{\eta}(\mathbf{r})$ and E_{η} as defined previously. (There are ensemble densities that cannot be represented by a single Slater determinant. However, its pure state component density can

always be reproduced by an S system).

For *each* degenerate state η , the density $\rho_\eta(\mathbf{r})$ and energy E_η can be constructed from an S system as described in part A. Thus, the ensemble density and energy of Eqs. (11) and (12) may be obtained from g S systems. Each S system contributing to the ensemble density may be in a ground or excited state. Note that the electron-interaction potential energy $v_{ee,\eta}(\mathbf{r})$ for *each* of the g S systems will be different. Further, $v_{ee,\eta}(\mathbf{r})$ will be different depending on whether the particular S system is in a ground or excited state as explained previously. Thus, the ensemble density and energy within Q-DFT are obtained by replacing the $\rho_\eta(\mathbf{r})$ and E_η on the right hand sides of Eqs. (11) and (12) by the corresponding S system equivalents of Eqs. (3) and (7), respectively.

C. The ensemble density and energy may also be determined from a noninteracting fermion system whose orbitals could be degenerate as constructed within Q-DFT. According to Chayes et al [11], the ground state ensemble density may be determined as a unique weighted sum of squares of a finite number g of degenerate wave functions of this system. The potential energy $v_{ee}(\mathbf{r})$ of these noninteracting fermions is then determined via Q-DFT as follows. Rewrite the interacting and noninteracting system differential virial theorems of Eq. (5) as

$$\rho_{ens}(\mathbf{r})\nabla v(\mathbf{r}) = \sum_{\eta=1}^g \omega_\eta \mathbf{f}_\eta(\mathbf{r}), \quad (13)$$

where $\mathbf{f}_\eta(\mathbf{r}) = \mathbf{e}_{ee,\eta}(\mathbf{r}) - \mathbf{d}_\eta(\mathbf{r}) - \mathbf{z}_\eta(\mathbf{r})$, and

$$\rho_{ens}(\mathbf{r})\nabla v(\mathbf{r}) = \sum_{\eta=1}^g \omega_\eta \mathbf{f}_{s,\eta}(\mathbf{r}), \quad (14)$$

where $\mathbf{f}_{s,\eta}(\mathbf{r}) = \rho_\eta(\mathbf{r})\nabla v_{ee,\eta}(\mathbf{r}) - \mathbf{d}_\eta(\mathbf{r}) - \mathbf{z}_{s,\eta}(\mathbf{r})$. Equating Eqs. (13) and (14) leads to

$$\sum_{\eta=1}^g \omega_\eta \rho_\eta(\mathbf{r})\nabla v_{ee,\eta}(\mathbf{r}) = \sum_{\eta=1}^g \omega_\eta \mathbf{q}_\eta(\mathbf{r}), \quad (15)$$

where $\mathbf{q}_\eta(\mathbf{r}) = \mathbf{e}_{ee,\eta}(\mathbf{r}) + \mathbf{z}_{t_e,\eta}(\mathbf{r})$, $\mathbf{z}_{t_e,\eta}(\mathbf{r}) = \mathbf{z}_{s,\eta}(\mathbf{r}) - \mathbf{z}_\eta(\mathbf{r})$. Eq. (15) is a consequence of the linearity of the differential virial theorem [12]. As we require a *single* effective potential energy $v_s(\mathbf{r}) = v(\mathbf{r}) + v_{ee}(\mathbf{r})$, we replace $v_{ee,\eta}(\mathbf{r})$, in Eq. (15) by $v_{ee}(\mathbf{r})$ to obtain

$$\nabla v_{ee}(\mathbf{r}) = -\mathbf{Q}(\mathbf{r}), \quad (16)$$

where $\mathbf{Q}(\mathbf{r}) = -(\sum_{\eta=1}^g \omega_{\eta} \mathbf{q}_{\eta}(\mathbf{r}))/\rho_{ens}(\mathbf{r})$, Thus, the electron-interaction potential energy $v_{ee}(\mathbf{r})$ is the work done in the conservative field $\mathbf{Q}(\mathbf{r})$:

$$v_{ee}(\mathbf{r}) = - \int_{\infty}^{\mathbf{r}} \mathbf{Q}(\mathbf{r}') \cdot d\mathbf{l}'. \quad (17)$$

Note that the components $\mathbf{q}_{\eta}(\mathbf{r})$ are conservative so that $\mathbf{Q}(\mathbf{r})$ is conservative, and hence $v_{ee}(\mathbf{r})$ is path independent.

For the occupation of orbitals we follow Ullrich-Kohn [4]. Accordingly, all levels are occupied except the highest (h) which is q - fold degenerate and partially occupied. The number of the model fermions in these levels are $N^h \leq 2q$. The ensemble density which is a weighted sum of the degenerate Slater determinants is

$$\rho_{ens}(\mathbf{r}) = \sum_{i,\sigma}^{N-N^h} |\phi_i(\mathbf{x})|^2 + \sum_{i,\sigma}^q f_i |\phi_i^h(\mathbf{x}; R)|^2, \quad (18)$$

with $0 \leq f_i \leq 1$, and $f_i = \sum_{\eta=1}^g \omega_{\eta} \theta_{i,\eta}$ where $\theta_{i,\eta} = 1$ if the orbital $\phi_i^h(\mathbf{x}; R)$ occurs in the determinant $\Phi_{\eta}\{\phi_i\}$, and 0 otherwise. Here the $\phi_i^h(\mathbf{x}; R)$ are appropriately rotated (R) orbitals determined self-consistently together with the f_i and the lower lying orbitals leading to the ensemble density. The ensemble energy is obtained from the g Slater determinants as in part B. Once again for an excited state ensemble density, the corresponding noninteracting system may be in a ground or excited state.

Note that the methodology of construction of the g S systems of part B also follows from Eq. (15). We believe that it is easier to construct the g S systems of part B than it is to construct the single noninteracting system of part C. This is because each of the g S systems may be constructed independently.

From the above degenerate state Q-DFT description it is then possible to provide a rigorous physical interpretation for *each* energy functional and functional derivative of the corresponding KS-DFT. In each case, the local potential energy of the model fermions is the work done in a conservative field. The energy in turn may be expressed in terms of the components of this field. Thus, for example, the KS-DFT degenerate *ground* state electron-interaction energy functional $E_{ee}^{KS}[\rho_{ens}]$ [4] of the ensemble density is the ensemble sum of the electron-interaction $E_{ee,\eta}$ and Correlation-Kinetic $T_{c,\eta}$

energies. The functional derivative is the work done to move the model fermion in the conservative field $\mathbf{Q}(\mathbf{r})$. The *same* interpretation applies to the *bidity* energy functional $E_{ee}^{KS}[\rho_{gr}, \rho_{ens}]$ [3] of degenerate *excited* state KS-DFT, and of its functional derivative.

The Q-DFT mapping from a pure degenerate excited state to an S system can be demonstrated via the first excited *triplet* state of the exactly solvable Hooke's atom [13]. This atom is comprised of two electrons with a harmonic external potential energy $v(\mathbf{r}) = \frac{1}{2}\omega r^2$. The triplet state wave functions are of the form

$$\Psi(\mathbf{r}_1\mathbf{r}_2) = C_0 e^{-\omega R^2} e^{-\omega r^2/4} [1 + C_1 \sqrt{\frac{\omega}{2}} r + C_2 (\frac{\omega}{2}) r^2 + C_3 (\frac{\omega}{2})^{3/2} r^3] Y_{l,m}(\theta, \phi), \quad (19)$$

where $Y_{l,m}(\theta, \phi)$ is the spherical harmonic with $l = 1, m = -1, 0, 1, \mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1, \mathbf{R} = (\mathbf{r}_2 + \mathbf{r}_1)/2$, and C_0, C_1, C_2, C_3 are constants. For each degenerate wave function, the corresponding S system, in either a ground or excited state, can be determined in a manner similar to the transformation of the first excited *singlet* state [10]. These results will be presented elsewhere. The ensemble density of these degenerate states then follows from the 3 S systems. (Note that the ensemble density for this two electron model is v-representable. However, the methodology for constructing the g S systems is the same whether or not the density is v-representable.) Another example is that of the noninteracting Be atom [4]. Here the ensemble density, which is not v-representable, is the weighted sum of the density of 4 S systems in the states $1s^2 2s^2, 1s^2 2p_i^2$ ($i = x, y, x$). This latter model atom is also an example [4] of the noninteracting fermion system that leads to the ensemble density with appropriately rotated highest occupied orbitals.

In conclusion, we have described via Q-DFT the physics of mapping from a degenerate state of Schrödinger theory to that of a model system of noninteracting fermions such that the equivalent density and energy are determined. The cases of *both pure* and *ensemble* v-representable densities are explained. The framework is general and formally the *same* for *both* degenerate *ground* and *excited* states.

This work was supported in part by the Research Foundation of CUNY.

-
- [1] W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
 - [2] M. Levy and A. Nagy, Phys. Rev. Lett. **83**, 4361 (1999), and references therein.
 - [3] A. Nagy and M. Levy, Phys. Rev. A **63**, 052502 (2001).
 - [4] C. A. Ullrich and W. Kohn, Phys. Rev. Lett. **87**, 093001 (2001); *ibid* **89**,156401(2002).
 - [5] W. Kohn in *Highlights of Condensed Matter Theory*, ed. by F. Bassani et al (North Holland, Amsterdam,1985).
 - [6] M. Levy, Phys. Rev. A **26**, 1200 (1982).
 - [7] E. H. Lieb, Int. J. Quantum Chem. **24**, 243 (1983).
 - [8] M. Levy, (private communication).
 - [9] V. Sahni, Phys. Rev. A **55**, 1846 (1997); Top. Curr. Chem. **182**, 1(1996); Z. Qian and V. Sahni, Phys. Rev. A **57**, 2557(1999); Phys. Rev. B **62**, 16364 (2000); Int. J. Quantum Chem. **80**, 555 (2000).
 - [10] V. Sahni, L. Massa, R. Singh, and M. Slamet, Phys. Rev. Lett. **87**, 113002 (2001); M. Slamet and V. Sahni, Int. J. Quantum Chem. **85**, 436 (2001).
 - [11] J. T. Chayes, L. Chayes, and M. B. Ruskai, J. Stat. Phys. **38**, 497(1985).
 - [12] A. Holas and N. H. March, Phys. Rev. A **51**, 2040(1995).
 - [13] M. Taut, Phys. Rev. A **48**, 3561(1993)