

# Universal reduction of pressure between charged surfaces by long-wavelength surface charge modulation

D. B. LUKATSKY AND S. A. SAFRAN

*Department of Materials and Interfaces, Weizmann Institute, Rehovot 76100, Israel*

PACS. 82.70.-y – Disperse systems; complex fluids.

PACS. 61.20.Qg – Structure of associated liquids: electrolytes, molten salts, etc..

**Abstract.** – We predict theoretically that long-wavelength surface charge modulations universally reduce the pressure between the charged surfaces with counterions compared with the case of uniformly charged surfaces with the same average surface charge density. The physical origin of this effect is the fact that surface charge modulations *always* lead to enhanced counterion localization near the surfaces, and hence, fewer charges at the midplane. We confirm the last prediction with Monte Carlo simulations.

*Introduction-* The interaction of charged interfaces is a basic problem in surface science, and is relevant to *e.g.*, colloid stabilization, membrane adhesion, microemulsion formation [1]. While many theories have focused on the simple case of homogeneously charged surfaces, real interfaces are characterized by discrete surface charge distributions. Theoretical analysis of the problem predicts two important properties of these systems: the spatial dependence of the counterion/salt density and the pressure between the interfaces [2,3,4,5,6,7,8,9,10,11,12,13,14,15]. The main question is whether one can provide *universal* predictions of the effect of surface charge discreteness (modulations) on the counterion spatial distribution and on the interaction law between the interfaces, independently of the details of the surface distribution of discrete charges (modulations). Our work shows that indeed, there are quite general effects induced by surface charge discreteness.

The principal conclusion of previous studies based on the solution of linear or quasi-linear Debye-Huckel equation with non-uniform surface charge distributions with either excess salt [2,3,15], or with counterions only [15], is the following: The discreteness (or modulations) of surface charge *always* leads to an *enhanced repulsive* interaction between the surfaces compared with the case of uniformly charged surfaces with the same average surface charge density, provided the charges (or charge modulations) on the two surfaces are in-phase. If the charges (charge modulations) on the two surfaces are out-of-phase, the repulsion can be *either enhanced or reduced* compared with the uniformly charged surface case. These studies [2,3,15] calculated the effects of charge modulations on the counterion spatial distribution and on the electrostatic potential to first order in the amplitude of modulations. The only relevant terms in the pressure are therefore due to the square of the non-zero components of the electric field,  $\vec{E}$ , at the midplane  $\sim E_x^2 + E_y^2 - E_z^2$  (see below). This contribution is always positive for the in-phase case, and leads to the *enhanced repulsion* in the in-phase case.



In this paper, we show that a consistent calculation of the pressure, including both the osmotic terms proportional to the counterion charge density at the midplane, and the electric field terms calculated *to second order in the amplitude of modulations*, results in a *universal reduction* of the pressure by long-wavelength surface charge modulations. The notion of a *universal* reduction means that the effect does not depend on a particular form, amplitude, and phase shift between surface charge modulations. The physical origin of the predicted reduction of the pressure is due to the osmotic effect and arises because surface charge modulations *always* enhance counterion localization in the vicinity of surfaces, and hence, *always* reduce the midplane counterion density. We confirm this prediction for the charge density with Monte-Carlo (MC) simulations. We emphasize that the predicted effects are due to the correlations between the counterions and the inhomogeneities of the surface charge distribution.

*Two surfaces with charge modulations-* In this section we consider two, non-uniformly charged planar surfaces with fixed surface charge densities  $-e\sigma_1(\vec{\rho})$  and  $-e\sigma_2(\vec{\rho})$  [where  $\vec{\rho} = (x, y)$ ], respectively. The surfaces whose average charges are equal and are taken to be negative, are located at  $z = -\frac{h}{2}$  and  $z = \frac{h}{2}$ , respectively. The surface electrostatic repulsion is screened by the positively charged counterions of valence  $Z$ ; the counterions are located only in the water between the surfaces, and the dielectric constant,  $\epsilon$ , ( $\epsilon \simeq 80$  for water), is assumed to be homogeneous throughout the system. The charge densities  $\sigma_1(\vec{\rho})$  and  $\sigma_2(\vec{\rho})$  obey the condition:  $\int \sigma_i(\vec{\rho}) d\vec{\rho} / A_0 = \sigma_0$ , where  $A_0$  is the surface area, and  $\sigma_0 = ZN / 2A_0$  is the average number of surface charge per unit area, with  $N$  being the total number of counterions in the system.

The system is fully specified within mean-field theory, by solving the differential equation for the potential [17],  $\varphi(\vec{\rho}, z)$ , with the boundary conditions implied by the delta function, spatially dependent surface charge distributions  $\sigma_1(\vec{\rho})$  and  $\sigma_2(\vec{\rho})$ :

$$\nabla^2 \phi(\rho, z) = -4\pi\ell_B Z \left[ Z n_0 e^{-\phi} - \delta(z + \frac{h}{2})\sigma_1(\vec{\rho}) - \delta(z - \frac{h}{2})\sigma_2(\vec{\rho}) \right], \quad |z| \leq \frac{h}{2}, \quad (1)$$

and  $\nabla^2 \phi(\vec{\rho}, z) = 4\pi\ell_B Z [\delta(z + \frac{h}{2})\sigma_1(\vec{\rho}) + \delta(z - \frac{h}{2})\sigma_2(\vec{\rho})]$ , for  $|z| \geq \frac{h}{2}$ , where there are no counterions [16]. Here  $\phi(\vec{\rho}, z) = eZ\varphi(\vec{\rho}, z)/k_B T$  is the reduced electrostatic potential,  $\ell_B = \frac{e^2}{\epsilon k_B T}$  is the Bjerrum length, and  $\delta(z)$  is the Dirac delta function. In what follows, we consider a general, modulated charge distribution described by a Fourier decomposition [14]:  $\sigma_1(\vec{\rho}) = \sigma_0 \left[ 1 + \sum_{\vec{Q} \neq 0} \varepsilon(\vec{Q}) e^{i\vec{Q} \cdot \vec{\rho}} \right]$  and  $\sigma_2(\vec{\rho}) = \sigma_0 \left[ 1 + \sum_{\vec{Q} \neq 0} \varepsilon(\vec{Q}) e^{i\vec{Q} \cdot (\vec{\rho} + \vec{b})} \right]$ , where  $\vec{b}$  is a phase shift of the modulations on the two surfaces. The coefficient  $\varepsilon(\vec{Q})$  is the amplitude of the corresponding  $\vec{Q}$  mode; for example, on a square lattice,  $\vec{Q} = 2\pi\vec{m}/a$ , and  $\vec{m} = (m_x, m_y)$ , where  $m_x$  and  $m_y$  are integers, and  $a$  is a lattice constant. For a periodic lattice of surface charges, one must keep the infinite sum over reciprocal lattice vectors,  $\vec{Q}$ , and set  $\varepsilon(\vec{Q}) = 1$ . In analogy with the analysis performed earlier for the case of a single, isolated surface [14], we expand the potential,  $\phi(\vec{\rho}, z)$ , in powers of the charge modulation amplitude,  $\varepsilon(\vec{Q})$ :  $\phi(\vec{\rho}, z) \approx \phi_0(z) + \phi_1(\vec{\rho}, z) + \phi_2(\vec{\rho}, z) + \dots$ , where  $\phi_0(z) = 2 \ln \cos k_0 z$  is the solution of the PB equation [17] with uniform surface charge densities,  $-e\sigma_0$ . The parameter  $k_0$  is determined by the charge conservation condition [17]:  $k_0 \lambda_0 \tan(\frac{k_0 h}{2}) = 1$ , where  $\lambda_0 = 1/2\pi\ell_B Z \sigma_0$ . Our objective is to obtain perturbatively the corrections  $\phi_1(\vec{\rho}, z)$  [linear in  $\varepsilon(\vec{Q})$ ] and  $\phi_2(\vec{\rho}, z)$  [quadratic in  $\varepsilon(\vec{Q})$ ]. Substituting the expansion of  $\phi(\vec{\rho}, z)$  into Eq. (1), and collecting terms linear in  $\varepsilon(\vec{Q})$ , we find that  $\phi_1(\vec{Q}, z)$  [the two-dimensional Fourier transform of  $\phi_1(\vec{\rho}, z)$ ] obeys



the following equation for  $|z| \leq \frac{h}{2}$ :

$$\left[ -\frac{\partial^2}{\partial z^2} + Q^2 + \frac{2k_0^2}{\cos^2 k_0 z} \right] \phi_1(\vec{Q}, z) = -\frac{2\varepsilon(\vec{Q})}{\lambda_0} \left[ \delta(z + \frac{h}{2}) + \delta(z - \frac{h}{2}) e^{i\vec{Q} \cdot \vec{b}} \right], \quad \vec{Q} \neq 0. \quad (2)$$

From here on, all the lengths in the problem will be rescaled in the units of  $\lambda_0$ :  $\mathcal{Q} \equiv Q\lambda_0$ ,  $\tilde{z} \equiv z/\lambda_0$ ,  $\vec{\rho} \equiv \vec{\rho}/\lambda_0$ ,  $k_0 \equiv k_0\lambda_0$ ,  $h \equiv h/\lambda_0$ , etc. The solution of this equation,  $\phi_1(\vec{Q}, \tilde{z})$ , is given in the *Appendix*.


The equation for the contribution to the potential that is second order in the surface charge modulation amplitude,  $\phi_2(\vec{\rho}, \tilde{z})$ , is obtained in a similar manner; collecting the terms quadratic in  $\varepsilon(\vec{Q})$  we find for  $|\tilde{z}| \leq \frac{h}{2}$ :  $\left[ -\nabla^2 + \frac{2k_0^2}{\cos^2 k_0 \tilde{z}} \right] \phi_2(\vec{\rho}, \tilde{z}) = \frac{k_0^2}{\cos^2 k_0 \tilde{z}} \phi_1^2(\vec{\rho}, \tilde{z})$ . This equation is solved, using the Green function for the homogeneous equation, [18]. In what follows, for the sake of simplicity, we consider two opposite cases: (i) in-phase, *i.e.*, maximum interaction,  $\vec{b} = (0, 0)$ , and (ii) out-of-phase, *i.e.*, minimum interaction,  $\vec{b} = (\frac{a}{2}, \frac{a}{2})$ . The case of arbitrary  $\vec{b}$  may be solved analytically, as well. We will consider surface charge modulations described by a single symmetric  $\vec{Q}$  mode:  $\vec{Q} = \{(\pm \frac{2\pi}{a}, 0), (0, \pm \frac{2\pi}{a})\}$ .

The average of  $\phi_2$  in the  $x-y$  plane is given in the *Appendix* in both cases. Our main prediction is the *enhancement* of the counterion density at each surface, averaged over the  $x-y$  plane,  $\langle n(\vec{\rho}, \tilde{z} = \pm \frac{h}{2}) \rangle_{\vec{\rho}}$ . The physical importance of this result is that by charge conservation it leads to the *reduction* of the midplane average density compared to the case of uniformly charged surfaces in both the in-phase,  $\vec{b} = (0, 0)$ , and out-of-phase,  $\vec{b} = (\frac{a}{2}, \frac{a}{2})$ , cases. This conclusion comes from the analysis of  $\langle n(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}}$  which takes the following form to quadratic order in  $\varepsilon(\vec{Q})$ :  $\langle n(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}} = n_0(\tilde{z})[1 + \frac{1}{2} \langle \phi_1^2(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}} - \langle \phi_2(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}}]$ , where  $n_0(\tilde{z}) = n_0 / \cos^2 k_0 \tilde{z}$  is the counterion density between uniformly charged surfaces [17],  $k_0^2 = 2\pi\ell_B Z^2 n_0$ . We emphasize that the first-order term,  $\phi_1(\vec{\rho}, \tilde{z})$ , vanishes after averaging with respect to  $\vec{\rho}$ , and that only quadratic terms in  $\varepsilon(\vec{Q})$  contribute to the average density; this observation is crucial for the correct calculation of the osmotic pressure, as we show below. The enhancement of the counterion density in the vicinity of the surfaces [and the consequent reduction of the midplane density] is due to the correlations between the counterions and the inhomogeneities of the surface charge distribution. In particular, in the asymptotic limit of small inter-surface separation,  $h \ll 1$ ,  $\mathcal{Q}h \ll 1$  the normalized average midplane counterion density has the form: (i) in the in-phase case,  $\frac{\langle n(\vec{\rho}, 0) \rangle_{\vec{\rho}}}{n_0(0)} \approx 1 - \frac{2}{3} \sum_{\vec{Q}} \frac{|\varepsilon(\vec{Q})|^2}{(2+\mathcal{Q})^2} h + \mathcal{O}(h^2)$ ; (ii) in the out-of-phase case,  $\frac{\langle n(\vec{\rho}, 0) \rangle_{\vec{\rho}}}{n_0(0)} \approx 1 - \frac{1}{6} \sum_{\vec{Q}} |\varepsilon(\vec{Q})|^2 h^2 + \mathcal{O}(h^3)$ . We stress that the strongest *reduction* of the midplane counterion density [and hence, the strongest *enhancement* of the contact density] occurs for the long-wavelength modulations. This is in a qualitative agreement with the predictions obtained for a single, isolated surface [14]. In the opposite limit of large separations,  $h \gg 1$ ,  $\mathcal{Q}h \gg 1$ , the midplane density is reduced with an asymptotic form that is identical in both limits, (i) and (ii):  $\frac{\langle n(\vec{\rho}, 0) \rangle_{\vec{\rho}}}{n_0(0)} \approx 1 - \sum_{\vec{Q}} \frac{4|\varepsilon(\vec{Q})|^2 (1+\mathcal{Q})}{(1+2\mathcal{Q}+2\mathcal{Q}^2)^2} \frac{1}{h} + \mathcal{O}(\frac{1}{h^2})$ .

To verify the predictions of the theory, we have performed MC simulations of the counterion density profiles between non-uniformly charged surfaces. The non-uniform surface charge densities were chosen to have the form:  $\sigma_1(\vec{\rho})/\sigma_0 = 1 + \varepsilon_0 \sum_{\vec{Q}} e^{i\vec{Q} \cdot \vec{\rho}}$ , and  $\sigma_2(\vec{\rho})/\sigma_0 = 1 + \varepsilon_0 \sum_{\vec{Q}} e^{i\vec{Q} \cdot (\vec{\rho} + \vec{b})}$ , respectively, where  $\vec{Q}$  takes on the four symmetric values:  $\{(\pm \frac{2\pi}{a}, 0), (0, \pm \frac{2\pi}{a})\}$ . The counterions interact via an unscreened Coulomb potential, and an Ewald 2D algorithm [19] was used to sum the Coulomb interactions; each counterion interacts with the surfaces via the potential,  $eZ\varphi_s$  generated by the *exact* solution of the Poisson equation with the chosen surface charge densities:  $\frac{eZ\varphi_s}{k_B T} = -\varepsilon_0 \sum_{\vec{Q}} e^{i\vec{Q} \cdot \vec{\rho}} (e^{-\mathcal{Q}|\tilde{z}+h/2|} + e^{i\vec{Q} \cdot \vec{b}} e^{-\mathcal{Q}|\tilde{z}-h/2|}) / \mathcal{Q}$ .



Fig. 1 – Normalized counterion density in the in-phase case,  $\tilde{n}(\tilde{z}) \equiv 2\pi\ell_B Z^2 \lambda_0^2 \langle n(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}}$  from the MC simulations (symbols) and the corresponding predictions of the analytic, perturbation theory (curves). The parameters used are:  $g \equiv \frac{\ell_B Z^2}{\lambda_0} = 0.5$ ,  $\varepsilon_0 = 1$ ,  $Q = 1$  (diamonds and dashed line). The counterion profile between uniformly charged surfaces with the same average surface densities of charge (stars and solid line) is shown for comparison. The simulations were performed with 100 particles. Inset: The pressure difference diagram,  $p - p_0 = 0$ , as a function of the rescaled wavelength of modulations,  $Q$ , and the inter-surface separation,  $h$ , in the in-phase case, where  $p_0$  is the pressure between uniformly charged surfaces. We stress that already at  $h \gtrsim 1.25$ , charge modulations reduce the pressure for any values of  $Q$ .



Using the simulations we compare the counterion density averaged in  $x-y$  plane. We plot the normalized dimensionless density,  $\tilde{n}(\tilde{z}) \equiv 2\pi\ell_B Z^2 \lambda_0^2 \langle n(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}}$  vs. the rescaled length,  $\tilde{z} \equiv z/\lambda_0$ . The analytic expressions for  $\langle n(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}}$  are given above (see also the *Appendix*) for both the in-phase and the out-of-phase cases.

The dimensionless parameter which determines the strength of the Coulomb interaction in the simulations is  $g \equiv \frac{\ell_B Z^2}{\lambda_0}$ ; other two important parameters are the amplitude of the surface charge modulation,  $\varepsilon_0$  and the lengthscale,  $Q$ , of the surface charge modulations. In Fig. 1 we plot the simulated, average density,  $\tilde{n}(\tilde{z})$ , in the in-phase case,  $\vec{b} = (0, 0)$  for values of  $g = 0.5$ ,  $\varepsilon_0 = 1$ , and  $Q = 1$  (diamonds), as well as the simulated counterion profile between uniformly charged surfaces with the same average surface charge density (stars). Additional simulations showed that the average density is higher for the smaller values of  $Q$ , similar to the conclusion obtained earlier in the case of a single, isolated surface [14]. We also note that the quantitative difference between average density profiles predicted by theory and confirmed by simulations in the in-phase and the out-of-phase cases is very small. The contact (midplane) density is always slightly higher (smaller) in the in-phase case compared with the out-of-phase case, since in the out-of-phase case the broken planar symmetry leads to the existence of an oscillating  $z$ -component of the electric field at the midplane (absent in the in-phase case), that attracts more counterions to the midplane due to the enhanced correlations. We stress again that



the perturbation theory prediction of the enhancement of the contact counterion density is universal in the sense that the enhancement occurs for all values of the parameters, *i.e.*,  $\mathcal{Q}$ ,  $h$ , and  $\varepsilon$ , in both the in-phase and the out-of-phase cases. We emphasize that ion fluctuation and correlation effects neglected in the mean-field approach we adopt, become important at high values of  $g \gg 1$ , *i.e.*, in the strong coupling regime [13]. Finally we note, that qualitatively similar conclusions with respect to the enhancement of the contact counterion density have recently been obtained by numerical solutions of the integral equation theory [11]; however some of their effects are related to the fact that both the counterions and the surface charges have an excluded volume in [11].

*Pressure between charge modulated surfaces-* In this section, we show why surface charge modulations in the in-phase case (under certain conditions), and in the out-of-phase case (for all values of the parameters) *always reduce* the repulsive pressure between surfaces with counterions compared with the case of uniformly charged surfaces.

The free energy,  $F$ , of the system has the form [17]:  $\frac{F}{k_B T} = \int n(\vec{r}) [\ln(n(\vec{r})v_0) - 1] d\vec{r} + \frac{1}{2} \int \phi(\vec{r}) n_{tot}(\vec{r}) d\vec{r}$ , where the total system charge density:  $n_{tot}(\vec{r}) = n(\vec{r}) - \frac{1}{2} [\sigma_1(\vec{\rho})\delta(z + \frac{h}{2}) + \sigma_2(\vec{\rho})\delta(z - \frac{h}{2})]$ ,  $v_0$  is an ion-size volume, and  $\vec{r} = (\vec{\rho}, z)$ . Keeping terms up to second order in the amplitude of surface charge modulations,  $\varepsilon(\vec{Q})$ , one notes that  $F$  can be expressed only in terms of the  $x-y$  average of the potentials as obtained above,  $\langle \phi_2(\vec{r}) \rangle_{\vec{\rho}}$ ,  $\langle \phi_1^2(\vec{r}) \rangle_{\vec{\rho}}$  and  $\langle \sigma_i(\vec{\rho}) \phi_1(\vec{r}) \rangle_{\vec{\rho}}$ ; after straightforward but lengthy integrations, one obtains the following dimensionless free energy,  $f \equiv \frac{F}{k_B T A_0 (\sigma_0/Z)}$ , per counterion  $f_+$  and  $f_-$  in the in-phase and in the out-of-phase case, respectively:

$$f_{\pm} = f_0 + \sum_{\vec{Q}} 2|\varepsilon(\vec{Q})|^2 \frac{1 + \mathcal{Q} \pm e^{-\mathcal{Q}h} (\mathcal{Q} - 1)}{1 + 2\mathcal{Q} + 2\mathcal{Q}^2 \mp e^{-\mathcal{Q}h} + (1 \mp e^{-\mathcal{Q}h}) k_0^2}, \quad (3)$$

where  $f_0$  is the free energy of uniformly charged surfaces with counterions [17]. The normalized, dimensionless osmotic pressure between the surfaces,  $p = -\partial f / \partial h$ , is the main quantity of interest measured in experiments. Our principle observations follow intuitively from the asymptotic analysis of the pressure: (i) in the *in-phase* case, in the limit of small separations,  $h \ll 1$ ,  $\mathcal{Q}h \ll 1$ , one has,

$$p - p_0 = \sum_{\vec{Q}} |\varepsilon(\vec{Q})|^2 \frac{\mathcal{Q}^2 - \frac{4}{3}}{(2 + \mathcal{Q})^2} + \mathcal{O}(h), \quad (4)$$

where  $p_0 = k_0^2$  is the normalized, dimensionless pressure [20] between uniformly charged surfaces [17]. In the *out-of-phase* case, when  $h \ll 1$ ,  $\mathcal{Q}h \ll 1$  we obtain,  $p - p_0 = -\sum_{\vec{Q}} |\varepsilon(\vec{Q})|^2 + \mathcal{O}(h)$ . Therefore in this asymptotic limit of small inter-surface separations, in the in-phase case the pressure between the surfaces is reduced for long-wavelength surface charge modulations; in the out-of-phase case, the pressure is reduced for any wavelength. This effect [quadratic in  $\varepsilon(\vec{Q})$ ] is related to the reduction of the midplane counterion density induced by surface charge modulations upon decreasing of  $\mathcal{Q}$ . In the limit of large inter-surface separations,  $h \gg 1$ ,  $\mathcal{Q}h \gg 1$ , in both cases, (i) and (ii), the pressure is *always reduced*:

$$p - p_0 = -\sum_{\vec{Q}} \frac{4|\varepsilon(\vec{Q})|^2 (1 + \mathcal{Q})}{(1 + 2\mathcal{Q} + 2\mathcal{Q}^2)^2} \frac{\pi^2}{h^3} + \mathcal{O}\left(\frac{1}{h^4}\right). \quad (5)$$

We note that this asymptotic form is similar to the universal, Casimir-type attractive fluctuation pressure,  $\sim -1/h^3$  (see *e.g.*, [21,22]) between the surfaces with boundary conditions that



suppress or modify fluctuations of the medium; the amplitude of this pressure is non-universal, of course, since the spectrum of modes is restricted by the fixed surface charge modulations in the present case.

It is interesting to note, that corresponding asymptotic results for the pressure [20] between *uniformly* charged surfaces [17],  $p_0$  [ $p_0 = \frac{2}{h}$ , if  $h \ll 1$ , and  $p_0 = \frac{\pi^2}{h^2}$ , if  $h \gg 1$ ] show that the modulation correction,  $p - p_0$ , can *never* overcome the overall repulsion in these limiting cases. For intermediate values of  $h$ , however, the total pressure,  $p$ , can change sign and become attractive, even in the in-phase case, if we extrapolate our perturbative results to values of  $\varepsilon(\vec{Q})$ , which are not necessarily small. We also emphasize again that we use a mean-field approach, and neglect the *fluctuation* contribution to the pressure [21] that further reduces the repulsive interaction.

The generic behavior of the pressure difference,  $p - p_0$ , in the in-phase case for arbitrary  $h$  and  $Q$  is shown in the inset of Fig. 1. There is a broad range of parameters where the repulsion between two surfaces is reduced by the effect of surface charge modulations. It is interesting to note, that already at  $h \gtrsim 1.25$ , charge modulations *reduce* the pressure for any value of  $Q$ . We have checked that a qualitatively similar reduction of the repulsive pressure occurs in the case of a square lattice of discrete charges (infinite sum with respect to all  $\vec{Q}$  modes).

Our prediction of the *reduction* of pressure in the in-phase case differs from the conclusions of Refs. [2, 3, 15]. The difference between those works and our predictions can be understood using an alternative definition of the pressure via the stress tensor [23]:  $p = [\tilde{n} + \frac{1}{4} \langle \phi_x^2 + \phi_y^2 - \phi_z^2 \rangle_{\vec{p}}]_{\vec{z}=0}$ , where  $\tilde{n} \equiv 2\pi\ell_B Z^2 \lambda_0^2 \langle n \rangle_{\vec{p}}$  is the normalized, dimensionless density [20]; this gives results identical to the ones presented above. In the in-phase case,  $\phi_z = 0$ , as implied by the symmetry, and the electrostatic contribution to the pressure is purely repulsive, arising from terms in the electric field that are *first order* in the amplitude of surface charge modulations,  $\varepsilon(\vec{Q})$  [the pressure is, of course, always quadratic in  $\varepsilon(\vec{Q})$ , since  $p \sim \phi_x^2 + \phi_y^2$ ]. This is the only contribution that was taken into account in Refs. [2, 3, 15], where only first order corrections to the counterion density were calculated. As far as the osmotic term is concerned, the first order corrections vanish when the average over the  $x-y$  plane is performed, and hence, it is necessary to calculate the counterion density to *second order* in the surface charge modulation amplitude,  $\varepsilon(\vec{Q})$ . It is this osmotic contribution, calculated here, that is the origin of the *universal reduction* of the pressure, since  $\tilde{n}$  at the midplane is always reduced by surface charge modulations, as we have shown; this effect is quadratic in  $\varepsilon(\vec{Q})$ , and beyond the linear approach of Refs. [2, 3, 15]. It is the competition between the osmotic and electrostatic contribution that leads to the diagram represented in the inset of Fig. 1. In the out-of-phase case, where  $\phi_z \neq 0$ , the electrostatic contribution to  $p$  is reduced compared with the in-phase case, and further reduction of the midplane density by charge modulations *always* leads to the reduction of the pressure, compared with the case of uniformly charged surfaces.

In summary, we emphasize that our main result, the prediction of a *universal* reduction of the osmotic pressure by long-wavelength, quenched surface charge modulations, is naturally related to effects of fluctuating surface charges on the pressure. The unifying theme is that systems with thermally *fluctuating* surface charge density are qualitatively similar to systems with quenched, *modulated* surface charge density: in both situations, the counterions are more localized near the surfaces compared with the case of uniformly charged surfaces, and in both cases this leads to a reduction of the pressure.

\* \* \*



We are grateful to P. Pincus, C. Jeppesen, J.-P. Hansen, A.W.C. Lau, R. Netz, and T.O. White. We acknowledge the support from Israel Binational Science Foundation (BSF) grant 98-00063, and the support of the Schmidt Minerva Center.

*Appendix-* In this appendix, we summarize the results for perturbative corrections,  $\phi_1(\vec{\rho}, \tilde{z})$ , and  $\langle \phi_2(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}}$ , to the potential,  $\phi_0(\tilde{z})$ , induced by surface charge modulations.

In particular, one finds:  $\phi_1(\vec{Q}, \tilde{z}) = \mathcal{A} v_1(Q, \tilde{z}) + \mathcal{B} v_2(Q, \tilde{z})$ , where  $\mathcal{A} = \frac{2\varepsilon(Q)}{Q} \frac{\alpha_1 + \alpha_2 e^{i\vec{Q} \cdot \vec{b}}}{\alpha_1^2 - \alpha_2^2}$ ,  $\mathcal{B} = \frac{2\varepsilon(Q)}{Q} \frac{\alpha_1 e^{i\vec{Q} \cdot \vec{b}} + \alpha_2}{\alpha_1^2 - \alpha_2^2}$ ,  $\alpha_1 = \exp(-\frac{Qh}{2})(k_0^2 + 1)/Q^2$ ,  $\alpha_2 = \exp(\frac{Qh}{2})(k_0^2 + 1 + 2Q + 2Q^2)/Q^2$ , and  $v_1(Q, \tilde{z}) = e^{Q\tilde{z}}(1 + (k_0/Q) \tan k_0 \tilde{z})$ ,  $v_2(Q, \tilde{z}) = e^{-Q\tilde{z}}(1 - (k_0/Q) \tan k_0 \tilde{z})$ .

The average potential,  $\langle \phi_2(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}}$  is also obtained in a simple, closed form in terms of elementary functions:  $\langle \phi_2(\vec{\rho}, \tilde{z}) \rangle_{\vec{\rho}} = \sum_{\vec{Q}} \frac{16|\varepsilon(\vec{Q})|^2 \Gamma_{\pm}(Q)}{Q^2 (\alpha_1 \mp \alpha_2)^2}$ , where  $\Gamma_+(Q)$  and  $\Gamma_-(Q)$  corresponds to the in-phase case and the out-of-phase case, respectively. Here we use the notations:  $\Gamma_{\pm}(Q) = u_2(\tilde{z})[\frac{1}{\eta} \mathcal{G}_{\pm}(\frac{k_0 h}{2}) - \mathcal{F}_{\pm}(\frac{k_0 h}{2}) + \mathcal{F}_{\pm}(k_0 \tilde{z})] - u_1(\tilde{z}) \mathcal{G}_{\pm}(k_0 \tilde{z})$ ,  $u_1(\tilde{z}) = \tan k_0 \tilde{z}$ ,  $u_2(\tilde{z}) = 1 + k_0 \tilde{z} \tan k_0 \tilde{z}$ ,  $\eta = \frac{k_0}{1+k_0} + \frac{k_0 h}{2}$ ,  $\mathcal{G}_{\pm}(x) = \frac{\cosh 2\theta x \mp 1}{8\theta^2 \cos^2 x} [\tan x - x(1 - \tan^2 x)] + \frac{\sinh 2\theta x}{4\theta \cos^2 x} [1 + x \tan x] \pm \frac{1}{4} [\frac{x}{\cos^2 x} + \tan x]$ , and  $\mathcal{F}_{\pm}(x) = \frac{\cosh 2\theta x \mp 1}{8\theta^2} \frac{\tan^4 x}{\cos^2 x} + \frac{\sinh 2\theta x \tan x}{4\theta \cos^2 x} \pm \frac{1}{4 \cos^2 x} - \frac{\cosh 2\theta x}{8\theta^2}$ , where  $\theta = Q/k_0$ .

## REFERENCES

- [1] J. N. Israelachvili, *Intermolecular and Surface Forces* (Academic Press Inc., San Diego, 1992).
- [2] P. Richmond, J. Chem. Soc. Faraday Trans. II **71**, 1154 (1975).
- [3] M. Kostoglou and A. J. Karabelas, J. Colloid Interface Sci. **151**, 534 (1992).
- [4] D. Y. Chan, D. J., Mitchel, and B. W. Ninham, J. Chem. Phys. **72**, 5159 (1980).
- [5] R. M. Peitzsch, M. Eisenberg, K. A. Sharp, and S. McLaughlin, Biophys. J. **68**, 729 (1995).
- [6] R. Kjellander and S. Marcelja, J. Chem. Phys. **88**, 7138 (1988).
- [7] W. van Megen and I. Snook, J. Chem. Phys. **73**, 4656 (1980).
- [8] H. Wennerstrom and B. Jonsson, J. Phys. France **49**, 1033 (1988).
- [9] G. D. Guttman and D. Andelman, J. Phys. France II **3**, 1411 (1993).
- [10] L. I. Daikhin and M. Urbakh, Phys. Rev. E **59**, 1921 (1999).
- [11] O. Gonzalez-Amezcu, M. Hernandez-Contreras, and P. Pincus, Phys. Rev. E **64**, 041603 (2001).
- [12] R. Messina, C. Holm, and K. Kremer, Eur. Phys. J. E, **4**, 363 (2001); R. Messina, C. Holm, and K. Kremer, Phys. Rev. E, **64**, 021405 (2001).
- [13] A. G. Moreira and R. R. Netz, Europhys. Lett., **57**, 911 (2002); see also Europhys. Lett. **52**, 705 (2000); Phys. Rev. Lett. **87**, 078301 (2001).
- [14] D. B. Lukatsky, S. A. Safran, A. W. C. Lau, and P. Pincus, Europhys. Lett., **58**, 785 (2002).
- [15] T. O. White and J.-P. Hansen, J. Phys. Cond. Matt., **14**, 7649 (2002).
- [16] The case where the counterions are located only in the space between the surfaces is chosen for convenience to enable comparison with MC simulations. The case where the counterions are located in the whole space leads to similar qualitative conclusions.
- [17] S. A. Safran, *Statistical Thermodynamics of Surfaces, Interfaces, and Membranes* (Addison-Wesley Publishing Com., Reading, MA 1994).
- [18] D. B. Lukatsky, PhD thesis, Weizmann Institute, work in preparation.
- [19] D. M. Heyes, M. Barber, and J. H. R. Clarke, J. Chem. Soc., Faraday Trans. II, **73**, 1485 (1977).
- [20] We note again that the normalized, dimensionless pressure,  $p$ , is related to the physical pressure,  $\Pi$ , by:  $p \equiv \frac{\Pi}{k_B T 2\pi \ell_B \sigma_0^2}$ . In the case of uniformly charged surfaces [17],  $\Pi_0 = k_B T n_0$ , and, consequently,  $p_0 = k_0^2$ , where  $k_0$  [defined in the text after Eq. (1)] is dimensionless [since we express the lengths in units of  $\lambda_0$ , as specified in the text after Eq. (2)].
- [21] P. Pincus and S. A. Safran, Europhys. Lett. **42**, 103 (1998).



- [22] M. Kardar and R. Golestanian, Rev. Mod. Phys. **71**, 1233 (1999).
- [23] W. B. Russel, D. A. Saville, and W. R. Schowalter, *Colloidal Dispersions* (Cambridge University Press, Cambridge 1989).