

Spin-Wave Description of Haldane-gap Antiferromagnets

Shoji Yamamoto and Hiroshi Hori

Division of Physics, Hokkaido University, Sapporo 060-0810, Japan

(Received 13 December 2002)

Modifying the conventional antiferromagnetic spin-wave theory which is plagued by the difficulty of the zero-field sublattice magnetizations diverging in one dimension, we describe magnetic properties of Haldane-gap antiferromagnets. The modified spin waves, constituting a grand canonical bosonic ensemble so as to recover the sublattice symmetry, not only depict well the ground-state correlations but also give useful information on the finite-temperature properties.

PACS numbers: 75.10.Jm, 05.30.Jp, 75.40.Mg

Haldane's conjecture [1,2] that one-dimensional Heisenberg antiferromagnets should exhibit qualitatively different low-energy structures based on whether the constituent spins are integral or fractional sparked renewed interest in the field of quantum magnetism. An energy gap immediately above the ground state was indeed observed in a quasi-one-dimensional Heisenberg antiferromagnet $\text{Ni}(\text{C}_2\text{H}_8\text{N}_2)_2\text{NO}_2$ (CDO) [3] and a rigorous example of such a massive phase was also found out [4,5]. The energy gaps in magnetic excitation spectra, that is, spin gaps, are now one of the most attractive and important topics. In the context of theoretical progress, we may be reminded of quantized plateaux in the ground-state magnetization curves [6], a dramatic crossover from one- to two-dimensional quantum antiferromagnets [7], and an antiferromagnetic excitation gap accompanied by ferromagnetic background [8-11]. From the experimental point of view, metal oxides such as spin-Peierls compounds $\text{Cu}_{1-x}\text{M}_x\text{GeO}_3$ ($\text{M} = \text{Zn}; \text{Mg}$) [12,13], Haldane-gap antiferromagnets R_2BaNdO_5 ($\text{R} = \text{rare earth}$) [14] and ladder materials $\text{Sr}_{n-1}\text{Cu}_{n+1}\text{O}_{2n}$ ($n = 3; 5; 7$; [15] have significantly contributed to systematic investigations of the mechanism of gap formation.

In studying Haldane-gap antiferromagnets, numerical tools such as quantum Monte Carlo and density-matrix renormalization-group methods are indeed useful, but analytic approaches still play an important role. The nonlinear model technique [12,16-19] is powerful enough to investigate low-energy structures with particular emphasis on the competition between massive and massless phases. It is the valence-bond-solid description [4,5] of integer-spin chains that enables us to readily understand the hidden order [20] inherent in the Haldane massive phase. However, these methods are not so useful for exploring their all properties as they are in investigating the ground-state properties. Although we may consider applying the spin-wave scheme to Haldane-gap antiferromagnets in such circumstances, the difficulty of the zero-field sublattice magnetizations diverging in one dimension has been a problem in the conventional spin-wave theory [21,22] for years. Therefore, it was a major breakthrough that spin waves, being constrained to keep zero magnetization, succeeded in precisely describing the low-temperature thermodynamics of

the one-dimensional Heisenberg ferromagnet [23]. This new spin-wave scheme, which is now referred to as the modified spin-wave theory, was further applied to quantum antiferromagnets and ferrimagnets. The modified spin-wave scheme is highly successful for extensive ferrimagnets [24-29] and still applies well for two-dimensional antiferromagnets [30-34]. As for one-dimensional antiferromagnets, there exists a pioneering argument [35], but it looks unsatisfactory for interpreting experimental and numerical observations.

In this article, we aim to do away with our vague but persistent impression that the spin-wave scheme hardly works for one-dimensional quantum antiferromagnets. Spin-2 Haldane-gap antiferromagnets [36,37], as well as those of spin 1, have recently been synthesized and more explorations into novel quantum phenomena in one dimension are expected in the future. In such circumstances, we make our first attempt to construct a modified spin-wave theory for Haldane-gap antiferromagnets.

We consider integer-spin antiferromagnetic Heisenberg chains

$$H = J \sum_{j=1}^{X^L} \mathbf{S}_j \cdot \mathbf{S}_{j+1}; \quad \mathbf{S}_j^2 = S(S+1); \quad (1)$$

We define bosonic operators for the spin deviation in each sublattice via

$$\begin{aligned} \mathbf{S}_{2n-1}^+ &= \frac{1}{q} \sum_{\mathbf{k}} 2S \frac{\mathbf{a}_{\mathbf{k}}^\dagger \mathbf{a}_n}{\mathbf{a}_{\mathbf{k}}^\dagger \mathbf{a}_n}; \quad \mathbf{S}_{2n-1}^z = S - \mathbf{a}_{\mathbf{k}}^\dagger \mathbf{a}_n; \\ \mathbf{S}_{2n}^+ &= \mathbf{b}_n^\dagger \sum_{\mathbf{k}} 2S \frac{\mathbf{b}_{\mathbf{k}}^\dagger}{\mathbf{b}_{\mathbf{k}}^\dagger}; \quad \mathbf{S}_{2n}^z = S + \mathbf{b}_{\mathbf{k}}^\dagger \mathbf{b}_n; \end{aligned} \quad (2)$$

The Fourier-transformed operators are introduced as

$$\begin{aligned} \mathbf{a}_{\mathbf{k}} &= \frac{1}{N} \sum_{n=1}^{X^N} e^{ik(n-1/4)} \mathbf{a}_n; \\ \mathbf{b}_{\mathbf{k}} &= \frac{1}{N} \sum_{n=1}^{X^N} e^{ik(n+1/4)} \mathbf{b}_n; \end{aligned} \quad (3)$$

where twice the lattice constant $2a$ is set equal to unity and therefore $k = 2\pi n/N$ ($n = 0; 1; \dots; N-1; N=L=2$). Through the Bogoliubov transformation

$$\begin{aligned} \mathbf{a}_{\mathbf{k}}^\dagger &= \frac{\gamma_{\mathbf{k}}}{\cosh k} \cosh k + \frac{k}{\sinh k} \sinh k; \\ \mathbf{b}_{\mathbf{k}}^\dagger &= \frac{\gamma_{\mathbf{k}}}{\cosh k} \cosh k - \frac{k}{\sinh k} \sinh k; \end{aligned} \quad (4)$$

we obtain

$$H = E_N + E_1 + E_0 + H_1 + H_0 + O(S^{-1}); \quad (5)$$

where

$$E_N = 2NJS^2; \quad (6a)$$

$$E_1 = 4NJS(\quad); \quad (6b)$$

$$E_0 = 2NJ(\quad)^2; \quad (6c)$$

$$H_1 = J \sum_k \left[\frac{1}{2} (k) \frac{y}{k} k + \frac{y}{k} k \right. \\ \left. + \frac{1}{2} (k) \frac{y}{k} k + \frac{y}{k} \frac{y}{k} \right]; \quad (7a)$$

$$H_0 = J \sum_k \left[\frac{1}{2} (k) \frac{y}{k} k + \frac{y}{k} k \right. \\ \left. + \frac{1}{2} (k) \frac{y}{k} k + \frac{y}{k} \frac{y}{k} \right]; \quad (7b)$$

with

$$= \frac{1}{2N} \sum_k \cos \frac{k}{2} \sinh 2k; \quad (8a)$$

$$= \frac{1}{2N} \sum_k (\cosh 2k - 1); \quad (8b)$$

$$\frac{1}{2} (k) = 2S \cosh 2k \cos \frac{k}{2} \sinh 2k; \quad (9a)$$

$$\frac{1}{2} (k) = 2(\quad) \cosh 2k \cos \frac{k}{2} \sinh 2k; \quad (9b)$$

$$\frac{1}{2} (k) = 2S \cos \frac{k}{2} \cosh 2k \sinh 2k; \quad (10a)$$

$$\frac{1}{2} (k) = 2(\quad) \cos \frac{k}{2} \cosh 2k \sinh 2k; \quad (10b)$$

The naivest diagonalization of the Hamiltonian (5), whether up to $O(S^1)$ or up to $O(S^0)$, results in diverging sublattice magnetizations even at zero temperature. Although eq. (2) assumes that $a_n^\dagger a_n = 2S$ and $b_n^\dagger b_n = 2S$, the conventional spin-wave theory cannot reasonably control the boson numbers. Then we consider introducing a grand canonical constraint to the noncompact Hamiltonian (5).

Isotropic magnets should lie in the state of zero magnetization $\sum_j S_j^z = 0$ and the minimization of the free energy under such a condition indeed yields an excellent description of the low-temperature thermodynamics of ferromagnets [23]. However, in the cases with antiferromagnetic exchange interactions, the zero-magnetization constraint, claiming that $\sum_n (a_n^\dagger a_n - b_n^\dagger b_n) = 0$, still fails to overcome the divergence of the numbers of the sublattice bosons. Hence we minimize the free energy constraining the sublattice magnetizations to be zero:

$$\sum_n a_n^\dagger a_n = \sum_n b_n^\dagger b_n = SN; \quad (11)$$

Within the conventional spin-wave theory, spins on one sublattice point predominantly up, while those on the other predominantly down. This condition (11) restores the sublattice symmetry. In order to enforce the constraint (11), we first introduce a Lagrange multiplier and diagonalize

$$\mathbb{H} = H + 2J \sum_k (a_k^\dagger a_k + b_k^\dagger b_k); \quad (12)$$

Then the ground-state energy and the dispersion relation are obtained as

$$E_g = E_N + \mathbb{E}_1; \quad \mathbb{E}_1 = E_1 + 4NJ; \quad (13)$$

$$\frac{1}{2} (k) = \frac{1}{2} (k); \quad \frac{1}{2} (k) = \frac{1}{2} (k) + 2 \cosh 2k; \quad (14)$$

within the linear modified spin-wave scheme and as

$$E_g = E_N + \mathbb{E}_1 + E_0; \quad (15)$$

$$\frac{1}{2} (k) = \frac{1}{2} (k) + \frac{1}{2} (k); \quad (16)$$

within the up-to- $O(S^0)$ interacting modified spin-wave scheme. For eqs. (13) and (14), $\frac{1}{2} (k)$ is given by $\frac{1}{2} (k) = 2 \sinh 2k \cosh 2k \cos \frac{k}{2} = 0$, whereas for eqs. (15) and (16), $\frac{1}{2} (k)$ may be determined in two ways. One idea is the perturbational treatment of H_0 , which is referred to as the perturbational interacting modified spin-wave scheme, where $\frac{1}{2} (k)$ is still given by $\frac{1}{2} (k) = 0$ and the $O(S^0)$ quantum correction is the $O(S^1)$ -eigenstate average of H_0 . The other is the full diagonalization of $H_1 + H_0$, which is referred to as the full-diagonalization interacting modified spin-wave scheme, where $\frac{1}{2} (k)$ is given by $\frac{1}{2} (k) + \frac{1}{2} (k) = 0$. Once $\frac{1}{2} (k)$ is given, we calculate the free energy and obtain the optimum thermal distribution functions as

$$\langle a_k^\dagger a_k \rangle = \langle b_k^\dagger b_k \rangle = n_k = \frac{1}{e^{J \frac{1}{2} (k) / k_B T} - 1}; \quad (17)$$

where $\frac{1}{2} (k)$ is self-consistently determined by the condition

$$\sum_k (2n_k + 1) \cosh 2k = (2S + 1)N; \quad (18)$$

First, let us evaluate the ground-state energy. We compare the modified spin-wave calculations with the highly accurate quantum Monte Carlo estimates [38] in Table I. The modified spin-wave findings are generally in good agreement with the quantum Monte Carlo results. The interacting modified spin waves describe the ground-state correlations much better than the linear ones and their description becomes increasingly refined with increasing S . The interacting modified spin-wave findings miss the correct value by 0.5 percent for $S = 1$ and by only 0.008 percent for $S = 3$.

Secondly, we consider the Haldane gap Δ ($ak = \pi$)

(I). Table II shows that the modified spin-wave scheme, in contrast with the conventional spin-wave theory, succeeds in generating the gap but considerably underestimates it. Unavailability of the absolute energy

level structure is an inevitable consequence of our employing the effective Hamiltonian (12). Then we examine the present scheme by scaling $\Delta(T)$ to its zero-temperature value $\Delta(T=0) = \Delta_0$. Such an argument is quite usual with field-theoretical investigations, which do not lead to an estimate of the normalization factor but derive finite-temperature expressions involving only ratios such as $k_B T = \Delta_0$. In Fig. 1, we compare the modified spin-wave calculations with numerical [39], field-theoretical [40] and experimental [41] findings. Now the superiority of the modified spin-wave scheme is clear at a glance. Of all the theoretical tools, the full-diagonalization interacting modified spin-wave approach is the most successful to reproduce the observed upward behavior of $\Delta(T)$ with increasing temperature [41,43]. The nonlinear ϕ -model treatment [40,44] is justified well in the low-temperature region $k_B T \ll \Delta_0$, while the maximum-entropy technique [39] works less well with increasing temperature. Nickel compounds such as Y_2BaNiO_5 [41] and $\text{Ni}(\text{C}_2\text{H}_8\text{N}_2)_2\text{NO}_2$ (CD_4) [42] are good candidates for spin-1 Haldane-gap antiferromagnets, but magnetic anisotropy and interchain interaction are not negligible there. They split the lowest excitation gap of the isotropic chain (1) into several levels and smear the intrinsic behavior of the ideal integer-spin chains. Considering such practical factors, the present theory satisfactorily interprets the observations.

Lastly, we show the modified spin-wave calculations of the magnetic susceptibility in Fig. 2. Considering the total breakdown of the conventional spin-wave theory in one-dimensional thermodynamic calculations, the modified spin-wave achievement is highly successful. All the calculations are guaranteed to reproduce the paramagnetic susceptibility $\chi = Lg^2 \frac{S(S+1)}{2} = S(S+1)/3k_B T$ at high temperatures. Since the interacting modified spin-wave scheme gives better estimates of the gap than the linear modified spin-wave one (Table I), it is somewhat better at describing the low-temperature behavior. With increasing S , the activation-type initial behavior is suppressed and the antiferromagnetic peak is broadened. The $S = 1$ modified spin-wave calculations are in agreement with the quantum Monte Carlo findings over a wide temperature range, while those for $S = 2$ look somewhat poorer at intermediate temperatures. It may be closely related to the fact that the modified spin-wave estimates of the $S = 2$ gap are worse than those of the $S = 1$ gap (Table II). However, Table II suggests that the validity of the modified spin-wave scheme for excitations significantly improves with increasing S , possibly in a staggered way. Thermodynamic calculations, whether by quantum Monte Carlo or density-matrix renormalization group, for systems with large degrees of freedom are less feasible numerically, in particular, at low temperatures. The present scheme has the advantage of saving time and computational resources.

We have demonstrated the applicability of the new spin-wave scheme to Haldane-gap antiferromagnets. This is the first comprehensive attempt to describe one-

dimensional spin-gapped antiferromagnets in terms of spin waves. The modified spin waves are free from their thermal as well as quantum divergence and can therefore microscopically interpret various magnetic properties. Besides the magnetic susceptibility, the spin correlation function and the nuclear spin-lattice relaxation time can be revealed. Unfortunately, we have less information on the bare energy spectrum, because we get rid of the quantum divergence at the cost of the original Hamiltonian. However, with the zero-temperature spin gap, which is readily and precisely available through numerical calculations within a canonical ensemble [38], we can still argue the energy structure quantitatively.




FIG. 1. Temperature dependences of the spin-1 Haldane gap calculated by the linear (LMSW), perturbational interacting (PIMSW) and full-diagonalization interacting (FDIMSW) modified spin-wave schemes, a quantum Monte Carlo (QMC) method combined with the maximum-entropy technique [39] and the nonlinear ϕ -model (NSM) approach [40]. Inelastic neutron-scattering measurements (Exp.) on Y_2BaNiO_5 [41] are also shown for reference.




FIG. 2. The linear (LMSW), perturbational interacting (PIMSW) and full-diagonalization interacting (FDIMSW) modified spin-wave calculations of the zero-field magnetic susceptibility as a function of temperature compared with quantum Monte Carlo (QMC) estimates.

Haldane-gap antiferromagnets in a magnetic field provide further interesting issues. With the increase of an applied field, the gap is reduced and the ground state is mixed increasingly with the first excited state. Indeed the nuclear spin-lattice relaxation is accelerated in the vicinity of the critical field, but its overall behavior as a function of an applied field and temperature [45] is far from understandable at a glance. We may expect the present new scheme to open the way for the total understanding of low-dimensional spin-gapped antiferromagnets.

[1] F.D.M. Haldane: Phys. Lett. 93A (1983) 464.
[2] F.D.M. Haldane: Phys. Rev. Lett. 50 (1983) 1153.
[3] J.P. Renard, M. Verdaguer, L.P. Regnault, W.A.C. Erkelens, J. Rossat-Mignod and W.G. Stirling: Europhys. Lett. 3 (1987) 945.
[4] I. A. A. eck, T. Kennedy, E.H. Lieb and H. Tasaki: Phys. Rev. Lett. 59 (1987) 799.
[5] I. A. A. eck, T. Kennedy, E.H. Lieb and H. Tasaki: Commun. Math. Phys. 115 (1988) 477.
[6] M. Oshikawa, M. Yamanka and I. A. A. eck: Phys. Rev. Lett. 78 (1997) 1984.
[7] E. Dagotto and T.M. Rice: Science 271 (1996) 618.
[8] S. Maslov and A. Zheludev: Phys. Rev. B 57 (1998) 68.
[9] S. Yamamoto and T. Fukui: Phys. Rev. B 57 (1998) R14008.
[10] S. Yamamoto, T. Fukui, K. M. Aisinger and U. Schollwöck: J. Phys.: Condens. Matter 10 (1998) 11033.
[11] Y. Takushima, A. Koga and N. Kawakami: Phys. Rev. B 61 (2000) 15189.
[12] M. Hase, I. Terasaki and K. Uchinokura: Phys. Rev. Lett. 70 (1993) 3651.
[13] M. Hase, I. Terasaki, Y. Sasago, K. Uchinokura and H. Obara: Phys. Rev. Lett. 71 (1993) 4059.
[14] J. D. Armitage and L.P. Regnault: Solid State Commun. 86 (1993) 409.
[15] M. Azuma, Z. Hiroi, M. Takano, K. Ishida and Y. Kitaoka: Phys. Rev. Lett. 73 (1994) 3463.
[16] I. A. A. eck: Nucl. Phys. B 257 (1985) 397.
[17] I. A. A. eck: Nucl. Phys. B 265 (1985) 409.
[18] T. Fukui and N. Kawakami: Phys. Rev. B 55 (1997) R14709.
[19] T. Fukui and N. Kawakami: Phys. Rev. B 56 (1997) 8799.
[20] M. den Nijs and K. Rommelse, Phys. Rev. B 40 (1989) 4709.
[21] P.W. Anderson: Phys. Rev. 86 (1952) 694.
[22] R. Kubo: Phys. Rev. 87 (1952) 568.
[23] M. Takahashi: Phys. Rev. Lett. 58 (1987) 168.
[24] S. Yamamoto, T. Fukui and T. Sakai: Eur. Phys. J. B 15 (2000) 211.
[25] S. Yamamoto: Solid State Commun. 117 (2001) 1.
[26] T. Nakanishi, S. Yamamoto and T. Sakai: J. Phys. Soc. Jpn. 70 (2001) 1380.
[27] T. Nakanishi and S. Yamamoto: Phys. Rev. B 65 (2002) 214418.
[28] S. Yamamoto and T. Nakanishi: Phys. Rev. Lett. 89

(2002) 157603.
[29] A.S. Ovchinnikov, I.G. Bostrem, V.E. Sinitsyn, A.S. Boyarchenkov, N.V. Baranov and K. Inoue: J. Phys.: Condens. Matter 14 (2002) 8067.
[30] M. Takahashi: Phys. Rev. B 40 (1989) 2494.
[31] J.E. Hirsch and S. Tang: Phys. Rev. B 40 (1989) 4769.
[32] S. Tang, M.E. Lazzouni and J.E. Hirsch: Phys. Rev. B 40 (1989) 5000.
[33] H.A. Ceccatto, C.J. Gazza and A.E. Trumper: Phys. Rev. B 45 (1992) 7832.
[34] A.V. Dotsenko and O.P. Sushkov: Phys. Rev. B 50 (1994) 13821.
[35] S.M. Rezende: Phys. Rev. B 42 (1990) 2589.
[36] G.E. Granroth, M.W. Meisel, M. Chaparala, Th. Jolicur, B.H. Ward and D.R. Talham: Phys. Rev. Lett. 77 (1996) 1616.
[37] H. Yamazaki and K. Katsumata: Phys. Rev. B 54 (1996) R6831.
[38] S. Todo and K. Kato: Phys. Rev. Lett. 87 (2001) 047203.
[39] J. Deisz, M. Jarrell and D.L. Cox: Phys. Rev. B 48 (1993) 10227.
[40] Th. Jolicur and O. Golinelli: Phys. Rev. B 50 (1994) 9265.
[41] T. Sakaguchi, K. Kakurai, T. Yokoo and J. Akimitsu: J. Phys. Soc. Jpn. 65 (1996) 3025.
[42] J.P. Renard, M. Verdaguer, L.P. Regnault, W.A.C. Erkelens, J. Rossat-Mignod, J. Ribas, W.G. Stirling and C. Vettier: J. Appl. Phys. 63 (1988) 3538.
[43] Z. Tun, W.J.L. Buyers, R.L. Armstrong, K. Kakurai and B. Briat: Phys. Rev. B 42 (1990) 4677.
[44] I. A. A. eck: Phys. Rev. Lett. 62 (1989) 474.
[45] N. Fujiwara, T. Goto, S. Maegawa and T. Kohmoto: Phys. Rev. B 47 (1993) 11860.

TABLE I. The linear (LM SW), perturbational interacting (PIM SW) and full-diagonalization interacting (FDIM SW) modified spin-wave calculations of the ground-state energy per site compared with quantum Monte Carlo (QMC) estimates [38].

	S = 1	S = 2	S = 3
LM SW	1:361879	4:726749	10:0901
PIM SW	1:394853	4:759760	10:1231
FDIM SW	1:394617	4:759759	10:1231
QMC	1:401481 (4)	4:761249 (6)	10:1239 (1)

TABLE II. The linear (LM SW), perturbational interacting (PIM SW) and full-diagonalization interacting (FDIM SW) modified spin-wave calculations of the lowest excitation gap ($T = 0$) compared with quantum Monte Carlo (QMC) estimates [38].

	S = 1	S = 2	S = 3
LM SW	0:07200	0:00626	0:00279
PIM SW	0:07853	0:00655	0:00287
FDIM SW	0:08507	0:00683	0:00295
QMC	0:41048 (6)	0:08917 (4)	0:01002 (3)