

Quantum Orders and Spin Liquids in Cs_2CuCl_4

Yi Zhou^{1,*} and Xiao-Gang Wen^{2,†}

¹*Center for Advanced Study, Tsinghua University, Beijing 100084, P. R. China*

²*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

(Dated: March 22, 2022)

Motivated by experiments on Cs_2CuCl_4 samples, we studied and classified the symmetric spin liquids on triangular lattice. We identified 63 Z_2 spin liquids, 30 $U(1)$ spin liquids and 2 $SU(2)$ spin liquids. All those spin liquids have the same symmetry but different quantum orders. We calculated the spin spectral functions in some simple spin liquids and compared them to the one measured by experiments on Cs_2CuCl_4 . We find that the $U(1)C\tau_+^0\tau_-^0\tau^1$ spin liquid or one of its relatives is consistent with observed properties of the spin liquid state in Cs_2CuCl_4 . We discussed the fine distinctions among those possible spin liquids and the spin liquids proposed using slave-fermion approach, so that future experiments can determine which spin liquids actually describe the spin state in Cs_2CuCl_4 .

PACS numbers: 73.43.Nq, 74.25.-q, 11.15.Ex

I. INTRODUCTION

Our understanding of states of matter and their internal orders has been dominated by Landau's symmetry breaking theory.[1, 2] We used to believe the all possible orders are described by various symmetry breaking states. However, after the discovery of fractional quantum Hall (FQH) states,[3, 4] we started to realize that FQH states contain a new kind of order - topological order[5] (which was first proposed to describe spin liquids found in research of high T_c superconductors[6]). What is new about topological orders is that topological orders cannot be characterized by symmetry breaking and the related local order parameters and long range correlations. Topological orders are characterized by new set of universal quantum numbers, such as ground state degeneracy, fractional statistics, edge excitations, *etc.*[5]

Recently, a concept of quantum order was proposed[7, 8] to describe non-symmetry breaking order that generally appear in a quantum state. The quantum order generalize the topological order to gapless states. To have a concrete description of quantum order without using order parameters, a new mathematical object - projective symmetry group (PSG) - was introduced. The concept of quantum order and its PSG characterization allow us to understand quantum phases and quantum phase transitions in a systematic way.

Just like the group theory allows us to classify symmetry breaking orders, the PSG characterization of quantum states allows us to classify different quantum orders[7] and can distinguish two different quantum phases even when they have exactly the same symmetry.

Also just like the symmetry description of classical order allows us to obtain low energy properties of system without knowing the details of the systems, the PSG de-

scription of quantum orders also allows us to obtain low energy properties of system without knowing the details of the systems. However, unlike symmetries which produce and protect gapless Nambu-Goldstone bosons,[9, 10] the quantum orders can produce and protect gapless collective modes which behave like light and other gapless gauge bosons.[11, 12] Those gapless collective modes can also be gapless/massless fermions, even when the original theory is purely bosonic.[13, 14, 15] More recently, it was realized that the quantum ordered states described by the PSG are actually string-net condensed states.[16] The emerging gauge bosons are the fluctuations of condensed string-nets and emerging fermions are the end of condensed strings.[17]

In this paper, we would like to further develop the PSG description of quantum orders. In Ref.[7], using PSG, a classification of quantum orders in symmetric spin liquids on 2D square lattice was given. Here, we would like to expand the results of Ref.[7] to 2D spin liquids which break the parities $P_x: x \rightarrow -x$ and $P_y: y \rightarrow -y$. We will assume the spin liquids have the following symmetries: two translation symmetries T_x, T_y , two parities $P_{xy}: (x, y) \rightarrow (y, x)$ and $P_{x\bar{y}}: (x, y) \rightarrow (-y, -x)$, and a time reversal symmetry. Such type of 2D spin liquids was observed recently in Cs_2CuCl_4 sample.[18, 19] By classifying the quantum orders in those spin liquids, we hope to identify the quantum order in the Cs_2CuCl_4 sample.

The spin dynamics in Cs_2CuCl_4 can be described by a 2D spin-1/2 system on a square lattice with nearest neighbor coupling $J' = 0.125\text{meV}$ and one diagonal coupling $J = 0.375\text{meV}$ in the $\hat{x} + \hat{y}$ direction. At temperature $T > T_c = 0.62\text{K} = 0.053\text{meV}$, the 2D spin system was found to be in a liquid state. Since T_c is a small energy scale, we will regard the finite temperature spin liquid state as a zero temperature quantum state. (More precisely, we assume that we can add additional frustrations to lower T_c to zero.) The theoretical spin-wave calculations[20, 21] and series expansion calculations[22] also suggest the existence of spin liquids in the above J - J' model. In this paper, we are going to concentrate on the

*Electronic address: yizhou@castu.tsinghua.edu.cn

†URL: <http://dao.mit.edu/~wen>

following physical issue: what is this spin liquid in the Cs_2CuCl_4 sample. There are several possibilities.

Since J' is small, one possibility is that the spin liquid, at low energies, just behaves like a decoupled 1D spin-1/2 chains. In other words, the spin liquids is in the same universality class of the decoupled 1D spin-1/2 chains. It is also possible the spin liquid is not purely 1D and has intrinsic 2D correlations. The 2D correlation can lead to 2D dispersion for low lying excitations. Furthermore, there can be several different types of 2D spin liquids that have the same symmetry as the purely 1D spin liquid. Since all those possible spin liquids have the same symmetry, it is difficult to study them without knowing how to characterize them.

In this paper, using quantum order and its PSG, we construct and characterize a large class of 2D spin liquids that have the same symmetry as the purely 1D spin liquid. In section II, we introduce mean-field ansatz that describe the symmetric spin liquids - the spin liquids that do not break any symmetries. In section III we discuss how to use PSG's to characterize different mean-field phases (or the universal classes of the mean-field ansatz). In section IV, we find all the Z_2 and $SU(2)$ PSG's and a large class of $U(1)$ PSG's within the $SU(2)$ slave-boson approach. This tells us the possible spin liquids that can be constructed use the $SU(2)$ slave-boson theory.

Constructing a large class of symmetric spin liquids and obtaining their PSG characterization are useful in the following sense. If a spin liquid state is found in Cs_2CuCl_4 or some other samples which do not break any symmetry, then the spin liquid has a good chance to be in the class that we obtained. Identifying the PSG that characterizes the constructed spin liquid will allow us to identify many universal properties of the spin liquid. We can check those universal properties experimentally which allow us to identify the PSGs for the experimentally observed spin liquids. The classification will also help us the study the phase transitions between symmetric spin liquids that do not change any symmetries.

In section V, we discuss some simple ansatz that realize some of the classified mean-field spin liquids. We calculate the mean-field energies of the constructed spin liquids to determine which spin liquids are likely to appear in the Cs_2CuCl_4 system. We also calculate spin correlation in those likely spin liquids in section VI. This allows experimentists to determine which spin liquid actually describe Cs_2CuCl_4 system (above T_c) using neutron scattering and other techniques. Section VII discusses connection of our results with experiments and previous theoretical results.

The main point of our calculation is to identify universal properties of various spin liquid states. This point is highly non-trivial since the involved spin liquids all have the same symmetry. We show that the crystal momenta of gapless spin-1 excitations can be used to experimentally distinguish different spin liquids.

We would like to pointed out that there are two ways to study possible spin liquid states in Cs_2CuCl_4 . The first

approach is the slave-fermion approach, where one start with a spin ordered state (such as the spiral state) and then study the spin liquid induced by strong spin wave fluctuations.[23, 24, 25] One can obtain the spin spectral function from the slave-fermion approach. The calculated spin spectral function[24, 25] agrees well with observed spin spectral function.[18, 19] In the slave-fermion approach, the spin disordered state (the spin liquid state) always has a gap, with low lying bosonic excitations.

In this paper, we are going to use slave-boson approach to study spin liquid states. In the slave-boson approach, the spin liquid state can either be gapped or gapless. The low lying excitations are fermions. In general slave-boson approach can generate more exotic states then the slave-fermion approach. We will see later that the spin spectral function obtained from the slave-boson approach also agrees well with observed spin spectral function.[18, 19]

However, the spin liquid obtained in Ref. [23, 24] and the spin liquids obtained in this paper are really different. For one thing, the spin liquid obtained in Ref. [23, 24] has a gap, while the spin liquids obtained in this paper is gapless. Because the gap is small, the two spin liquids behave similarly at finite temperatures.

What is the relation between the slave-boson approach and the slave-fermion approach? To understand the relation, we need use the result in Ref. [16] where it was shown that the spin liquids obtained from the slave-boson approach and the slave-fermion approach have a condensation of nets of closed strings. The quantum orders in the spin liquids obtained by both slave-fermion and slave-boson approaches can be described by PSG's. PSG's simply characterize different string-net condensations.

In the spin liquids obtained by slave-fermion approach, we concentrate on one type of condensed strings whose ends are bosons (we will call those strings bosonic strings). The PSG is nothing but the symmetry group of the hopping Hamiltonian of the ends of the condensed strings.[16] For different string condensations, the PSG for the ends of the condensed strings are different. Hence PSG can be used to characterize different string condensations (or different quantum orders).[16] In slave-fermion approach, we concentrate on spin liquids with condensation of fermionic strings. The ends of fermionic strings are fermions. We can use the PSG for the ends of fermionic strings to characterize different string condensed (or quantum ordered) states.

Some spin liquids contain condensation of both bosonic strings and fermionic strings. Those states can be characterized by either the PSG for the ends of the bosonic strings or the PSG for the ends of the fermionic strings.[16] Those states can be constructed either by slave-fermion approach or slave-boson approach.




FIG. 1: The nearest neighbor spin model on a triangular lattice.

II. SYMMETRIC SPIN LIQUIDS ON A SPIN-1/2 SYSTEM

We consider the spin-1/2 system on a 2D square lattice with nearest-neighbor coupling J' and next-nearest-neighbor coupling J on only one of the diagonal link $(i_x, i_y) \rightarrow (i_x + 1, i_y + 1)$. Such a lattice can also be regarded as a triangular lattice which does not have the 60° rotation symmetry. Within the $SU(2)$ slave-boson approach[26][27], general spin wave function can be constructed by introducing a mean-field Hamiltonian

$$H_{mean} = - \sum_{\langle i,j \rangle} \left(\psi_i^\dagger u_{ij} \psi_j + h.c. \right) + \sum_i a_0^l \psi_i^\dagger \tau^l \psi_i \quad (1)$$

where $\psi^T = (\psi_1, \psi_2)$, $u_{ji}^\dagger = u_{ij}$, $\tau^{1,2,3}$ are the Pauli matrices, and u_{ij} are 2×2 complex matrices. The collection $(u_{ij}, a_0^l \tau^l)$ is called a mean-field ansatz[7][8]. For each mean-field ansatz, we can obtain a mean-field ground state by filling the negative energy levels of H_{mean} with the spinons $\psi : |\Psi_{mean}^{(u_{ij}, a_0^l \tau^l)}\rangle$. The physical spin wave function Ψ_{spin} can now be obtained by performing a projection (see [7]):

$$\Psi_{spin}^{(u_{ij}, a_0^l \tau^l)}(\{r_i\}) = \langle 0 | \prod_{i=1}^{N_{up}} (\psi_{1,r_i} \psi_{2,r_i}) | \Psi_{mean}^{(u_{ij}, a_0^l \tau^l)} \rangle \quad (2)$$

where r_i is the coordinate of the i^{th} up-spin and N_{up} is the total number of the up-spin. Here an empty site of ψ represents a down-spin and a double occupied site represents an up-spin. Single occupied sites are unphysical states and are projected out.

Our representation does not have explicit spin rotation symmetry and it is hard to see spin rotation symmetry from the mean-field ansatz. This problem is solved in Ref. [7]. We find that a spin liquid is spin-rotation symmetric iff u_{ij} satisfies

$$\begin{aligned} u_{ij} &= i \rho_{ij} W_{ij} \\ \rho_{ij} &= \text{non-negative real number} \\ W_{ij} &\in SU(2) \end{aligned} \quad (3)$$

In this paper, we will only consider spin liquids with spin rotation symmetry, and we will assume that the ansatz satisfies the above conditions.

If we regard the ansatz $(u_{ij}, a_0^l \tau^l)$ as a label of physical wave function, $\Psi_{spin}^{(u_{ij}, a_0^l \tau^l)}(\{r_i\})$, then such a label is not a one-to-one label. It has a $SU(2)$ gauge structure. *i.e.* two ansatz, $(u_{ij}, a_0^l \tau^l)$ and $(W(u_{ij}), W(a_0^l \tau^l))$, related by a $SU(2)$ gauge transformation label the same physical wave function:

$$\begin{aligned} \Psi_{spin}(\{\alpha_i\}) &= \langle 0 | \prod_i f_{i\alpha_i} | \Psi_{mean}^{(W(u_{ij}), W(a_0^l \tau^l))} \rangle \\ &= \langle 0 | \prod_i f_{i\alpha_i} | \Psi_{mean}^{(u_{ij}, a_0^l \tau^l)} \rangle \end{aligned} \quad (4)$$

where $W(u_{ij}) = W_i u_{ij} W_j^\dagger$, $W(a_0^l(i) \tau^l) = W_i a_0^l(i) \tau^l W_i^\dagger$, and $W_i \in SU(2)$.

III. QUANTUM ORDER AND PROJECTIVE SYMMETRY GROUP

For the symmetry breaking states, the symmetry is a universal property shared by all the states in the same phase. Because of this, the symmetry provides a quantum number that characterizes different symmetry breaking orders. Hence we can say symmetry group (SG) describes the internal orders of symmetry breaking states. To characterize and classify different quantum orders which contain no symmetry breaking, we should construct some universal quantum numbers which describe different classes of quantum entanglement in the many-body ground state wave function. Ref. [7] and [8] propose that the symmetry of the mean-field ansatz $(u_{ij}, a_0^l \tau^l)$ is a universal property and serves as a quantum number that characterizes the quantum order in spin liquids. The symmetry of the group of the ansatz will be called the projective symmetry group (PSG). An element of PSG is a combined operation of a symmetry transformation followed by a gauge transformation. By definition, a PSG is formed by all the combined operations that leave the ansatz unchanged.

Because of the $SU(2)$ gauge structure, the $(u_{ij}, a_0^l \tau^l)$ labeling of the physical spin wave function $\Psi_{spin}(\{\alpha_i\})$ is not a one-to-one labeling. Two mean-field ansatzes differing by a $SU(2)$ gauge transformation give rise to the same spin wave function. Because of this, it is a non-trivial task to find out the symmetry of a spin liquid label by an ansatz $(u_{ij}, a_0^l \tau^l)$. In order for a spin wave function to have a symmetry, its corresponding ansatz is only required to be invariant under the symmetric transformation followed by a proper $SU(2)$ gauge transformation. Thus given two spin wave functions with the same symmetry, their ansatz can be invariant under the same symmetric transformations followed by *different* gauge transformations. In this case, the two spin liquids with the same symmetry can have different PSG's. We see

that the PSG characterization is more refined than the symmetry group characterization. PSG contains some information about the phase of spin wave function.

To understand the precise relation between PSG and SG, we need to introduce a subgroup of PSG - the invariant gauge group (IGG).[7][8] IGG is a special subgroup of PSG, which is formed by pure gauge transformations that leave the ansatz unchanged,

$$IGG \equiv \{W_i | W_i u_{ij} W_j^\dagger = u_{ij}, W_i \in SU(2)\} \quad (5)$$

Then SG and PSG are related by $SG = PSG/IGG$.

Spin liquids may support a special kind of low energy collective excitations - gauge fluctuations.[28] It was shown that[7, 29] the gauge group of those low energy gauge fluctuations is nothing but the IGG of the corresponding ansatz. We see that IGG of an ansatz is very important. In this paper, we will consider only three kinds of IGG, $SU(2)$, $U(1)$ and Z_2 . Correspondingly, we call the corresponding spin liquids $SU(2)$, $U(1)$ and Z_2 spin liquids.

Mathematically, PSG is defined as,

$$PSG \equiv \{G_U | G_U U(u_{ij}) = u_{ij}, G_U(i) \in SU(2)\} \quad (6)$$

where $U(u_{ij}) = \tilde{u}_{ij} \equiv u_{U(i),U(j)}$, $G_U U(\tilde{u}_{ij}) \equiv G_U(i) \tilde{u}_{ij} G_U^\dagger(j)$, U generates the symmetry transformation and G_U is the associated gauge transformation.

A PSG may change under a gauge transformation W . From $W G_U(u_{ij}) = W(u_{ij})$, where $W(u_{ij}) \equiv W_i u_{ij} W_j^\dagger$, we find that $W G_U U W^{-1} W(u_{ij}) = W(u_{ij})$. Therefore if $G_U U$ is in the PSG of ansatz u_{ij} , then $W G_U U W^{-1}$ is in the PSG of the gauge transformed ansatz $W(u_{ij})$. We see that the gauge transformation G_U associated with the transformation U changes in the following way

$$G_U(i) \rightarrow W(i) G_U(i) W(U(i))^\dagger \quad (7)$$

under a $SU(2)$ gauge transformation.

Since PSG is a property of an ansatz, we can group all ansatz sharing the same PSG together to form a class. Such a class is the universal class of quantum states that corresponds a quantum phase.

Now let us consider some simple examples of spin liquids described by the ansatz $(u_{ij}, a_0^l \tau^l)$. The first example is

$$u_{i,i+m} = u_m = u_m^l \tau^l \quad (8)$$

where u_m^l are real, $l = 1, 2, 3$. It is easy to obtain the spinon dispersion of such an ansatz,

$$E_\pm(k) = \pm \sqrt{\sum_l (u_m^l - a_0^l)^2}$$

where $u_k^l = \sum_m u_m^l e^{ik \cdot m}$. The Brillouin zone is $k_x, k_y \in (-\pi, \pi)$.

The second example is

$$\begin{aligned} u_{i,i+\hat{x}} &= i\chi\tau^0 + \eta\tau^3, \\ u_{i,i+\hat{y}} &= (-)^{i_x} (i\chi\tau^0 + \eta\tau^3), \\ u_{i,i+\hat{x}+\hat{y}} &= (-)^{i_x} \lambda\tau^1, \end{aligned} \quad (9)$$

where χ, η and λ are real. Its spinon spectrum is determined by

$$\begin{aligned} H(k) &= (\chi \sin k_x \tau^0 + \eta \cos k_x \tau^3) \otimes \tau^1 \\ &+ (\chi \sin k_y \tau^0 + \eta \cos k_y \tau^3) \otimes \tau^3 \\ &+ \lambda \tau^1 \otimes \tau^2 \sin(k_x + k_y) + a_0^l \tau^l \otimes \tau^0 \end{aligned}$$

where the Brillouin zone is $k_x \in (-\pi/2, \pi/2), k_y \in (-\pi, \pi)$. The four bands of spinon dispersion have a form of $\pm E_1(k), \pm E_2(k)$. It seems strange to find that the spinon spectrum is defined only on half of the lattice Brillouin zone. However, this is not inconsistent with the translation symmetry since the single spinon excitation is not physical. Only two-spinon excitations correspond to physical excitations and their spectrum should be defined on the full Brillouin zone. The two-spinon spectrum defined on the full Brillouin zone can be constructed from single-spinon spectrum[7]

$$\begin{aligned} E_{2s}(k) &= E_{\alpha_1}(k_1) + E_{\alpha_2}(k_2) \\ k &= k_1 + k_2 + n\pi\hat{x} \end{aligned} \quad (10)$$

where $n = 0, 1$ and $\alpha_{1,2} = 1, 2$ is the sub-index of the single spinon dispersion $\pm E_1(k), \pm E_2(k)$. We note that the physical spin-1 excitations are formed by two-spinon excitations in the mean-field theory. Thus the two-spinon spectrum E_{2s} is also the spectrum of physical spin-1 excitation and can be measured in experiments.

IV. CLASSIFICATION OF SYMMETRIC SPIN LIQUIDS

In this section, we classify symmetric spin liquids on triangular lattices. We consider the spin liquids which are invariant under translation transformation $T_x(i \rightarrow i + \hat{x})$ and $T_y(i \rightarrow i + \hat{y})$, parity transformation $P_{xy}((i_x, i_y) \rightarrow (i_y, i_x))$ and $P_{x\bar{y}}((i_x, i_y) \rightarrow (-i_y, -i_x))$, spin rotation transformation, spin-parity transformation $T^*(S_x \rightarrow S_x, S_y \rightarrow -S_y, S_z \rightarrow S_z)$. The spin rotation symmetry requires the ansatz take the form of Eq.(3). Under the spin-parity transformation T^* , the three components of spin change as $S_x \rightarrow S_x, S_y \rightarrow -S_y, S_z \rightarrow S_z$. This transformation can be realized through wavefunction transformation $\Phi \rightarrow \Phi^*$,

$$\langle \vec{S} \rangle = \frac{1}{2} \int \Phi^* f^\dagger \vec{\sigma} f \Phi \rightarrow \frac{1}{2} \int \Phi f^\dagger \vec{\sigma} f \Phi^*,$$

under the S_z representation f and f^\dagger are real, then

$$\langle \vec{S} \rangle = \langle \vec{S} \rangle^* = \frac{1}{2} \int \Phi^* f^\dagger \vec{\sigma} f \Phi$$

$$\begin{aligned}
& \rightarrow \frac{1}{2} \int \Phi f^\dagger \bar{\sigma} f \Phi^* = \left(\frac{1}{2} \int \Phi f^\dagger \bar{\sigma} f \Phi^* \right)^* \\
& = \frac{1}{2} \int \Phi^* f^\dagger \bar{\sigma}^* f \Phi,
\end{aligned}$$

which leads to $S_x \rightarrow S_x$, $S_y \rightarrow -S_y$, $S_z \rightarrow S_z$. The ansatz u_{ij} transfer as $u_{ij} \rightarrow u_{ij}^*$, combined with a gauge transformation $u_{ij} \rightarrow (i\tau^2) u_{ij} (-i\tau^2)$, we denote this combined transformation as

$$T^* : u_{ij} \rightarrow -u_{ij}. \quad (11)$$

Let us firstly discuss the method to classify the PSG briefly and leave the details to the appendix. For two given symmetry transformations, whose corresponding elements in a PSG, must satisfy some algebraic relations. Solving these equations allows us to construct a PSG which will be called the algebraic PSG. The new name algebraic PSG is introduced to distinguish them from the invariant PSG defined in the previous section. Any invariant PSG will be algebraic PSG. But an algebraic PSG may not be an invariant PSG unless there does exist an ansatz such that the algebraic PSG is the total symmetry group of the ansatz.

As an example, let us consider the two translations T_x and T_y , which satisfy the following relation

$$T_x T_y T_x^{-1} T_y^{-1} = 1. \quad (12)$$

From the definition of PSG, we find that the two elements of PSG, $G_x T_x$ and $G_y T_y$, must satisfy

$$\begin{aligned}
& G_x T_x G_y T_y (G_x T_x)^{-1} (G_y T_y)^{-1} \\
& = G_x T_x G_y T_y T_x^{-1} G_x^{-1} T_y^{-1} G_y^{-1} \\
& = G_x (i) G_y (i - \hat{x}) G_x^{-1} (i - \hat{y}) G_y^{-1} (i) \in \mathcal{G}. \quad (13)
\end{aligned}$$

Hereafter we will denote the IGG as \mathcal{G} . Each solution of the equation (13) corresponds to an algebraic PSG for T_x and T_y .

By adding other symmetry transformations, we can find and classify all the algebraic PSG associated with a given symmetry group. Since an invariant PSG is always an algebraic PSG, we can check whether the algebraic PSG is an invariant PSG through constructing an explicit ansatz u_{ij} . If an algebraic PSG supports an ansatz u_{ij} with no addition symmetry, then it is an invariant PSG. Through this method, we can classify the symmetric spin liquids through PSG.

In the appendix, we classify spin liquids with the above mentioned symmetries through PSG's. We limit ourselves to $SU(2)$, $U(1)$ and Z_2 spin liquids (we only consider those PSG's whose IGG is one of $SU(2)$, $U(1)$ and Z_2). We found that there are 63 kinds of Z_2 spin liquids, 30 kinds of $U(1)$ spin liquids and 2 kinds of $SU(2)$ spin liquids with $T_{x,y}$, $P_{xy,x\bar{y}}$, T^* and spin rotation symmetries.

V. APPLICATION TO NEAREST-NEIGHBOR SPIN COUPLING MODEL

In this section, we present our classification results via some simple examples. We will assume only $u_{i,i+x}$, $u_{i,i+y}$, and $u_{i,i+x+y}$ are no zero. In this case, we have the following 7 kinds of Z_2 spin liquids and 3 kinds of $U(1)$ (Other Z_2 and $U(1)$ spin liquids require non-vanishing u_{ij} on longer bonds.)

The 7 Z_2 spin liquids are:

(1) $Z2A\tau_+^0\tau_+^3$ spin liquid

$$\begin{aligned}
u_{i,i+\hat{x}} &= \chi\tau^1 + \eta\tau^2, \\
u_{i,i+\hat{y}} &= \chi\tau^1 + \eta\tau^2, \\
u_{i,i+\hat{x}+\hat{y}} &= \lambda\tau^1, \\
a_0^1 &= a_1, a_0^2 = a_2, a_0^3 = 0
\end{aligned} \quad (14)$$

(2) $Z2A\tau_+^1\tau_+^3$ spin liquid

$$\begin{aligned}
u_{i,i+\hat{x}} &= \chi\tau^1 + \eta\tau^2, \\
u_{i,i+\hat{y}} &= \chi\tau^1 - \eta\tau^2, \\
u_{i,i+\hat{x}+\hat{y}} &= \lambda\tau^1, \\
a_0^1 &= a_1, a_0^{2,3} = 0
\end{aligned} \quad (15)$$

(3) $Z2A\tau_-^0\tau_-^3$ spin liquid

$$\begin{aligned}
u_{i,i+\hat{x}} &= i\chi\tau^0 + \eta\tau^3, \\
u_{i,i+\hat{y}} &= i\chi\tau^0 + \eta\tau^3, \\
u_{i,i+\hat{x}+\hat{y}} &= \lambda\tau^1, \\
a_0^1 &= a_1, a_0^{2,3} = 0
\end{aligned} \quad (16)$$

(4) $Z2A\tau_-^1\tau_-^3$ spin liquid

$$\begin{aligned}
u_{i,i+\hat{x}} &= i\chi\tau^0 + \eta\tau^3, \\
u_{i,i+\hat{y}} &= i\chi\tau^0 - \eta\tau^3, \\
u_{i,i+\hat{x}+\hat{y}} &= \lambda\tau^1, \\
a_0^1 &= a_1, a_0^{2,3} = 0
\end{aligned} \quad (17)$$

(5) $Z2B\tau_-^3\tau_+^3$ spin liquid

$$\begin{aligned}
u_{i,i+\hat{x}} &= \chi\tau^1 + \eta\tau^2, \\
u_{i,i+\hat{y}} &= (-)^{i_x} (-\chi\tau^1 - \eta\tau^2), \\
u_{i,i+\hat{x}+\hat{y}} &= (-)^{i_x} \lambda\tau^1, \\
a_0^{1,2,3} &= 0
\end{aligned} \quad (18)$$

(6) $Z2B\tau_-^1\tau_+^3$ spin liquid

$$\begin{aligned}
u_{i,i+\hat{x}} &= \chi\tau^1 + \eta\tau^2, \\
u_{i,i+\hat{y}} &= (-)^{i_x} (\chi\tau^1 - \eta\tau^2), \\
u_{i,i+\hat{x}+\hat{y}} &= (-)^{i_x} \lambda\tau^2, \\
a_0^{1,2,3} &= 0
\end{aligned} \quad (19)$$

(7) $Z2B\tau^3\tau_-^1\tau_-^3$ spin liquid

$$\begin{aligned} u_{i,i+\hat{x}} &= i\chi\tau^0 + \eta\tau^3, \\ u_{i,i+\hat{y}} &= (-)^{i_x} (i\chi\tau^0 + \eta\tau^3), \\ u_{i,i+\hat{x}+\hat{y}} &= (-)^{i_x} \lambda\tau^1, \\ a_0^{1,2,3} &= 0. \end{aligned} \quad (20)$$

The 3 $U(1)$ spin liquids are:

(1) $U1A\tau^0\tau_+^0\tau_+^1$ spin liquids:

$$\begin{aligned} u_{i,i+\hat{x}} &= \chi\tau^3, \\ u_{i,i+\hat{y}} &= \chi\tau^3, \\ u_{i,i+\hat{x}+\hat{y}} &= \lambda\tau^3, \\ a_0^{1,2} &= 0, a_0^3 = a_3. \end{aligned} \quad (21)$$

(2) $U1B\tau^1\tau_-^0\tau_+^1$ spin liquids:

$$\begin{aligned} u_{i,i+\hat{x}} &= \chi\tau^3, \\ u_{i,i+\hat{y}} &= -(-)^{i_x} \chi\tau^3, \\ u_{i,i+\hat{x}+\hat{y}} &= (-)^{i_x} \lambda\tau^3, \\ a_0^{1,2,3} &= 0. \end{aligned} \quad (22)$$

(3) $U1C\tau_+^0\tau_-^0\tau_-^1$ spin liquids:

$$\begin{aligned} u_{i,i+\hat{x}} &= \chi\tau^2, \\ u_{i,i+\hat{y}} &= -\chi\tau^2, \\ u_{i,i+\hat{x}+\hat{y}} &= \lambda\tau^3, \\ a_0^{1,2} &= 0, a_0^3 = a_3. \end{aligned} \quad (23)$$

The labeling scheme of these spin liquids is defined in the appendix.

We have performed the self-consistent mean-field calculation for our J - J' model. We found many self-consistent mean-field solutions. The energies of four of them are plotted in Fig. 2, where we have set $J+J'=1$. To understand the physical properties of those mean-field states, we need to find their PSGs.

The A phase in Fig. 2 is a one-dimensional state. Its IGG is $SU^\infty(2)$, one $SU(2)$ for each decoupled 1D chain. We will call such a state $SU^\infty(2)$ spin liquid. Its ansatz is given by

$$u_{i,i+\hat{x}+\hat{y}} = \lambda\tau^3, a_0^{1,2,3} = 0, \quad (24)$$

with spinon dispersion

$$E(k) = \pm 2\lambda \cos(k_x + k_y) \quad (25)$$

This phase, favored by small J' , corresponds to the phase of decoupled 1D spin chain mentioned earlier.

The B phase is the $U1B\tau^1\tau_-^0\tau_+^1$ spin liquid. Its four spinon bands are given by

$$\pm 2\sqrt{\chi^2(\cos^2 k_x + \cos^2 k_y) + \lambda^2 \sin^2(k_x + k_y)} \quad (26)$$




FIG. 2: The mean-field energies for various phases in the nearest neighbor spin coupling system. (A) the $SU(2) \times SU(2)$ spin liquids in Eq.(24). (B) the $U1B\tau^1\tau_-^0\tau_+^1$ state in Eq.(22). (C) the $U1C\tau_+^0\tau_-^0\tau_-^1$ state in Eq.(23). (D) the π -flux phase ($SU2B$ state) in Eq.(28).




FIG. 3: (a) Contour plot of the spinon dispersion $E_+(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $U1B\tau^1\tau_-^0\tau_+^1$ state. (b) Lower edge of the two-spinon spectrum $E_{2s}(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $U1B\tau^1\tau_-^0\tau_+^1$ state.

The spinon dispersion $\min(E_1(k), E_2(k))$ and the lower edge of the spectrum $E_{2s}(k)$ of the physical spin-1 excitations are plotted as functions of $(k_x/\pi, k_y/\pi)$ in Fig. 3. This phase is one of 2D spin liquid phases.

The C phase has a PSG $U1C\tau_+^0\tau_-^0\tau_-^1$. The spinon dispersion is given by

$$E_+(k) = 2\sqrt{[\lambda \cos(k_x + k_y) - a_3]^2 + \chi^2 (\cos k_x - \cos k_y)^2} \quad (27)$$

The spinon dispersion $E_+(k)$ and the lower edge of the spin-1 spectrum $E_{2s}(k)$ are plotted as functions of $(k_x/\pi, k_y/\pi)$ in Fig. 4. This phase is another 2D spin liquid phase.

The D phase in Fig.2 is the π -flux phase ($SU2B$ spin




FIG. 4: (a) Contour plot of the spinon dispersion $E_+(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $U1C\tau_+^0\tau_-^0\tau^1$ state. (b) Lower edge of the two-spinon spectrum $E_{2s}(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $U1C\tau_+^0\tau_-^0\tau^1$ state.

liquid), which is given by

$$\begin{aligned} u_{i,i+\hat{x}} &= \chi\tau^1, \\ u_{i,i+\hat{y}} &= \chi\tau^2, \\ a_0^{1,2,3} &= 0. \end{aligned} \quad (28)$$

The spinon dispersion is

$$E_+(k) = 2\chi\sqrt{\cos^2 k_x + \cos^2 k_y} \quad (29)$$

The spinon dispersion $E_+(k)$ and the lower edge of the spin-1 spectrum $E_{2s}(k)$ are plotted as functions of $(k_x/\pi, k_y/\pi)$ in Fig. 5.




FIG. 5: (a) Contour plot of the spinon dispersion $E_+(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the π -flux state. (b) Lower edge of the two-spinon spectrum $E_{2s}(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the π -flux state.

Now we consider two other Z_2 spin liquids. The first one is the $Z2A\tau^0\tau_+^0\tau_+^3$ spin liquid state in Eq.(14). The energy dispersion of the $Z2A\tau^0\tau_+^0\tau_+^3$ state is given by

$$\begin{aligned} E_+(k) &= 2\sqrt{\epsilon_1^2 + \epsilon_2^2} \\ \epsilon_1 &= \chi(\cos k_x + \cos k_y) + \lambda \cos(k_x + k_y) - a_1 \\ \epsilon_2 &= \eta(\cos k_x + \cos k_y) \end{aligned} \quad (30)$$

The spinon dispersion $E_+(k)$ and the lower edge of the spin-1 spectrum $E_{2s}(k)$ are plotted as functions of $(k_x/\pi, k_y/\pi)$ in Fig. 6.




FIG. 6: ((a) Contour plot of the spinon dispersion $E_+(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $Z2A\tau^0\tau_+^0\tau_+^3$ state. (b) Lower edge of the two-spinon spectrum $E_{2s}(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $Z2A\tau^0\tau_+^0\tau_+^3$ state.

The second one is the $Z2A\tau^0\tau_+^0\tau_+^3$ spin liquid state in Eq.(15). Energy dispersion of the $Z2A\tau^0\tau_+^0\tau_+^3$ state is given by

$$\begin{aligned} E_+(k) &= 2\sqrt{\epsilon_1^2 + \epsilon_2^2} \\ \epsilon_1 &= \chi(\cos k_x + \cos k_y) + \lambda \cos(k_x + k_y) - a_1 \\ \epsilon_2 &= \eta(\cos k_x - \cos k_y) \end{aligned} \quad (31)$$

The spinon dispersion $E_+(k)$ and the lower edge of the spin-1 spectrum $E_{2s}(k)$ are plotted as functions of $(k_x/\pi, k_y/\pi)$ in Fig. 7.




FIG. 7: (a) Contour plot of the spinon dispersion $E_+(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $Z2A\tau^1\tau_+^1\tau_+^3$ state. (b) Lower edge of the two-spinon spectrum $E_{2s}(k)$ as a function of $(k_x/\pi, k_y/\pi)$ for the $Z2A\tau^1\tau_+^1\tau_+^3$ state.

If the mean-field state is stable against the gauge fluctuations, we expect that the mean-field spin-1 spectrum E_{2s} should qualitatively agree with the real spin-1 spectrum.

VI. SPIN SPECTRAL FUNCTION

The quantity of interest for comparison with experiments is the spin spectral function in (q, ω) space $S(q, \omega)$. $S(q, \omega)$ can be calculated from the Fourier transformation of two spin correlation function in real space

$\langle \vec{S}_i(t) \cdot \vec{S}_j(0) \rangle$. Due to the spin-rotation invariance, we obtain that

$$\langle \vec{S}_i \cdot \vec{S}_j \rangle = 3\langle S_i^z \cdot S_j^z \rangle = 3\langle S_i^x \cdot S_j^x \rangle = 3\langle S_i^y \cdot S_j^y \rangle$$

The spin spectral function at zero temperature is given by

$$S(q, \omega) = 3S^{zz}(q, \omega) \quad (32)$$

where

$$S^{zz}(q, \omega) = \sum_{\lambda} 2\pi |\langle \lambda | S^z(q) | G \rangle|^2 \delta(E_G + \omega - E_{\lambda}), \quad (33)$$

$$S^z(q) = \sum_j e^{iq \cdot j} S_j^z = \frac{1}{2} \left(\sum_{k, \alpha} \psi_{k+q, \alpha}^{\dagger} \psi_{k, \alpha} - N \delta_{q, 0} \right), \quad (34)$$

the summation is over all the eigenstates $|\lambda\rangle$ and N is the number of lattice sites.

Numerical calculation indicates that the neutron scattering spectrum predicted by the dynamics spin correlation contains continuous modes besides the main sharp peak. The Fig. 8-11 show the spin spectral function $S^{zz}(q, \omega)$ in the q -space. The point Γ is $(0, 0)$, M is $(0, \pi)$, and X is (π, π) . Darker shade represents a larger spectral function. These results can be compared with the inelastic neutron scattering experiments. The scan from Γ to X corresponds to the scan from $k = 0$ to $k = 2\pi/b$ in Fig. 2a of Ref. [18].




FIG. 8: The spin spectral function $S(q, \omega)$ of the $SU^{\infty}(2)$ spin liquids in Eq.(24).

VII. COMPARISON WITH EXPERIMENTS AND PREVIOUS THEORETICAL RESULTS

Our J - J' model is designed to describe the spin fluctuations in the Cs_2CuCl_4 sample. The spin dynamics in




FIG. 9: The spin spectral function $S(q, \omega)$ of the $U1B\tau^1\tau^0\tau^1_+$ state in Eq.(22). The solid lines mark the lower edge of the spectrum.




FIG. 10: The spin spectral function $S(q, \omega)$ of the $U1C\tau^0_+\tau^0_-\tau^1$ state in Eq.(23). The solid lines mark the lower edge of the spectrum.

Cs_2CuCl_4 at low temperatures were measured by neutron scattering.[18, 19] To the first order approximation, the measured spin spectral function is similar to that of decoupled 1D spin chain (or the $SU^{\infty}(2)$ spin liquid) (see Fig. 8). The decoupled 1D spin chain has gapless spin-1 excitations along the lines $k_x + k_y = \pi$ and $k_x + k_y = 2\pi$. If we examine the measured spectral function more closely to determine the low energy boundary of spin-1 excitation $\varepsilon_1(k_x, k_y)$, we find the gapless spin-1 excitations actually only appear at certain isolated points, implying that the spin state is 2-dimensional. The




FIG. 11: The spin spectral function $S(q, \omega)$ of the π -flux phase ($SU2B$ state) in Eq.(28)

experiment[18, 19] strongly suggests that there is a gapless spin-1 excitation at $(k_x, k_y) = (\pi/2 + \epsilon, \pi/2 + \epsilon)$ where $\epsilon \sim 0.04$. However, it is unclear whether the spin-1 excitation near $(k_x, k_y) = (\pi, \pi)$ is gapless or has a small gap. The experiment also determines the high energy boundary $\varepsilon_2(k_x, k_y)$ of spin-1 excitations. It was found that $\varepsilon_2(k_x, k_y)$ has maxima near $(k_x, k_y) = \pm(\pi/2, \pi/2)$ (the location is not known precisely).

Using this information, we can determine which spin liquids classified in this paper cannot describe the spin liquid state in Cs_2CuCl_4 . First the π -flux phase has no low energy spin-1 excitations near $(\pi/2, \pi/2)$. Thus it cannot describe the spin liquid state in Cs_2CuCl_4 . The $U1B\tau^1\tau^0\tau^1_+$ also does not work since the spin-1 excitations open a small gap near $(\pi/2, \pi/2)$.

It appears that the $U1C\tau^0_+\tau^0_-\tau^1$ spin liquid and some of its close relatives such as $Z2A\tau^0\tau^0_+\tau^3_+$, $Z2A\tau^1\tau^1_+\tau^3_+$, $Z2A\tau^1\tau^1_-\tau^3_-$ and $Z2A\tau^0\tau^0_-\tau^3_-$ spin liquids are consistent with observed spin spectral function. Let us list the universal characters of the above five spin liquids. The $U1C\tau^0_+\tau^0_-\tau^1$ and $Z2A\tau^0\tau^0_+\tau^3_+$ spin liquids only have gapless spin-1 excitations at $(0, 0)$, (π, π) , $\pm(\pi/2 + \epsilon, \pi/2 + \epsilon)$ and $\pm(\pi/2 - \epsilon, \pi/2 - \epsilon)$. The $Z2A\tau^1\tau^1_+\tau^3_+$ spin liquid only has gapless spin-1 excitations at $(0, 0)$, $\pm(\pi - \epsilon', \pi - \epsilon')$, $\pm(\pi/2 + \epsilon, \pi/2 + \epsilon)$ and $\pm(\pi/2 - \epsilon, \pi/2 - \epsilon)$. The $Z2A\tau^1\tau^1_-\tau^3_-$ and $Z2A\tau^0\tau^0_-\tau^3_-$ spin liquids only have gapless spin-1 excitations in small patches near $(0, 0)$, (π, π) , and $\pm(\pi/2, \pi/2)$. By measuring the locations of gapless spin-1 excitations and their spectral functions, we hope further experiments can determine which spin liquid actually describes the sample.

However, to compare the spin spectral function quantitatively with experimental measured results,[18, 19] we need to include fluctuations around the mean-field state. The interaction due to fluctuations enhances the spin fluctuations at $(\pi/2, \pi/2)$. The strong enough interac-

tion can drive the transition from the spin liquid state to the spiral state.

To include the interaction enhancement of the $(\pi/2, \pi/2)$ spin fluctuations, we consider the effects of the J -term (the diagonal coupling term): $\sum_i J \mathbf{S}_i \cdot \mathbf{S}_{i+\mathbf{x}+\mathbf{y}}$. Within the random phase approximation (RPA), the spin correlation function $\pi(\mathbf{k}, \omega)$ can be obtained from the mean-field spin correlation function $\pi_0(\mathbf{k}, \omega)$:

$$\pi(\mathbf{k}, \omega) = \frac{\pi_0(\mathbf{k}, \omega)}{1 - \tilde{J} \cos(k_x + k_y) \pi_0(\mathbf{k}, \omega)} \quad (35)$$

The imaginary part of $\pi(\mathbf{k}, \omega)$ gives us the RPA spin spectral function. To be definite, we start with the mean-field $U1C\tau^0_+\tau^0_-\tau^1$ spin liquid state and choose $\pi_0(\mathbf{k}, \omega)$ to be the mean-field spin correlation function of the $U1C\tau^0_+\tau^0_-\tau^1$ state. The experiments[18, 19] indicate that the system is close to the spiral phase. So here we choose the value of \tilde{J} to make the system close to the spiral phase instability. The resulting RPA spin spectral function is plotted in Fig. 12 and Fig. 13. The spin spectral functions along the lines A, C, and D in Fig. 12 were measured by experiments.[18] The simple RPA results are quite similar to the measured results (see Fig. 3 in Ref. [18]).




FIG. 12: The spin spectral function $S(q, \omega)$ of the $U1C\tau^0_+\tau^0_-\tau^1$ state obtained by RPA. The short solid lines mark the lower edge of the spectrum.

We would like to remark that the low energy effective theory for the $U1C\tau^0_+\tau^0_-\tau^1$ state contain gapless fermions interacting with $U(1)$ gauge field. Depending on that strength of the $U(1)$ gauge fluctuations, the $U(1)$ gauge interaction may or may not destabilize the $U1C\tau^0_+\tau^0_-\tau^1$ state. It is shown that if the spinons has a very anisotropic dispersion, the instanton effect of the 2+1 $U(1)$ gauge theory is irrelevant.[7] In this case, the $U1C\tau^0_+\tau^0_-\tau^1$ state is an algebraic spin liquid,[30] which contains gapless spin-1 excitations, but non of the gapless excitations are described by well defined quasiparticles. If the spinons has a more isotropic dispersion, the instanton effect is relevant, which lead to a confinement of the




FIG. 13: The RPA spin spectral function $S(q, \omega)$ along the lines A, C, D, G, and J in Fig. 12 for the $U1C\tau_+^0\tau_-^0\tau^1$ state.

$U(1)$ gauge field. In this case, the $U1C\tau_+^0\tau_-^0\tau^1$ state may become one of the stable Z_2 spin liquid states or other symmetry breaking states (such as the spiral state) below a transition temperature T_c .

Even when the $U1C\tau_+^0\tau_-^0\tau^1$ state is unstable, it is still useful for understanding the system at finite temperatures. The $U1C\tau_+^0\tau_-^0\tau^1$ state may control the dynamical properties of the Cs_2CuCl_4 samples above the T_c , provided that T_c is much smaller than the band width of the spin fluctuations.

VIII. CONCLUSION

In this paper, we use the projective symmetry group (PSG) to characterize quantum orders in symmetric spin liquids on 2D triangular lattice. We classify these symmetric spin liquids through PSG, and find that there are 63 kinds of Z_2 spin liquids, 30 kinds of $U(1)$ spin liquids and 2 kinds of $SU(2)$ spin liquids. The mean-field phase diagram Fig. 2. for the nearest neighbor spin coupling system is calculated. One-spinon and two-spinon excitation spectrum for some Z_2 and $U(1)$ spin liquids are inspected. We show that the gapless spinon-1 excitation spectrum can be used to physically measure the quantum order. By examining the measured spin spectral function in Cs_2CuCl_4 sample,[18] we find that it is possible that the spin liquid state in Cs_2CuCl_4 is described by the $U1C\tau_+^0\tau_-^0\tau^1$ spin liquid or one of its relatives.

We would like to point out that the RPA spin spectral function for the $U1C\tau_+^0\tau_-^0\tau^1$ spin liquid is very close to the spin spectral function[25] obtained from the slave-fermion approach.[23, 24] In the experiments,[18, 19] the spin liquid was observed only at a low but finite temperature. At that temperature, it is hard to distinguish to the two proposed states. However, if we find a spin liquid state at zero temperature, then we can tell which of the two proposed spin liquids can describe the experiments. This is because at zero temperature, the $U1C\tau_+^0\tau_-^0\tau^1$ spin liquid (and its relatives) and the spin liquid from the

slave-fermion approach are qualitatively different. The $U1C\tau_+^0\tau_-^0\tau^1$ spin liquid (and its relatives) contains gapless spin-1 excitations while the spin liquid from the slave-fermion approach is fully gapped.

XGW is supported in part by NSF Grant No. DMR-01-23156 and by NSF-MRSEC Grant No. DMR-02-13282.

APPENDIX A: CLASSIFICATION OF Z_2 SPIN LIQUIDS

In this appendix we consider the connected ansatz only, which is interesting and cover a wide range of spin liquids in fact. For any two sites i and j , a connected ansatz will offer a path $(i_1 i_2 \cdots i_n)$, which satisfy that $u_{i_k, i_{k+1}} \neq 0$, $k = 1, 2, \dots, n$, $i_1 = i$ and $i_n = j$. Due to the translation symmetry of the ansatz, we can choose a gauge in which all the loop operators of the ansatz are translation invariant. We will call such a gauge uniform gauge. Hereafter we will work in uniform gauge. According to the definition of PSG and uniform gauge,

$$\begin{aligned} G_x(i)P_{T_x C}G_x^\dagger(i) &= G_x(i)P_C G_x^\dagger(i) = P_C, \\ G_x(i)P_C &= P_C G_x(i). \end{aligned}$$

For Z_2 spin liquids, different loop operators basing at the same base point do not commute. We see that translation invariance of P_C in the uniform gauge requires that

$$G_x(i), G_y(i) \in \mathcal{G} = \{\pm\tau^0\}.$$

Gauge transformations $W(i) = (-)^{f(i)}\tau^0$ ($f(i) = \pm 1$) do not change the translation invariant property of the loop operators. We can always choose a gauge which satisfy that :

$$G_y(i) = \tau^0. \quad (\text{A1})$$

Gauge transformation $W(i) = W(i_x)$ does not change the condition $G_y(i) = \tau^0$, hence we can gauge fix that

$$G_x(i_x, i_y) = \tau^0. \quad (\text{A2})$$

From the relations (13), it is easy to see that

$$G_x(i)G_y(i - \hat{x})G_x^{-1}(i - \hat{y})G_y^{-1}(i) = \pm\tau^0. \quad (\text{A3})$$

Substituting Eqs. (A1) and (A2) into the above, one sees that

$$G_x(i)G_x^{-1}(i - \hat{y}) = \pm\tau^0. \quad (\text{A4})$$

Therefore there are two classes of PSG belong to translation symmetry,

$$G_x(i) = G_y(i) = \tau^0 \quad (\text{A5})$$

and

$$G_x(i) = (-)^{i_y}\tau^0, G_y(i) = \tau^0. \quad (\text{A6})$$

We will label the former $Z2A$ and the later $Z2B$.

Then we consider the spin parity symmetry T^* which satisfies that

$$T^* T_x T^{*-1} T_x^{-1} = T^* T_y T^{*-1} T_y^{-1} = T^{*2} = \pm 1. \quad (A7)$$

In a similar way, we have the following relations

$$\begin{aligned} G_{T^*} T^* G_x T_x (G_{T^*} T^*)^{-1} (G_x T_x)^{-1} &\in \mathcal{G}, \\ G_{T^*} T^* G_y T_y (G_{T^*} T^*)^{-1} (G_y T_y)^{-1} &\in \mathcal{G}, \\ G_{T^*} T^* G_{T^*} T^* &\in \mathcal{G}, \end{aligned} \quad (A8)$$

then

$$\begin{aligned} G_{T^*}(i) G_x(i) G_{T^*}^{-1}(i - \hat{x}) G_x^{-1}(i) &= \eta_{xt} \tau^0, \\ G_{T^*}(i) G_y(i) G_{T^*}^{-1}(i - \hat{y}) G_y^{-1}(i) &= \eta_{yt} \tau^0, \\ G_{T^*}(i)^2 &= \pm \tau^0, \end{aligned} \quad (A9)$$

$$G_{T^*}(i) = \eta_{xt}^{i_x} \eta_{yt}^{i_y} g_{T^*}, \quad (A10)$$

where $\eta_{xt} = \pm 1$, $\eta_{yt} = \pm 1$ and $g_{T^*}^2 = \pm \tau^0$. Two gauge inequivalent choices of g_{T^*} are $g_{T^*} = \tau^0$ and $g_{T^*} = i\tau^3$.

Next we add two types parity transformations $P_{xy}(i_x, i_y) = (i_y, i_x)$ and $P_{x\bar{y}}(i_x, i_y) = (-i_y, -i_x)$, which satisfy that

$$\begin{aligned} T_x P_{xy} T_y^{-1} P_{xy}^{-1} &= 1, \\ T_y P_{xy} T_x^{-1} P_{xy}^{-1} &= 1, \\ T_x P_{x\bar{y}} T_y P_{x\bar{y}} &= 1, \\ T_x P_{x\bar{y}} T_y P_{x\bar{y}} &= 1, \\ T^* P_{xy} T^{*-1} P_{xy}^{-1} &= 1, \\ T^* P_{x\bar{y}} T^{*-1} P_{x\bar{y}} &= 1, \end{aligned} \quad (A11)$$

From the above equations, we find that

$$\begin{aligned} (G_x T_x)(G_{P_{xy}} P_{xy})(G_y T_y)^{-1} (G_{P_{xy}} P_{xy})^{-1} &\in \mathcal{G} \\ (G_y T_y)(G_{P_{xy}} P_{xy})(G_x T_x)^{-1} (G_{P_{xy}} P_{xy})^{-1} &\in \mathcal{G} \\ (G_x T_x)(G_{P_{x\bar{y}}} P_{x\bar{y}})(G_y T_y)(G_{P_{x\bar{y}}} P_{x\bar{y}})^{-1} &\in \mathcal{G} \\ (G_y T_y)(G_{P_{x\bar{y}}} P_{x\bar{y}})(G_x T_x)(G_{P_{x\bar{y}}} P_{x\bar{y}})^{-1} &\in \mathcal{G} \\ G_{T^*} T^* (G_{P_{xy}} P_{xy})(G_{T^*} T^*)^{-1} (G_{P_{xy}} P_{xy})^{-1} &\in \mathcal{G} \\ G_{T^*} T^* (G_{P_{x\bar{y}}} P_{x\bar{y}})(G_{T^*} T^*)^{-1} (G_{P_{x\bar{y}}} P_{x\bar{y}})^{-1} &\in \mathcal{G} \end{aligned} \quad (A12)$$

and

$$\begin{aligned} G_x(i) G_{P_{xy}}(i - \hat{x}) G_y^{-1}(P_{xy}(i)) G_{P_{xy}}^{-1}(i) &= \eta_{pxy} \tau^0 \\ G_y(i) G_{P_{xy}}(i - \hat{y}) G_x^{-1}(P_{xy}(i)) G_{P_{xy}}^{-1}(i) &= \eta_{ypxy} \tau^0 \\ G_x(i) G_{P_{x\bar{y}}}(i - \hat{x}) G_y^{-1}(P_{x\bar{y}}(i) - \hat{x}) G_{P_{x\bar{y}}}^{-1}(i) &= \eta_{px\bar{y}} \tau^0 \\ G_y(i) G_{P_{x\bar{y}}}(i - \hat{y}) G_x^{-1}(P_{x\bar{y}}(i) - \hat{y}) G_{P_{x\bar{y}}}^{-1}(i) &= \eta_{ypx\bar{y}} \tau^0 \\ G_{T^*}(i) G_{P_{xy}}(i) G_{T^*}^{-1}(P_{xy}(i)) G_{P_{xy}}^{-1}(i) &= \eta_{tpxy} \tau^0 \\ G_{T^*}(i) G_{P_{x\bar{y}}}(i) G_{T^*}^{-1}(P_{x\bar{y}}(i)) G_{P_{x\bar{y}}}^{-1}(i) &= \eta_{tpx\bar{y}} \tau^0 \end{aligned} \quad (A13)$$

where $\eta_{pxy} = \pm 1$...

For $Z2A$ type PSG's,

$$\begin{aligned} G_{P_{xy}}(i) &= \eta_{xpxy}^{i_x} \eta_{ypxy}^{i_y} g_{P_{xy}}, \\ G_{P_{x\bar{y}}}(i) &= \eta_{xpx\bar{y}}^{i_x} \eta_{ypx\bar{y}}^{i_y} g_{P_{x\bar{y}}}. \end{aligned} \quad (A14)$$

For $Z2B$ type PSG's,

$$\begin{aligned} G_{P_{xy}}(i) &= (-)^{i_x i_y} \eta_{xpxy}^{i_x} \eta_{ypxy}^{i_y} g_{P_{xy}}, \\ G_{P_{x\bar{y}}}(i) &= (-)^{i_x i_y} \eta_{xpx\bar{y}}^{i_x} \eta_{ypx\bar{y}}^{i_y} g_{P_{x\bar{y}}}. \end{aligned} \quad (A15)$$

We also have that

$$P_{xy}^2 = P_{x\bar{y}}^2 = P_{xy} P_{x\bar{y}} P_{xy}^{-1} P_{x\bar{y}}^{-1} = 1 \quad (A16)$$

which leads to

$$\begin{aligned} G_{P_{xy}}(i) G_{P_{xy}}(P_{xy}(i)) &= \pm \tau^0, \\ G_{P_{x\bar{y}}}(i) G_{P_{x\bar{y}}}(P_{x\bar{y}}(i)) &= \pm \tau^0, \end{aligned}$$

and

$$\begin{aligned} (\eta_{xpxy} \eta_{ypxy})^{i_x + i_y} g_{P_{xy}}^2 &= \pm \tau^0, \\ (\eta_{xpx\bar{y}} \eta_{ypx\bar{y}})^{i_x + i_y} g_{P_{x\bar{y}}}^2 &= \pm \tau^0, \end{aligned}$$

The above implies that

$$\begin{aligned} \eta_{xpxy} &= \eta_{ypxy} = \eta_{pxy}, \\ \eta_{xpx\bar{y}} &= \eta_{ypx\bar{y}} = \eta_{px\bar{y}}, \\ g_{P_{xy}}^2 &= \pm \tau^0, g_{P_{x\bar{y}}}^2 = \pm \tau^0. \end{aligned}$$

By the similar way, one sees that

$$\eta_{xt} = \eta_{yt} = \eta_t. \quad (A17)$$

Gauge transformation $W(i) = (-)^{i_x}$ will change η_{pxy} to $-\eta_{pxy}$, therefore we can chose a gauge in which $\eta_{pxy} = 1$.

In conclusion, we have the following 60 kinds of $Z2A$ algebraic PSG's and 60 kinds of $Z2B$ algebraic PSG's on our triangular lattices:

$$Z2A g_{P_{xy}}(g_{P_{x\bar{y}}}) \eta_{px\bar{y}}(g_{T^*}) \eta_t$$

$$\begin{aligned} G_x(i) &= G_y(i) = \tau^0, \\ G_{T^*}(i) &= \eta_t^i g_{T^*}, \eta_t = \pm 1, \\ G_{P_{xy}}(i) &= g_{P_{xy}}, \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i g_{P_{x\bar{y}}}, \eta_{px\bar{y}} = \pm 1, \end{aligned} \quad (A18)$$

and

$$Z2B g_{P_{xy}}(g_{P_{x\bar{y}}}) \eta_{px\bar{y}}(g_{T^*}) \eta_t$$

$$\begin{aligned} G_y(i) &= \tau^0, G_x(i) = (-)^{i_y} \tau^0, \\ G_{T^*}(i) &= \eta_t^i g_{T^*}, \eta_t = \pm 1, \\ G_{P_{xy}}(i) &= (-)^{i_x i_y} g_{P_{xy}}, \\ G_{P_{x\bar{y}}}(i) &= (-)^{i_x i_y} \eta_{px\bar{y}}^i g_{P_{x\bar{y}}}, \eta_{px\bar{y}} = \pm 1, \end{aligned} \quad (A19)$$

where g_{T^*} , $g_{P_{xy}}$, and $g_{P_{x\bar{y}}}$ satisfy that

$$\begin{aligned}
g_{T^*}^2 &= \pm \tau^0, \\
g_{P_{xy}}^2 &= \pm \tau^0, \\
g_{P_{x\bar{y}}}^2 &= \pm \tau^0, \\
g_{T^*} g_{P_{xy}} g_{T^*}^{-1} g_{P_{xy}}^{-1} &= \pm \tau^0, \\
g_{T^*} g_{P_{x\bar{y}}} g_{T^*}^{-1} g_{P_{x\bar{y}}}^{-1} &= \pm \tau^0, \\
g_{P_{xy}} g_{P_{x\bar{y}}} g_{P_{xy}}^{-1} g_{P_{x\bar{y}}}^{-1} &= \pm \tau^0.
\end{aligned} \tag{A20}$$

All the gauge inequivalent g_{T^*} , $g_{P_{xy}}$, and $g_{P_{x\bar{y}}}$ are given by the following table

g_{T^*}	$g_{P_{xy}}$	$g_{P_{x\bar{y}}}$	$g_{P_{xy}} g_{P_{x\bar{y}}}$
τ^0	τ^0	τ^0	τ^0
τ^0	τ^0	$i\tau^3$	$i\tau^3$
τ^0	$i\tau^3$	τ^0	$i\tau^3$
τ^0	$i\tau^3$	$i\tau^3$	τ^0
τ^0	$i\tau^3$	$i\tau^1$	$i\tau^2$
$i\tau^3$	τ^0	τ^0	τ^0
$i\tau^3$	τ^0	$i\tau^3$	$i\tau^3$
$i\tau^3$	τ^0	$i\tau^1$	$i\tau^1$
$i\tau^3$	$i\tau^3$	τ^0	$i\tau^3$
$i\tau^3$	$i\tau^3$	$i\tau^3$	τ^0
$i\tau^3$	$i\tau^3$	$i\tau^1$	$i\tau^2$
$i\tau^3$	$i\tau^1$	τ^0	$i\tau^1$
$i\tau^3$	$i\tau^1$	$i\tau^3$	τ^2
$i\tau^3$	$i\tau^1$	$i\tau^1$	τ^0
$i\tau^3$	$i\tau^1$	$i\tau^2$	$i\tau^3$

After finding all the PSGs, next we need to find the ansatz that is invariant under those PSGs. This way we classify (mean-field) Z_2 spin liquids through PSG's. Let us first consider the translation symmetry. We have two classes of spin liquids,

Z2A spin liquids

$$\begin{aligned}
G_y(i) &= \tau^0, G_x(i) = \tau^0, \\
u_{i,i+m} &= u_m,
\end{aligned} \tag{A21}$$

and

Z2B spin liquids

$$\begin{aligned}
G_y(i) &= \tau^0, G_x(i) = (-)^{i_y} \tau^0, \\
u_{i,i+m} &= (-)^{m_y i_x} u_m.
\end{aligned} \tag{A22}$$

Considering the spin parity symmetry T^* , we have

$$G_{T^*} T^* (u_{i,i+m}) = u_{i,i+m}, \tag{A23}$$

which reads

$$- \eta_t^m g_{T^*} u_m g_{T^*}^{-1} = u_m \tag{A24}$$

Therefore, for different $(g_{T^*})_{\eta_t}$, we have the following table:

τ_{\pm}^0	$u_m = 0$
τ_{-}^0	$u_m = 0, \text{ if } m = \text{even}$
τ_{+}^3	$u_m = u_m^1 \tau^1 + u_m^2 \tau^2$
τ_{-}^3	$u_m = u_m^1 \tau^1 + u_m^2 \tau^2, \text{ if } m = \text{even}$ $u_m = u_m^0 \tau^0 + u_m^3 \tau^3, \text{ if } m = \text{odd}$

Now we consider the 180° rotation symmetry, which can be constructed through combining P_{xy} and $P_{x\bar{y}}$,

$$G_{P_{xy}} P_{xy} G_{P_{x\bar{y}}} P_{x\bar{y}} (u_{i,i+m}) = u_{i,i+m} \tag{A25}$$

the above leads to

$$\eta_{px\bar{y}}^m g_{P_{xy}} g_{P_{x\bar{y}}} u_m g_{P_{x\bar{y}}}^{-1} g_{P_{xy}}^{-1} = u_m^\dagger, \tag{A26}$$

for the *Z2A* spin liquids, and

$$(-)^{m_x m_y} \eta_{px\bar{y}}^m g_{P_{xy}} g_{P_{x\bar{y}}} u_m g_{P_{x\bar{y}}}^{-1} g_{P_{xy}}^{-1} = u_m^\dagger. \tag{A27}$$

for the *Z2B* spin liquids.

By writing all the ansatz, we find that although there are 120 kinds Z_2 algebraic PSG's, only 23 of them lead to *Z2A* spin liquids and 40 of them lead to *Z2B* spin liquids. Other algebraic PSG's lead to vanishing ansatz, $U(1)$ or $SU(2)$ spin liquids.

APPENDIX B: CLASSIFICATION OF $U(1)$ SPIN LIQUIDS

For $U(1)$ spin liquids, we can choose a gauge so that u_{ij} take the form

$$u_{ij} = i\rho_{ij} e^{i\theta_{ij}\tau^3} = u_{ij}^0 \tau^0 + u_{ij}^3 \tau^3 \tag{B1}$$

We will call this gauge canonical gauge. In the canonical gauge, IGG has a form of $\mathcal{G} = \{e^{i\theta\tau^3}, \theta \in [0, 2\pi)\}$, where θ_i is real for each i . Due to the translation symmetry of the ansatz, the loop operator P_C have a form

$$P_{T_x C} = (\tau^1)^{n_i} P_C (\tau^1)^{n_i} \tag{B2}$$

where $n_i = 0, 1$. The gauge transformation $G_{x,y}$ associated with the translation take the following form

$$\begin{aligned}
G_x(i) &= (-i\tau^1)^{n_i} e^{i\theta_x(i)\tau^3} (i\tau^1)^{n_{i-\hat{x}}} \\
G_y(i) &= (-i\tau^1)^{n_i} e^{i\theta_y(i)\tau^3} (i\tau^1)^{n_{i-\hat{y}}}
\end{aligned} \tag{B3}$$

in the canonical gauge. We note that a gauge transformations keep u_{ij} to have the form in the canonical gauge must have one of the following two forms

$$W_i = e^{i\theta_i \tau^3} \tag{B4}$$

$$W_i = e^{i\theta_i \tau^3} (i\tau^1) \tag{B5}$$

For spin liquids with connected u_{ij} , $G_{x,y}$ must take one of the above two forms in the canonical gauge. Thus n_i can only be one of the following four choices: $n_i = 0$, $n_i = (1 - (-)^i)/2$, $n_i = (1 - (-)^{i_x})/2$, and $n_i = (1 - (-)^{i_y})/2$. In these four cases, $G_{x,y}$ take one of the above two forms and u_{ij} can be connected.

Let us consider those cases in turn. We will work in the canonical gauge.

When $n_i = 0$, $G_{x,y}$ have a form

$$\begin{aligned} G_x(i) &= e^{i\theta_x(i)\tau^3} \\ G_y(i) &= e^{i\theta_y(i)\tau^3} \end{aligned} \quad (\text{B6})$$

by (B4) we can gauge fix them up to $\theta_x(i_x, 0) = 0$ and $\theta_y(i) = 0$. The relation (13) requires that

$$\theta_x(i) + \theta_y(i - \hat{x}) - \theta_x(i - \hat{y}) - \theta_y(i) = \varphi \quad (\text{B7})$$

Therefore $G_{x,y}$ have a form

$$\begin{aligned} G_x(i) &= e^{i(i_y\varphi + \theta_x)\tau^3} \\ G_y(i) &= e^{i\theta_y\tau^3} \end{aligned} \quad (\text{B8})$$

The ansatz with translation symmetry has a form

$$u_{i,i+m} = i\rho_m e^{i(-m_y i_x \varphi + \phi_m)\tau^3} \quad (\text{B9})$$

When $n_i = (1 - (-)^i)/2$,

$$\begin{aligned} G_x(i) &= e^{i\theta_x(i)\tau^3} (i\tau^1) \\ G_y(i) &= e^{i\theta_y(i)\tau^3} (i\tau^1) \end{aligned} \quad (\text{B10})$$

then

$$\begin{aligned} G_x(i) &= e^{i((-)^{i_y}\varphi + \theta_x)\tau^3} (i\tau^1) \\ G_y(i) &= e^{i\theta_y\tau^3} (i\tau^1) \end{aligned} \quad (\text{B11})$$

Using gauge transformation $W_i = e^{i((-)^{i_y-1}\varphi/2)\tau^3}$, the above change to

$$\begin{aligned} G_x(i) &= e^{i\theta_x\tau^3} (i\tau^1) \\ G_y(i) &= e^{i\theta_y\tau^3} (i\tau^1) \end{aligned} \quad (\text{B12})$$

The ansatz with translation symmetry has a form

$$u_{i,i+m} = i\rho_m e^{i(-)^i \phi_m \tau^3} \quad (\text{B13})$$

When $n_i = (1 - (-)^{i_x})/2$,

$$\begin{aligned} G_x(i) &= e^{i\theta_x(i)\tau^3} (i\tau^1) \\ G_y(i) &= e^{i\theta_y(i)\tau^3} \end{aligned} \quad (\text{B14})$$

then

$$\begin{aligned} G_x(i) &= e^{i(i_y\varphi + \theta_x)\tau^3} (i\tau^1) \\ G_y(i) &= e^{i\theta_y\tau^3} \end{aligned} \quad (\text{B15})$$

Using gauge transformation $W_i = e^{i(-)^{i_y}\varphi/2}\tau^3$, the above change to

$$\begin{aligned} G_x(i) &= e^{i\theta_x\tau^3} (i\tau^1) \\ G_y(i) &= e^{i\theta_y\tau^3} \end{aligned} \quad (\text{B16})$$

The ansatz with translation symmetry has a form

$$u_{i,i+m} = i\rho_m e^{i(-)^{i_x} \phi_m \tau^3} \quad (\text{B17})$$

When $n_i = (1 - (-)^{i_y})/2$,

$$\begin{aligned} G_x(i) &= e^{i\theta_x(i)\tau^3} \\ G_y(i) &= e^{i\theta_y(i)\tau^3} (i\tau^1) \end{aligned} \quad (\text{B18})$$

then

$$\begin{aligned} G_x(i) &= e^{i((-)^{i_y}\varphi + \theta_x)\tau^3} \\ G_y(i) &= e^{i\theta_y\tau^3} (i\tau^1) \end{aligned} \quad (\text{B19})$$

Using gauge transformation $W_i = e^{i((-)^{i_y-1}\varphi/2)\tau^3}$, the above change to

$$\begin{aligned} G_x(i) &= e^{i\theta_x\tau^3} \\ G_y(i) &= e^{i\theta_y\tau^3} (i\tau^1) \end{aligned} \quad (\text{B20})$$

The ansatz with translation symmetry has a form

$$u_{i,i+m} = i\rho_m e^{i(-)^{i_y} \phi_m \tau^3} \quad (\text{B21})$$

Now we add parity symmetry P_{xy} . Considering the flux in a plaquette, the parity symmetry requires $\varphi = 0$ or π in (B9). From the relations

$$\begin{aligned} G_x(i) G_{P_{xy}}(i - \hat{x}) G_y^{-1}(P_{xy}(i)) G_{P_{xy}}^{-1}(i) &= e^{i\theta_{xpy}\tau^3} \\ G_y(i) G_{P_{xy}}(i - \hat{y}) G_x^{-1}(P_{xy}(i)) G_{P_{xy}}^{-1}(i) &= e^{i\theta_{ypy}\tau^3} \\ G_x(i) G_{P_{x\bar{y}}}(i - \hat{x}) G_y^{-1}(P_{x\bar{y}}(i) - \hat{x}) G_{P_{x\bar{y}}}^{-1}(i) &= e^{i\theta_{xpx\bar{y}}\tau^3} \\ G_y(i) G_{P_{x\bar{y}}}(i - \hat{y}) G_x^{-1}(P_{x\bar{y}}(i) - \hat{y}) G_{P_{x\bar{y}}}^{-1}(i) &= e^{i\theta_{ypx\bar{y}}\tau^3} \\ G_{P_{xy}}(i) G_{P_{xy}}(P_{xy}(i)) &= e^{i\theta_{pxy}\tau^3} \\ G_{P_{x\bar{y}}}(i) G_{P_{x\bar{y}}}(P_{x\bar{y}}(i)) &= e^{i\theta_{px\bar{y}}\tau^3} \\ G_{P_{xy}}(i) G_{P_{x\bar{y}}}(P_{xy}(i)) G_{P_{xy}}^{-1}(P_{x\bar{y}}(i)) G_{P_{x\bar{y}}}^{-1}(i) &= e^{i\theta_p\tau^3} \end{aligned} \quad (\text{B22})$$

we will find only $n_i = 0$ and $n_i = (1 - (-)^i)/2$ lead to the existence of $G_{P_{xy}}(i)$. Then we can classify the $U(1)$ spin liquids with parity symmetry into three classes according to their translation property.

$U(1A)$ spin liquid:

$$\begin{aligned} G_x(i) &= e^{i\theta_x\tau^3}, \\ G_y(i) &= e^{i\theta_y\tau^3}, \\ u_{i,i+m} &= i\rho_m e^{i\phi_m\tau^3}. \end{aligned} \quad (\text{B23})$$

U1B spin liquid:

$$\begin{aligned} G_x(i) &= e^{i(i_y\pi+\theta_x)\tau^3}, \\ G_y(i) &= e^{i\theta_y\tau^3}, \\ u_{i,i+m} &= i(-)^{m_y i_x} \rho_m e^{i\phi_m\tau^3}. \end{aligned} \quad (\text{B24})$$

U1C spin liquid:

$$\begin{aligned} G_x(i) &= e^{i\theta_x\tau^3}(i\tau^1), \\ G_y(i) &= e^{i\theta_y\tau^3}(i\tau^1), \\ u_{i,i+m} &= i\rho_m e^{i(-)^i\phi_m\tau^3}. \end{aligned} \quad (\text{B25})$$

Next, we should find all the gauge inequivalent PSG's $G_{P_{xy}}(i)$ and $G_{P_{x\bar{y}}}(i)$ for the above cases respectively. In fact, as proved in Ref.[7], it exists the following relations between *U1A* and *U1B* PSG's:

$$\begin{aligned} \tilde{G}_{P_{xy}}(i) &= (-)^{i_x i_y} G_{P_{xy}}(i), \\ \tilde{G}_{P_{x\bar{y}}}(i) &= (-)^{i_x i_y} G_{P_{x\bar{y}}}(i), \\ \tilde{G}_{T^*}(i) &= G_{T^*}(i) \end{aligned} \quad (\text{B26})$$

where $G_{P_{xy}}, G_{P_{x\bar{y}}}$ are *U1A* PSG's. Hence we can consider the *U1A* PSG's only, and using the relations, we can construct the *U1B* PSG's.

1) *U1A* spin liquid, $G_x(i) = e^{i\theta_x\tau^3}$, $G_y(i) = e^{i\theta_y\tau^3}$, Eq.(B22) reads

$$G_{P_{xy}}(i) = e^{i((i_x-i_y)\varphi_{pxy}+\phi_{pxy})\tau^3} \quad (\text{B27})$$

$$\text{or } G_{P_{x\bar{y}}}(i) = e^{i((i_x+i_y)\varphi_{pxy}+\phi_{pxy})\tau^3}(i\tau^1), \quad (\text{B28})$$

and

$$G_{P_{x\bar{y}}}(i) = e^{i((i_x+i_y)\varphi_{px\bar{y}}+\phi_{px\bar{y}})\tau^3}$$

$$\text{or } G_{P_{xy}}(i) = e^{i((i_x-i_y)\varphi_{px\bar{y}}+\phi_{px\bar{y}})\tau^3}(i\tau^1)$$

Gauge transformation $W(i) = e^{-i(i_x\mp i_y)\varphi_{pxy}\tau^3/2}$ transfers (B27) and (B28) into

$$G_{P_{xy}}(i) = e^{i\phi_{pxy}\tau^3} \text{ or } G_{P_{x\bar{y}}}(i) = e^{i\phi_{pxy}\tau^3}(i\tau^1), \quad (\text{B29})$$

and do not change the forms of $G_{x,y,P_{x\bar{y}}}$. From the last equation of (B22), we obtain the following four cases,

a)

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}, \\ G_{P_{x\bar{y}}}(i) &= e^{i((i_x+i_y)\varphi_{px\bar{y}}+\phi_{px\bar{y}})\tau^3} \end{aligned}$$

Gauge transformation $W(i) = e^{-i(i_x+i_y)\varphi_{px\bar{y}}/2\tau^3}$ transfers the above into

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}, \\ G_{P_{x\bar{y}}}(i) &= e^{i\phi_{px\bar{y}}\tau^3} \end{aligned} \quad (\text{B30})$$

and does not change the forms of $G_{x,y}$.

b)

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}, \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i e^{i\phi_{px\bar{y}}\tau^3}(i\tau^1), \eta_{px\bar{y}} = \pm 1 \end{aligned} \quad (\text{B31})$$

c)

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}(i\tau^1), \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i e^{i\phi_{px\bar{y}}\tau^3}, \eta_{px\bar{y}} = \pm 1 \end{aligned} \quad (\text{B32})$$

d)

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}(i\tau^1), \\ G_{P_{x\bar{y}}}(i) &= e^{i((i_x-i_y)\varphi_{px\bar{y}}+\phi_{px\bar{y}})\tau^3}(i\tau^1) \end{aligned}$$

Gauge transformation $W(i) = e^{-i(i_x-i_y)\varphi_{px\bar{y}}/2\tau^3}$ transfers the above into

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}(i\tau^1), \\ G_{P_{x\bar{y}}}(i) &= e^{i\phi_{px\bar{y}}\tau^3}(i\tau^1) \end{aligned} \quad (\text{B33})$$

and does not change the forms of $G_{x,y}$.

Now we add the last symmetry spin parity T^* . For the same reason, $G_{T^*}(i)$ can take one of the two forms (B4) and (B5). The relations (A12) require that

$$\begin{aligned} G_{T^*}(i) G_x(i) G_{T^*}^{-1}(i-\hat{x}) G_x^{-1}(i) &= e^{i\theta_{xt}\tau^3} \\ G_{T^*}(i) G_y(i) G_{T^*}^{-1}(i-\hat{y}) G_y^{-1}(i) &= e^{i\theta_{yt}\tau^3} \\ G_{T^*}(i) G_{P_{xy}}(i) G_{T^*}^{-1}(P_{xy}(i)) G_{P_{xy}}^{-1}(i) &= e^{i\theta_{tpxy}\tau^3} \\ G_{T^*}(i) G_{P_{x\bar{y}}}(i) G_{T^*}^{-1}(P_{x\bar{y}}(i)) G_{P_{x\bar{y}}}^{-1}(i) &= e^{i\theta_{tpx\bar{y}}\tau^3} \\ G_{T^*}^2(i) &= e^{i\theta\tau^3}. \end{aligned} \quad (\text{B34})$$

Hence we have that all the *U1A* type PSG's:

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3} \\ G_{P_{x\bar{y}}}(i) &= e^{i\phi_{px\bar{y}}\tau^3} \\ G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_t^i e^{i\phi_t\tau^3}(i\tau^1) \end{aligned} \quad (\text{B35})$$

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3} \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i e^{i\phi_{px\bar{y}}\tau^3}(i\tau^1) \\ G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, e^{i\phi_t\tau^3}(i\tau^1) \end{aligned} \quad (\text{B36})$$

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}(i\tau^1) \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i e^{i\phi_{px\bar{y}}\tau^3} \\ G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, e^{i\phi_t\tau^3}(i\tau^1) \end{aligned} \quad (\text{B37})$$

$$\begin{aligned}
G_{P_{xy}}(i) &= e^{i\phi_{pxy}\tau^3}(i\tau^1) \\
G_{P_{x\bar{y}}}(i) &= e^{i\phi_{px\bar{y}}\tau^3}(i\tau^1) \\
G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_t^i e^{i\phi_t\tau^3}(i\tau^1)
\end{aligned} \tag{B38}$$

We can label these spin liquids as $U1Ag_{P_{xy}}(g_{P_{x\bar{y}}})_{\eta_{px\bar{y}}}(g_{T^*})_{\eta_t}$, where $g_{P_{xy}, P_{x\bar{y}}} = \tau^{0,1}$ corresponding to $e^{i\phi_{pxy}, px\bar{y}}\tau^3$ and $e^{i\phi_{pxy}, px\bar{y}}\tau^3(i\tau^1)$. Now we consider the form of ansatz that is invariant under the above PSG's. The translation symmetry requires that

$$u_{i,i+m} = u_m = u_m^0\tau^0 + u_m^3\tau^3. \tag{B39}$$

The 180° rotation symmetry requires that for $g_{P_{xy}}(g_{P_{x\bar{y}}})_{\eta_{px\bar{y}}} = \tau^0\tau_+^0, \tau^1\tau_+^1$,

$$u_m = u_{-m} = u_m^\dagger \tag{B40}$$

and for $g_{P_{xy}}(g_{P_{x\bar{y}}})_{\eta_{px\bar{y}}} = \tau^0\tau_{\eta_{px\bar{y}}}^1, \tau^1\tau_{\eta_{px\bar{y}}}^0$,

$$u_m = \eta_{px\bar{y}}^m \tau^1 u_{-m} \tau^1 = -\eta_{px\bar{y}}^m u_m \tag{B41}$$

The spin parity symmetry T^* requires that for $(g_{T^*})_{\eta_t} = \tau_-^0$

$$u_m = -(-)^m u_m \tag{B42}$$

and for $(g_{T^*})_{\eta_t} = \tau_{\eta_t}^1$

$$u_m = -\eta_t^m u_m^0\tau^0 + \eta_t^m u_m^3\tau^3 \tag{B43}$$

We find that the following 6 sets of ansatz that give rise to $U(1)$ symmetric spin liquids:

$$U1A[\tau_-^0\tau_+^1, \tau_-^1\tau_+^0]$$

$$\begin{aligned}
u_{i,i+m} &= u_m^0\tau^0 + u_m^3\tau^3 \\
u_m^{0,3} &= 0, \text{ if } m = \text{even}
\end{aligned} \tag{B44}$$

Here we have used the notation $U1A[ab, cd]e$ to represent the collection $U1Aabe$ and $U1Acde$.

$$U1A[\tau_+^0\tau_+^0, \tau_+^1\tau_+^1]$$

$$u_{i,i+m} = u_m^3\tau^3 \tag{B45}$$

$$U1A[\tau_+^0\tau_+^0, \tau_+^1\tau_+^1]$$

$$\begin{aligned}
u_{i,i+m} &= u_m^3\tau^3 \\
u_m^3 &= 0, \text{ if } m = \text{odd}
\end{aligned} \tag{B46}$$

The ansatz gives rise to $U(1) \times U(1)$ spin liquids since u_{ij} only connect points within two different sublattices.

Other PSG's lead to vanishing ansatz or $SU(2)$ spin liquids and can be dropped.

2) $U1B$ spin liquid, $G_x(i) = (-)^{i_y} e^{i\theta_x\tau^3}, G_y(i) = e^{i\theta_y\tau^3}$. From relations (B26), one can obtain all the $U1B$ PSG's easily,

$$\begin{aligned}
G_{P_{xy}}(i) &= (-)^{i_x i_y} e^{i\phi_{pxy}\tau^3}, \\
G_{P_{x\bar{y}}}(i) &= (-)^{i_x i_y} e^{i\phi_{px\bar{y}}\tau^3}, \\
G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_t^i e^{i\phi_t\tau^3}(i\tau^1)
\end{aligned} \tag{B47}$$

$$\begin{aligned}
G_{P_{xy}}(i) &= (-)^{i_x i_y} e^{i\phi_{pxy}\tau^3} \\
G_{P_{x\bar{y}}}(i) &= (-)^{i_x i_y} \eta_{px\bar{y}}^i e^{i\phi_{px\bar{y}}\tau^3}(i\tau^1) \\
G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, e^{i\phi_t\tau^3}(i\tau^1)
\end{aligned} \tag{B48}$$

$$\begin{aligned}
G_{P_{xy}}(i) &= (-)^{i_x i_y} e^{i\phi_{pxy}\tau^3}(i\tau^1) \\
G_{P_{x\bar{y}}}(i) &= (-)^{i_x i_y} \eta_{px\bar{y}}^i e^{i\phi_{px\bar{y}}\tau^3} \\
G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, e^{i\phi_t\tau^3}(i\tau^1)
\end{aligned} \tag{B49}$$

$$\begin{aligned}
G_{P_{xy}}(i) &= (-)^{i_x i_y} e^{i\phi_{pxy}\tau^3}(i\tau^1), \\
G_{P_{x\bar{y}}}(i) &= (-)^{i_x i_y} e^{i\phi_{px\bar{y}}\tau^3}(i\tau^1), \\
G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_t^i e^{i\phi_t\tau^3}(i\tau^1)
\end{aligned} \tag{B50}$$

We can label these spin liquids as $U1Bg_{P_{xy}}(g_{P_{x\bar{y}}})_{\eta_{px\bar{y}}}(g_{T^*})_{\eta_t}$. Now we consider the form of ansatz that is invariant under the above PSG's. The translation symmetry requires that

$$u_{i,i+m} = (-)^{m_y i_x} u_m = (-)^{m_y i_x} (u_m^0\tau^0 + u_m^3\tau^3). \tag{B51}$$

The 180° rotation symmetry requires that for $g_{P_{xy}}(g_{P_{x\bar{y}}})_{\eta_{px\bar{y}}} = \tau^0\tau_+^0, \tau^1\tau_+^1$,

$$u_m = (-)^{m_x m_y} u_m^\dagger \tag{B52}$$

and for $g_{P_{xy}}(g_{P_{x\bar{y}}})_{\eta_{px\bar{y}}} = \tau^0\tau_{\eta_{px\bar{y}}}^1, \tau^1\tau_{\eta_{px\bar{y}}}^0$,

$$u_m = -(-)^{m_x m_y} \eta_{px\bar{y}}^m u_m \tag{B53}$$

The spin parity symmetry T^* requires that for $(g_{T^*})_{\eta_t} = \tau_-^0$

$$u_m = -(-)^m u_m \tag{B54}$$

and for $(g_{T^*})_{\eta_t} = \tau_{\eta_t}^1$

$$u_m = -\eta_t^m u_m^0\tau^0 + \eta_t^m u_m^3\tau^3 \tag{B55}$$

We find that the following 10 sets of ansatz that give rise to $U(1)$ symmetric spin liquids:

$$U1B[\tau^0\tau_-^1, \tau^1\tau_-^0]\tau_-^0$$

$$\begin{aligned} u_{i,i+m} &= (-)^{m_y i_x} (u_m^0\tau^0 + u_m^3\tau^3) \\ u_m^{0,3} &= 0, \text{ if } m = \text{even} \end{aligned} \quad (\text{B56})$$

$$U1B[\tau^0\tau_+^0, \tau^1\tau_+^1]\tau_+^1$$

$$\begin{aligned} u_{i,i+m} &= (-)^{m_y i_x} u_m^3\tau^3 \\ u_m^3 &= 0, \text{ if } m_x = \text{odd and } m_y = \text{odd} \end{aligned} \quad (\text{B57})$$

$$U1B[\tau^0\tau_-^1, \tau^1\tau_-^0]\tau_-^1$$

$$\begin{aligned} u_{i,i+m} &= (-)^{m_y i_x} u_m^3\tau^3 \\ u_m^3 &= 0, \text{ if } m_x = \text{even and } m_y = \text{even} \end{aligned} \quad (\text{B58})$$

$$U1B[\tau^0\tau_+^0, \tau^1\tau_+^1]\tau_+^1$$

$$\begin{aligned} u_{i,i+m} &= (-)^{m_y i_x} u_m^3\tau^3 \\ u_m^3 &= 0, \text{ if } m_x = \text{odd or } m_y = \text{odd} \end{aligned} \quad (\text{B59})$$

The ansatz gives rise to $(U(1))^4$ spin liquids since u_{ij} only connect points within four different sublattices.

$$U1B[\tau^0\tau_-^1, \tau^1\tau_-^0]\tau_-^1$$

$$\begin{aligned} u_{i,i+m} &= (-)^{m_y i_x} (u_m^0\tau^0 + u_m^3\tau^3) \\ u_m^0 &= 0, \text{ if } m = \text{even} \\ u_m^3 &= 0, \text{ if } m_x = \text{even or } m_y = \text{even} \end{aligned} \quad (\text{B60})$$

Other PSG's lead to vanishing ansatz or $SU(2)$ spin liquids and can be dropped.

3) $U1C$ spin liquid, $G_x(i) = e^{i\theta_x\tau^3}(i\tau^1)$ and $G_y(i) = e^{i\theta_y\tau^3}(i\tau^1)$.

$$G_{P_{xy}}(i) = e^{i\theta_{pxy}(i)\tau^3}, e^{i\theta_{pxy}(i)\tau^3}(i\tau^1).$$

For the former four equations in (B22), we have that

$$\begin{aligned} \theta_{pxy}(i) &= (-)^i \varphi_{pxy} + \phi_{pxy}, \\ i_x\pi + (-)^i \varphi_{pxy} + \phi_{pxy} \end{aligned}$$

and the last three equations in (B22) require that

$$\begin{aligned} G_{P_{xy}}(i) &= \eta_{pxy}^i \eta_{xpxy}^{i_x} e^{i(\eta_{xpxy}\pi/4 + \phi_{pxy})\tau^3}, \\ &e^{i((-)^i \varphi_{pxy} + \phi_{pxy})\tau^3}(i\tau^1) \end{aligned} \quad (\text{B61})$$

and

$$\begin{aligned} G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i \eta_{xpx\bar{y}}^{i_x} e^{i(\eta_{xpx\bar{y}}\pi/4 + \phi_{px\bar{y}})\tau^3}, \\ &e^{i((-)^i \varphi_{px\bar{y}} + \phi_{px\bar{y}})\tau^3}(i\tau^1) \end{aligned} \quad (\text{B62})$$

where $\eta_{pxy,xpxy} = \pm 1, \eta_{px\bar{y},xpx\bar{y}} = \pm 1$. Now we add the last symmetry spin parity T^* . In a similar way, through the relations (A12), we have that

$$\begin{aligned} G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}, \\ &\eta_{xt}^{i_x} e^{i((-)^i \varphi_t + \phi_t)\tau^3}(i\tau^1) \end{aligned} \quad (\text{B63})$$

where $\eta_t = \pm 1, \eta_{xt} = \pm 1$.

It is noted that gauge transformation $W_i = e^{i(-)^i \phi\tau^3}$ and $W_i = \eta_{pxy}^{i_x}$ do not change $G_{x,y}$, and can be used to simplify the forms of $G_{P_{xy}, P_{x\bar{y}}, T^*}$. We conclude all the $U1C$ type PSG's:

$$\begin{aligned} G_{P_{xy}}(i) &= \eta_{xpxy}^{i_x} e^{i(\eta_{xpxy}\pi/4 + \phi_{pxy})\tau^3} \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i \eta_{xpx\bar{y}}^{i_x} e^{i(\eta_{xpx\bar{y}}\pi/4 + \phi_{px\bar{y}})\tau^3} \\ G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_{xpxy}^{i_x} e^{i\phi_t\tau^3}(i\tau^1) \end{aligned} \quad (\text{B64})$$

$$\begin{aligned} G_{P_{xy}}(i) &= \eta_{xpxy}^{i_x} e^{i(\eta_{xpxy}\pi/4 + \phi_{pxy})\tau^3} \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i \eta_{xpx\bar{y}}^{i_x} e^{i(\eta_{xpx\bar{y}}\pi/4 + \phi_{px\bar{y}})\tau^3}(i\tau^1) \\ G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_{xpxy}^{i_x} e^{i\phi_t\tau^3}(i\tau^1) \end{aligned} \quad (\text{B65})$$

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i(\eta_{xpxy}\pi/4 + \phi_{pxy})\tau^3}(i\tau^1) \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i \eta_{xpx\bar{y}}^{i_x} e^{i(\eta_{xpx\bar{y}}\pi/4 + \phi_{px\bar{y}})\tau^3} \\ G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_{xpxy}^{i_x} e^{i\phi_t\tau^3}(i\tau^1) \end{aligned} \quad (\text{B66})$$

$$\begin{aligned} G_{P_{xy}}(i) &= e^{i(\eta_{xpxy}\pi/4 + \phi_{pxy})\tau^3}(i\tau^1) \\ G_{P_{x\bar{y}}}(i) &= \eta_{px\bar{y}}^i \eta_{xpx\bar{y}}^{i_x} e^{i(\eta_{xpx\bar{y}}\pi/4 + \phi_{px\bar{y}})\tau^3}(i\tau^1) \\ G_{T^*}(i) &= \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}, \eta_{xpxy}^{i_x} e^{i\phi_t\tau^3}(i\tau^1) \end{aligned} \quad (\text{B67})$$

We can label these spin liquids as $U1C(g_{P_{xy}})_{\eta_{xpxy}}(g_{P_{x\bar{y}}})_{\eta_{px\bar{y}}}(g_{T^*})$. Now we consider the form of ansatz that is invariant under the above PSG's. The translation symmetry requires that

$$u_{i,i+m} = u_m^0\tau^0 + (-)^i u_m^3\tau^3. \quad (\text{B68})$$

The 180° rotation symmetry requires that for $G_{P_{xy}}(i) G_{P_{x\bar{y}}}(P_{xy}(i)) = \eta_{px\bar{y}}^i \tau^0$

$$u_m^0 = -\eta_{px\bar{y}}^m u_m^0, u_m^3 = (-\eta_{px\bar{y}})^m u_m^3; \quad (\text{B69})$$

for $G_{P_{xy}}(i) G_{P_{x\bar{y}}}(P_{xy}(i)) = \eta_{px\bar{y}}^i \eta_{xpxy}^{i_y}(i\tau^1)$

$$\begin{aligned} u_m^0 &= -\eta_{px\bar{y}}^m \eta_{xpxy}^{m_y} u_m^0, \\ u_m^3 &= -(-\eta_{px\bar{y}})^m \eta_{xpxy}^{m_y} u_m^3. \end{aligned} \quad (\text{B70})$$

The spin parity symmetry T^* requires that for $G_{T^*}(i) = \eta_t^i e^{i\phi_t\tau^3}|_{\eta_t=-1}$

$$u_{i,i+m} = -(-)^m u_{i,i+m} \quad (\text{B71})$$

and for $G_{T^*}(i) = \eta_{xpxy}^{i_x} e^{i\phi_t\tau^3}(i\tau^1)$

$$u_m^0 = -\eta_{xpxy}^{m_x} u_m^0, u_m^3 = \eta_{xpxy}^{m_x} u_m^3 \quad (\text{B72})$$

We find that the following 14 sets of ansatz that give rise to $U(1)$ symmetric spin liquids:

$$U1C[\tau_+^0\tau_-^0, \tau_-^0\tau_+^0, \tau_+^1\tau_-^1, \tau_-^1\tau_+^1]\tau^0$$

$$u_{i,i+m} = u_m^0\tau^0 + (-)^i u_m^3\tau^3$$

$$u_m^{0,3} = 0, \text{ if } m = \text{even}; \quad (\text{B73})$$

$$U1C[\tau_+^0\tau_+^0, \tau_+^1\tau_+^1]\tau^1$$

$$u_{i,i+m} = (-)^i u_m^3\tau^3$$

$$u_m^3 = 0, \text{ if } m = \text{odd} \quad (\text{B74})$$

The ansatz gives rise to $U(1) \times U(1)$ spin liquids.

$$U1C[\tau_+^0\tau_-^0, \tau_+^1\tau_-^1]\tau^1$$

$$u_{i,i+m} = (-)^i u_m^3\tau^3; \quad (\text{B75})$$

$$U1C[\tau_-^0\tau_+^0, \tau_-^1\tau_+^1]\tau^1$$

$$u_{i,i+m} = (-)^i u_m^3\tau^3$$

$$u_m^3 = 0, \text{ if } m_x = \text{odd or } m_y = \text{odd}; \quad (\text{B76})$$

The ansatz gives rise to $(U(1))^4$ spin liquids.

$$U1C[\tau_-^0\tau_-^0, \tau_-^1\tau_-^1]\tau^1$$

$$u_{i,i+m} = u_m^0\tau^0 + (-)^i u_m^3\tau^3$$

$$u_m^0 = 0, \text{ if } m_x = \text{even or } m_y = \text{odd}$$

$$u_m^3 = 0, \text{ if } m_x = \text{odd}; \quad (\text{B77})$$

$$U1C[\tau_-^0\tau_+^1, \tau_+^1\tau_-^0]\tau^1$$

$$u_{i,i+m} = u_m^0\tau^0 + (-)^i u_m^3\tau^3$$

$$u_m^0 = 0, \text{ if } m_x = \text{even}$$

$$u_m^3 = 0, \text{ if } m_x = \text{odd or } m_y = \text{even}; \quad (\text{B78})$$

Other PSG's lead to vanishing ansatz or $SU(2)$ spin liquids and can be dropped.

APPENDIX C: CLASSIFICATION OF $SU(2)$ SPIN LIQUIDS

We assume that for a $SU(2)$ spin liquid, one can always choose a gauge so that u_{ij} has a form

$$u_{ij} = u_{ij}^0\tau^0 \quad (\text{C1})$$

Hereafter we call this gauge canonical gauge. In the canonical gauge, IGG has a form $\mathcal{G} = SU(2)$. The gauge transformations that keep u_{ij} to have the form in the canonical gauge are given by

$$W_i = \eta(i)g \quad (\text{C2})$$

where $\eta(i) = \pm 1$ for each i and $g \in SU(2)$. The gauge transformation $G_{x,y}$ associated with the translation also take the above form

$$G_x(i) = \eta_x(i)g_x$$

$$G_y(i) = \eta_y(i)g_y \quad (\text{C3})$$

And we can gauge fix the above form as

$$G_x(i) = g_x$$

$$G_y(i) = \eta_y(i)g_y \quad (\text{C4})$$

where $\eta_y(i_x = 0, i_y) = 1$. Now the relation (13) reads

$$\eta_y(i - \hat{x})\eta_y(i) \in SU(2) \quad (\text{C5})$$

We find that there are only two different PSG's for translation symmetric ansatz

$$G_x(i) = g_x, G_y(i) = g_y \quad (\text{C6})$$

$$\text{and } G_x(i) = g_x, G_y(i) = (-)^{i_x} g_y \quad (\text{C7})$$

The two PSG's lead to the following two translation symmetric ansatz

$$u_{i,i+m} = u_m^0\tau^0 \quad (\text{C8})$$

$$\text{and } u_{i,i+m} = (-)^{m_x i_y} u_m^0\tau^0 \quad (\text{C9})$$

Similar to Z_2 case, adding more symmetries, we have all the $SU(2)$ PSG's belonged to triangular lattices.

$SU2A\eta_{px\bar{y}}\eta_t$

$$G_x(i) = g_x,$$

$$G_y(i) = g_y,$$

$$G_{T^*}(i) = \eta_t^i g_{T^*}, \quad (\text{C10})$$

$$G_{P_{xy}}(i) = g_{P_{xy}},$$

$$G_{P_{x\bar{y}}}(i) = \eta_{px\bar{y}}^i g_{P_{x\bar{y}}},$$

and

$SU2B\eta_{px\bar{y}}\eta_t$

$$G_x(i) = g_x,$$

$$G_y(i) = (-)^{i_x} g_y,$$

$$G_{T^*}(i) = \eta_t^i g_{T^*}, \quad (\text{C11})$$

$$G_{P_{xy}}(i) = (-)^{i_x i_y} g_{P_{xy}},$$

$$G_{P_{x\bar{y}}}(i) = (-)^{i_x i_y} \eta_{px\bar{y}}^i g_{P_{x\bar{y}}}$$

where $g_x, g_y, g_{T^*}, g_{P_{xy}}, g_{P_{x\bar{y}}} \in SU(2)$, $\eta_t = \pm 1$, and $\eta_{px\bar{y}} = \pm 1$. Moreover, only $\eta_t = -1$ and $\eta_{px\bar{y}} = -1$ lead to non vanishing ansatz. So there are only two kinds of $SU(2)$ spin liquids,

$SU2A$ spin liquids:

$$u_{i,i+m} = u_m^0\tau^0, m = \text{odd}, \quad (\text{C12})$$

$SU2B$ spin liquids:

$$u_{i,i+m} = (-)^{m_x i_y} u_m^0\tau^0, m = \text{odd}. \quad (\text{C13})$$

-
- [1] L. D. Landau, Phys. Zs. Sowjet **11**, 26 (1937).
 - [2] V. L. Ginzburg and L. D. Landau, J. exp. theor. Phys. **20**, 1064 (1950).
 - [3] D. C. Tsui, H. L. Stormer, and A. C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982).
 - [4] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
 - [5] X.-G. Wen, Advances in Physics **44**, 405 (1995).
 - [6] X.-G. Wen, Phys. Rev. B **40**, 7387 (1989).
 - [7] X.-G. Wen, Phys. Rev. B **65**, 165113 (2002).
 - [8] X.-G. Wen, Physics Letters A **300**, 175 (2002).
 - [9] Y. Nambu, Phys. Rev. Lett. **4**, 380 (1960).
 - [10] J. Goldstone, Nuovo Cimento **19**, 154 (1961).
 - [11] X.-G. Wen, Phys. Rev. Lett. **88**, 11602 (2002).
 - [12] O. I. Motrunich and T. Senthil, Phys. Rev. Lett. **89**, 277004 (2002).
 - [13] G. Kotliar and J. Liu, Phys. Rev. B **38**, 5142 (1988).
 - [14] L. Balents, M. P. A. Fisher, and C. Nayak, Int. J. Mod. Phys. B **12**, 1033 (1998).
 - [15] X.-G. Wen and A. Zee, Phys. Rev. B **66**, 235110 (2002).
 - [16] X.-G. Wen, hep-th p. 0302201 (2003).
 - [17] M. Levin and X.-G. Wen, cond-mat p. 0302460 (2003).
 - [18] R. Coldea, D. Tennant, A. Tsvelik, and Z. Tylczynski, Phys. Rev. Lett. **86**, 1335 (2001).
 - [19] R. Coldea, D. A. Tennant, K. Habicht, P. Smeibidl, C. Wolters, and Z. Tylczynski, Physical Review Letters **88**, 137203 (2002).
 - [20] A. E. Trumper, Phys. Rev. B **60**, 2987 (1999).
 - [21] J. Merino, R. H. McKenzie, J. B. Marston, and C. H. Chung, Journal of Physics: Condensed Matter **11**, 2965 (1999).
 - [22] W. Zheng, R. H. McKenzie, and R. R. P. Singh, Phys. Rev. B **59**, 14367 (1999).
 - [23] S. Sachdev, Phys. Rev. B **45**, 12377 (1992).
 - [24] C. H. Chung, J. B. Marston, and R. H. McKenzie, J. Phys.: Condens. Matter **13**, 5159 (2001).
 - [25] C.-H. Chung, K. Voelker, and Y. B. Kim, cond-mat/0302452 (2003).
 - [26] I. Affleck, Z. Zou, T. Hsu, and P. W. Anderson, Phys. Rev. B **38**, 745 (1988).
 - [27] E. Dagotto, E. Fradkin, and A. Moreo, Phys. Rev. B **38**, 2926 (1988).
 - [28] G. Baskaran and P. W. Anderson, Phys. Rev. B **37**, 580 (1988).
 - [29] X.-G. Wen, Phys. Rev. B **44**, 2664 (1991).
 - [30] W. Rantner and X.-G. Wen, Phys. Rev. B **66**, 144501 (2002).